



This work is protected by copyright and other intellectual property rights and duplication or sale of all or part is not permitted, except that material may be duplicated by you for research, private study, criticism/review or educational purposes. Electronic or print copies are for your own personal, non-commercial use and shall not be passed to any other individual. No quotation may be published without proper acknowledgement. For any other use, or to quote extensively from the work, permission must be obtained from the copyright holder/s.

**An epidemiological study of self-care restriction  
and joint pain in community-dwelling  
older people**

**Yeliz Prior**

**A thesis submitted for the degree of  
Doctor of Philosophy**

**June 2013**

**Arthritis Research UK Primary Care Centre  
Keele University**

## **Declaration**

This study involves secondary analysis of data collected in the North Staffordshire Osteoarthritis Project (NorStOP), which was funded by a Medical Research Council (MRC) programme grant (grant code: G9900220 and G0501798) and is hosted by the Arthritis Research UK Primary Care Centre, Keele University. The proposal for the PhD project was put forward by Dr Ross Wilkie, who obtained the Acorn funding for the studentship.

The NorStOP data were collected prior to the start of my PhD. Dr Ross Wilkie provided me with background information on the NorStOP study with regards to the methods of data collection and content of the surveys. I used the NorStOP data for secondary analysis of previously collected data. I planned and undertook all analyses for this study, with advice from the supervisory team: Dr Ross Wilkie, Dr Roger Beech and Professor George Peat. The supervisory team also advised me on the writing and presentations of the thesis chapters. I received guidance from Dr Rachel Gick on developing the search strategy for the systematic review. Dr Milisa Bucknall provided statistical advice throughout. The interpretation and the discussion of the findings are my own.

## **Abstract**

Self-care restriction is proposed in the World Health Organization's International Classification of Functioning (ICF) as an important functional consequence of health conditions. In contrast to traditional approaches which focus only on an individual's physical capacity, the ICF model also acknowledges the potential role of contextual factors such as environmental factors and personal needs. This thesis aims to understand the occurrence, course, and determinants of self-care restriction, as defined in the ICF, in community-dwelling adults aged 50 years and over, particularly in relation to joint pain and from a novel perspective: person-perceived participation restriction in which individuals judge whether their self-care needs are met "as and when I have wanted".

Data collected at three time points, by postal questionnaire across 6 years in the North Staffordshire Osteoarthritis project, were used in the analysis (n=7725 at baseline). The prevalence of person-perceived restricted self-care was 11.5% (95% CI 10.8, 12.2). A parsimonious model of factors associated with self-care restriction included age, activity limitation, depression, cognitive impairment, perceived inadequacy of income and low educational attainment. Older adults with joint pain were more likely to report self-care restriction, although this association was not independent of other factors. The frequencies of onset and persistence of restricted self-care at three years were 6.6% (95% CI 5.9, 7.5) and 38.9% (95% CI 33.7, 44.3) respectively. The key factors associated with restricted self-care at baseline were also predictive of the onset of self-care restriction at three years, but only depression was predictive of persistence.



Unmet need for help and assistance was associated with restricted self-care in older adults with joint pain. However, those who received help were also more likely to report restriction.

These analyses suggest a range of potential health and social targets for reducing self-care restriction in individuals with joint pain in middle and old age.

## **Acknowledgements**

Primarily, I would like to thank to my supervisors Dr Ross Wilkie, Dr Roger Beech and Professor George Peat for their invaluable advice and support throughout this study. I would like to thank to Dr Milisa Bucknall for her statistical advice and Professor Alison Hammond for her support and flexibility around my work commitments to allow me time to concentrate on writing my thesis.

I would like to extend my gratitude to Dr Sara Muller for her mentoring in the early stages of my PhD, and my fellow PhD candidates: Kate Watts for her compassion and understanding; Jemma Cowen for being there to listen; Andy Morden for his unfailing dry humour; and other research students for their friendship and peer support throughout the last four years. I would like to acknowledge my dear friend Zoe Mayson for being there for me in the good and the bad times, Ian Thomas for his unlimited good nature, and Evren Akkaya for being a true friend and inspiration. Furthermore, I am very grateful to the Arthritis Research UK, MRC and Acorn for funding this study and my studentship.

Finally, I would like to thank to my family for their support and encouragement, especially to my children, Max and Mia, for their continued patience, love and understanding over the last ten years. My utmost thanks go to my sister Filiz for being my rock, and a second mum to my kids when I worked in the holidays. Last, but not least, I would like to thank James Prior for taking the road less travelled with me and being his wonderful self; it made all the difference. Thank you.

## **Context of the thesis**

I studied for a BSc (Hons) in Occupational Therapy (OT) at the University of Salford. As part of my dissertation, I completed a research project proposal that spurred my interest in evidence based research. During this work, I recognised that there was a paucity of epidemiological research which underpinned the rationale behind the core principals of OT. I decided that I would like to contribute to my profession by getting actively involved in research. In order to do this, I needed to train to be a researcher. Thus, I started to search for funding opportunities to study for a research degree. Following the completion of my degree in 2008, I was successful in obtaining a PhD studentship at the Arthritis Research UK Primary Care Centre, Keele University to study for self-care restrictions and joint pain in older people. Self-care is an important topic in OT, as it is one of the three main occupations (i.e. self-care, productivity and leisure) in which therapists work together with individuals to overcome obstacles and optimise their involvement in activities of daily living.

After completing my PhD studentship at the Arthritis Research UK Primary Care Centre at Keele University, I secured a post as a research assistant to Professor Alison Hammond at the Centre of Health Sciences, University of Salford, where I continue to pursue research into the impact of musculoskeletal conditions further, and to build on the OT research in the assessment of activities of daily living and participation, through projects such as the UK-Evaluation of Daily Activity Questionnaire (UK-EDAQ), Valued Life Activities Scale (VLAS) and Work Rehabilitation in Inflammatory Arthritis (WORK-IA) .

## Contents

<b>Chapter 1 .....</b>	<b>20</b>
<b>Thesis statement and overview of research studies.....</b>	<b>20</b>
1.1 Introduction .....	20
1.2 Self-care .....	22
1.2.1 Epidemiology of self-care restriction .....	23
1.3 Thesis statement .....	25
1.4 Overview of research studies .....	26
1.4.1 Measuring the impact of musculoskeletal conditions in older adults: Models of disability [Chapter 2] .....	26
1.4.2 The impact of osteoarthritis on self-care in older adults: A systematic review [Chapter 3] .....	27
1.4.3 North Staffordshire Osteoarthritis Project (NorStOP): The survey overview and sample analysis [Chapter 4].....	28
1.4.4 Person-perceived self-care restriction in middle and old age: prevalence, distribution and associated factors [Chapter 5].....	29
1.4.5 Person-perceived self-care restriction in middle and old age: onset and persistence over a 3 year period [Chapter 6] .....	30
1.4.6 Potential predictors of the onset and persistence of restricted self-care in older adults with joint pain [Chapter 7].....	31
1.4.7 The concordance between activities and participation and the impact of environmental factors on self-care [Chapter 8] .....	33
1.4.8 Summary and discussion [Chapter 9] .....	35
<b>Chapter 2 .....</b>	<b>36</b>
2.1 Introduction .....	36
2.2 An overview of models of disability .....	38
2.2.1 The Nagi model of disablement (Nagi, 1965).....	38
2.2.2 The International Classification of Impairments, Disabilities, and Handicaps (ICIDH) .....	44
2.2.3 The International Classification of Functioning, Disability and Health (ICF) .....	46
2.2.4 Models of participation developed in Occupational Therapy .....	54

2.2.5	Overview of models of disability .....	56
2.3	The ICF model of self-care .....	62
<b>Chapter 3</b>	<b>.....</b>	<b>65</b>
<b>The impact of osteoarthritis on self-care in older adults: a systematic review .....</b>		<b>65</b>
3.1	The burden of musculoskeletal conditions in older adults.....	65
3.2	Osteoarthritis .....	65
3.2.1	Definition of osteoarthritis.....	66
3.2.2	Prevalence of osteoarthritis.....	69
3.2.3	The impact of osteoarthritis on self-care in older people.....	71
3.3	Aims and objectives .....	72
3.4	Methods .....	73
3.4.1	Search strategy .....	73
3.4.2	Selection criteria .....	75
3.4.3	Quality assessment and data extraction .....	77
3.4.4	Strength of the evidence .....	84
3.5	Results.....	85
3.5.1	Search Strategy .....	85
3.5.2	Articles Identified.....	89
3.5.3	Methodological quality .....	97
3.5.4	The association between self-care restriction and joint pain in community-dwelling older people.....	101
3.5.5	Determinants of self-care restrictions in older adults with joint pain... ..	103
3.5.6	Summary of results .....	103
3.6	Discussion .....	104
3.7	Conclusion .....	106
<b>Chapter 4</b>	<b>.....</b>	<b>109</b>
<b>North Staffordshire Osteoarthritis Project (NorStOP):</b>		
<b>The survey overview and sample analysis .....</b>		<b>109</b>
4.1	Introduction .....	109

4.2	Study design .....	110
4.2.1	Setting and sampling .....	110
4.2.2	Data administration .....	112
4.2.3	Ethical approval and informed consent .....	114
4.2.4	Follow-up of the NorStOP cohort .....	116
4.2.5	The Health and Regional Pains Survey Questionnaire Contents .....	117
4.2.6	Keele Assessment of Participation Restriction (KAP) .....	145
4.3	Response to NorStOP .....	155
4.3.1	Baseline survey response .....	155
4.3.2	Response at 3 years and 6 years follow-ups .....	157
4.3.3	Completeness of the data .....	159
4.4	The impact of non-response at baseline and attrition across the six years.. .....	160
4.4.1	Responders compared to non-responders at baseline .....	160
4.4.2	The NorStOP compared to the local and national population .....	161
4.5	Sample analysis.....	163
4.5.1	Descriptive characteristics of responders at baseline .....	163
4.5.2	Responders compared to lost at follow-up at 3 years .....	169
4.5.3	Responders compared to loss to follow-up at 6 years .....	171
4.6	Discussion .....	173
4.6.1	Representativeness of the general population .....	173
4.6.2	Study attrition .....	174
4.6.3	Data quality .....	174
4.6.4	Measurement of self-care restriction using KAP .....	176
4.7	Conclusions .....	178
<b>Chapter 5</b>	<b>.....</b>	<b>180</b>
<b>The prevalence and distribution of restricted self-care and associated factors in the general population at baseline.....</b>		<b>180</b>
5.1	Introduction .....	180
5.2	Aims and objectives .....	182
5.3	Methods .....	183

5.3.1	Dataset.....	183
5.3.2	Outcome .....	183
5.3.3	Covariates.....	183
5.4	Statistical analysis .....	185
5.4.1	Prevalence of person-perceived self-care restriction and its distribution by age and gender .....	185
5.4.2	Factors associated with restricted self-care in the general population.....	188
5.4.3	Parsimonious model of key factors associated with restricted self-care .....	193
5.4.4	Factors explaining the relationship between joint pain and self-care restriction.....	199
5.5	Results.....	200
5.5.1	Prevalence and age and gender distribution of restricted self-care ... ..	200
5.5.2	Factors associated with self-care restriction .....	202
5.5.3	Derivation of the parsimonious model of key factors associated with restricted self-care.....	213
5.5.4	Factors explaining the relationship between joint pain and self-care restriction.....	219
5.5.5	Summary of results .....	221
5.6	Discussion .....	222
5.6.1	Principal findings.....	222
5.6.2	Comparison with other studies.....	227
5.6.3	Limitations.....	231
5.7	Conclusions .....	234
<b>Chapter 6 .....</b>		<b>236</b>
<b>Person-perceived self-care restriction in the middle and old age: risk of onset and persistence over a 3 years period .....</b>		<b>236</b>
6.1	Introduction .....	236
6.2	Aims and objectives.....	243
6.3	Methods.....	244

6.3.1	Dataset.....	244
6.3.2	Outcome .....	244
6.3.3	Covariates.....	245
6.3.4	Statistical Analysis .....	247
6.4	Results.....	253
6.4.1	Onset self-care restriction at 3 years: complete-case analysis ....	253
6.4.2	Persistence of self-care restriction at 3 years .....	254
6.4.3	Association with baseline socio-economic and health characteristics .....	255
6.4.4	Association of the onset of self-care restriction with joint pain at baseline.....	257
6.4.5	Association of the persistence of self-care restriction with joint pain at baseline.....	259
6.4.6	Sensitivity analysis.....	262
6.4.7	Summary of results .....	262
6.5	Discussion .....	263
6.6	Conclusions .....	270
<b>Chapter 7</b>	<b>.....</b>	<b>272</b>
<b>Potential predictors of the risk of onset and persistence of restricted self-care in older people with joint pain ...272</b>		
7.1	Introduction .....	272
7.2	Aims and Objectives .....	276
7.3	Methods.....	277
7.3.1	Dataset.....	277
7.3.2	Outcome .....	277
7.3.3	Covariates.....	278
7.3.4	Statistical Analysis .....	281
7.4	Results.....	285
7.4.1	The risk of onset and persistence of self-care restriction at 3 years in those with joint pain.....	285
7.4.2	Association of the onset of self-care restriction at 3 years with joint specific characteristics at baseline .....	285



7.4.3	Factors associated with the onset of self-care restriction at 3 years ..	287
7.4.4	Association of the persistence of self-care restriction at 3 years with joint specific characteristics at baseline ..	289
7.4.5	Factors associated with the persistence of self-care restriction ...	290
	at 3 years ..	290
7.5	Summary of results ..	292
7.6	Discussion ..	293
7.6.1	Principal findings ..	293
7.6.2	Critical comparison against previous studies ..	293
7.6.3	Limitations ..	296
7.7	Conclusions ..	297
<b>Chapter 8</b>	<b>.....</b>	<b>299</b>
	<b>The concordance between activities and participation and the impact of environmental factors on self-care ..</b>	<b>299</b>
8.1	Introduction ..	299
8.2	Aims and Objectives ..	301
8.3	Methods ..	302
8.3.1	Dataset ..	302
8.3.2	Outcome ..	302
8.3.3	Covariates ..	303
8.3.4	Level of agreement between activities and participation ..	307
8.3.5	Statistical Analysis ..	309
8.4	Results ..	315
8.4.1	Comparison of person-perceived self-care restriction and task-specific activity limitation by age and gender ..	315
8.4.2	Self-care disability in the individual: examining the level of agreement between the constructs of activity and participation ..	320
8.4.3	The risk profile of person-perceived self-care restriction and task-specific activity limitation ..	326
8.4.4	Use of environmental facilitators ..	329
8.5	Summary of results ..	336
8.6	Discussion ..	337

8.6.1	Principal findings.....	337
8.6.2	Previous studies of activities and participation.....	343
8.6.3	Limitations.....	351
8.7	Conclusions .....	353
<b>Chapter 9</b>	<b>.....</b>	<b>355</b>
<b>Summary and discussion</b>	<b>.....</b>	<b>355</b>
9.1	Introduction .....	355
9.2	Summary of key findings.....	355
9.3	General discussion .....	359
9.3.1	The systematic review.....	359
9.3.2	The concept of person-perceived self-care restriction .....	365
9.3.3	The use of ICF as a framework.....	368
9.3.4	The use of the NorStOP dataset.....	370
9.4	Implications.....	372
9.4.1	Implications for practice (health and social care) .....	372
9.3.2	Implications for research .....	378
<b>References.....</b>		<b>381</b>

<b>List of abbreviations</b> .....	17
------------------------------------	----

## Figures

Figure 1.1	Expected population change in the United Kingdom, 2005-2051 .....	21
Figure 2.1	Nagi Model (Nagi, 1965) .....	39
Figure 2.2	The IOM Enabling-Disabling Process .....	41
Figure 2.3	The NCMRR Model.....	42
Figure 2.4	The ICIDH Model .....	44
Figure 2.5	The biopsychosocial model of health and illness .....	47
Figure 2.6	Diagram of interactions between the components of ICF .....	48
Figure 3.1	Flow diagram of papers accepted and rejected in the review.....	86
Figure 4.1	The pain manikin .....	125
Figure 4.2	Flowchart of response to NorStOP study at baseline.....	156
Figure 4.3	Study consort from baseline to 6 years follow-up.....	158
Figure 5.1	ICF Model of Human Function and Disability (WHO, 2001) .....	184

Figure 6.1	Flowchart of the onset and persistence of self-care restriction in the general population over a 3 years period.....	253
Figure 8.1	Comparison of age-related trends in prevalence of person-perceived self-care restriction (KAP restriction – solid bold black line) and number and severity of ADL limitations.....	319
Figure 8.2	The disability model illustrating relationship between contextual factors as scene-setters, contextual factors as facilitators and barriers, societal involvement, tasks, and acts.....	347

## Tables

Table 2.1	The ICF components and definitions.....	50
Table 2.2	The ICF concepts in context.....	51
Table 2.3	A critical review of the models of participation and disability .....	59
Table 2.4	The self-care and its components in the ICF framework .....	64
Table 3.1	The GDP diagnostic criteria for osteoarthritis.....	68
Table 3.2	Standardised prevalence rates of knee osteoarthritis.....	70
Table 3.3	Search strategy .....	75
Table 3.4	STROBE Statement—Checklist of items that should be included in reports of cross-sectional studies (an extract) .....	81
Table 3.5	The layout of the QUIPS-LBP: Quality Appraisal Exercise (an extract). .....	83
Table 3.6	Search 1: General population .....	87
Table 3.7	Search 2: Joint pain population.....	88
Table 3.8	Data extraction from the identified articles .....	94
Table 3.9	Overall summaries of the QUIPS-LBP ratings for each study .....	97
Table 3.10	Summary of key findings .....	108
Table 4.1	Health & Regional Pains Survey Questionnaire contents.....	118
Table 4.1b	Health & Regional Pains Survey Questionnaire contents .....	119
Table 4.1c	Health & Regional Pains Survey Questionnaire contents .....	120
Table 4.2	Categories of individual socio-economic classification .....	132
Table 4.3	Psychometric properties of the outcome measurement scales used in this study .....	135
Table 4.4	Strengths and weaknesses of the specifically developed single items in the survey.....	141
Table 4.5	A summary of the potential threats to the validity and reliability of self-reported survey data .....	144
Table 4.6	Keele Assessment of Participation .....	146
Table 4.7	Levels of missing data among items and scales within the NorStOP at baseline, 3 years and 6 years follow-up studies.....	159

Table 4.8	Age and gender distribution of respondents, exclusions, and non-responders of NorStOP Health Survey at baseline .....	160
Table 4.9	Comparison of the NorStOP responders to those of England and Staffordshire population by age, gender and ethnicity proportions .....	162
Table 4.10	Descriptive characteristics of NorStOP Health Survey baseline responders (n=7878): demographic and socio-economic characteristics .....	165
Table 4.11	Health characteristics of baseline responders to NorStOP Health Questionnaire (n=7878) .....	168
Table 4.12	Responders versus loss to follow-up at 3 years .....	170
Table 4.13	Responders versus loss to follow-up at 6 years .....	172
Table 4.14	Summary of key findings .....	179
Table 5.1	Age standardisation methods .....	186
Table 5.2	Variables included in the multivariable analysis .....	192
Table 5.3	The 4-week period prevalence of restricted self-care; overall and stratified by age and gender .....	200
Table 5.4	Standardisation of the prevalence of restricted self-care by age groups .....	202
Table 5.5	The standardised prevalence of restricted self-care .....	202
Table 5.6	Associations between restricted self-care and selected health conditions and impairments; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care .....	209
Table 5.7	Associations between restricted self-care and selected health conditions and impairments; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care in the univariable analysis .....	210
Table 5.8	Associations between restricted self-care and activity limitation; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care in the univariable analysis .....	211
Table 5.9	Associations between restricted self-care and contextual factors; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care in the univariable analysis .....	212
Table 5.10	Final (re-fitted) parsimonious model of key factors associated with restricted self-care in older adults in the general population .....	214
Table 5.11	Internal validity (NorStOP <sup>1</sup> ) of the parsimonious model .....	216
Table 5.12	External validity (NorStOP <sup>2</sup> ) scores of the parsimonious model .....	218
Table 5.13	Multivariable analysis of the associations of joint pain and restricted self-care adjusted for parsimonious factors .....	220
Table 5.14	Summary of key findings .....	235
Table 6.1	Index of Independence in Activities of Daily Living .....	238

Table 6.2	Onset of restricted self-care at 3 years; overall and stratified by age and gender .....	254
Table 6.3	Persistence of restricted self-care at 3 years; overall and stratified by age and gender.....	254
Table 6.4	Onset of self-care restriction at 3 years follow-up, by socio-economic status at baseline .....	255
Table 6.5	Distribution of the persistence of self-care restriction at 3 years by socio-demographic and health characteristics .....	256
Table 6.6	Onset of self-care restriction at 3 years and site of joint pain .....	258
Table 6.7	Onset of self-care restriction at 3 years and number of painful joint areas at baseline.....	258
Table 6.8	Onset of self-care restriction at 3 years and pattern of joint pain ....	259
Table 6.9	Persistence of self-care restriction at 3 years and site of joint pain .	260
Table 6.10	Persistence of self-care restriction at 3 years and number of painful joint areas at baseline .....	261
Table 6.11	Persistence of self-care restriction at 3 years and pattern of joint pain . at baseline .....	261
Table 6.12	Summary of key findings .....	271
Table 7.1	Joint-specific characteristics at baseline and onset of self-care restriction at 3 years.....	286
Table 7.2	Predicting the onset of self-care restriction at 3 years, multivariable analysis .....	288
Table 7.3	Joint-specific characteristics at baseline and persistence of self-care... restriction at 3 years.....	289
Table 7.4	Predicting the persistence of self-care restriction at 3 years, multivariable analysis.....	291
Table 7.5	Summary of key findings .....	298
Table 8.1	Key variables on self-care restriction, task-specific self-care activity limitation, and environmental facilitators/barriers and their mapping to ICF .....	305
Table 8.2	Other covariates used in this chapter .....	306
Table 8.3	Comparative occurrence and distribution by age and gender of self-care restriction and self-care activity limitation.....	317
Table 8.3b	Comparative occurrence and distribution by age and gender of self-care restriction and self-care activity limitation.....	318
Table 8.4	Level of agreement between person-perceived self-care restriction and task-specific activity limitation .....	325
Table 8.5	Comparison of risk profiles of person-perceived self-care restriction and task-specific activity limitation .....	328

Table 8.6	Cross-sectional association between restricted self-care and task-specific activity limitation in at least one self-care activity or more.....	330
Table 8.6b	Associations between restricted self-care and task-specific activity limitation; stratified by help and assistance status .....	330
Table 8.6c	Associations between restricted self-care and task-specific activity limitation; stratified by use of aids and appliances .....	331
Table 8.7	Socio-demographic characteristics of those who report limitation in washing oneself; and their help and assistance status .....	333
Table 8.8	Health characteristics of those who report limitation in washing oneself; and their help and assistance status .....	335
Table 8.9	Joint specific characteristics of those who report limitation in washing oneself; and their help and assistance status .....	336
Table 8.10	Distinguishing the components of functioning disability .....	348
Table 8.11	Summary of key findings.....	354
Table 9.1	Key findings from the thesis .....	358

## Appendices

Appendix 1: Baseline Health Questionnaire.....	447
Appendix 2: Health Questionnaire 3 years Follow-up .....	448
Appendix 3: Health Questionnaire 6 years Follow-up .....	449
Appendix 4: Baseline Regional Pains Questionnaire .....	450
Appendix 5: Regional Pains Questionnaire 3 years Follow-up .....	451
Appendix 6: Regional Pains Questionnaire 6 years Follow-up .....	452
Appendix 7: QUIPS-LBP Tool.....	453
Appendix 8: Published conference abstracts .....	454

## List of abbreviations

<b>ADL</b>	Activities of Daily Living
<b>AHEAD</b>	Asset and Health Dynamic Survey among the oldest old
<b>AIHW</b>	Australian Institute of Health and Welfare Knowledge
<b>AMED</b>	Allied and Alternative Medicine
<b>ASSIA</b>	Applied Social Sciences Index and Abstracts
<b>AOAT</b>	American Occupational Therapy Association
<b>AUSCAN</b>	Australian/Canadian Osteoarthritis Hand Index
<b>BAOT</b>	British Association of Occupational Therapists
<b>BPS</b>	British Pain Society
<b>CASP</b>	Critical Assessment Skills Programme
<b>CHPR</b>	Centre for Housing Planning and Research
<b>CI</b>	Confidence Interval
<b>CINAHL</b>	Cumulative Index to Nursing and Allied Health Literature
<b>CMOP</b>	Canadian Model of Occupational Performance
<b>COPM</b>	Canadian Occupational Performance Model
<b>COT</b>	College of Occupational Therapists
<b>EARRS</b>	Elderly At Risk Rating Scale
<b>EDAQ</b>	Evaluation of Daily Activity Questionnaire
<b>EMBASE</b>	Excerpta Medica Database
<b>ESCAP</b>	Economic and Social Commission for Asia and the Pacific
<b>FACS</b>	Fair Access to care Services
<b>FIM</b>	Functional Independence Measure
<b>GARS</b>	Groningen Activity Restriction Scale
<b>GMC</b>	General Medical Council
<b>GP</b>	General Practice
<b>HADs</b>	Hospital Anxiety and Depression Scale
<b>HAQ</b>	Health Assessment Questionnaire
<b>HRS</b>	Health Retirement Study
<b>HRQoL</b>	Health Related Quality of Life
<b>IADL</b>	Instrumental Activities of Daily Living
<b>ICD</b>	International Classification of Diseases
<b>ICE</b>	Imputations by Chained Equations
<b>ICF</b>	International Classification of Functioning
<b>ICIDH</b>	The International Classification of Impairments, Disabilities, and Handicaps
<b>ICR</b>	Intelligent Character Recognition
<b>IOM</b>	Institute of Medicine Model
<b>IPA</b>	Impact of Participation and Autonomy
<b>MCR</b>	Medical Components Summary
<b>MEDLINE</b>	Medical Literature Analysis and Retrieval System Online
<b>MESH</b>	Medical Subjects Headings
<b>MI</b>	Multiple Imputation
<b>MOHO</b>	Model of Human Occupation
<b>MOHOST</b>	The Model of Human Occupation Screening Tool
<b>MRC</b>	Medical Research Council

<b>NCMRR</b>	The National Centre for Medical Rehabilitation Research
<b>NIH</b>	National Institute of Health
<b>NHS</b>	National Health Service
<b>NORSTOP</b>	North Staffordshire Osteoarthritis Project
<b>NPSA</b>	National Patient Safety Agency
<b>NRES</b>	National Research Ethics Service
<b>NHSSTS</b>	National Health Service Strategic Tracing Service
<b>OCR</b>	Optic Character Recognition
<b>OMR</b>	Optic Mark Recognition
<b>ONS</b>	Office of National Statistics
<b>OT</b>	Occupational Therapy
<b>PsychINFO</b>	Psychological Information Database
<b>PCS</b>	Physical Components Summary
<b>PF</b>	Physical Functioning
<b>QUIPS</b>	Quality in Prognostic Studies
<b>QUORUM</b>	The Quality of Reporting of Meta-analyses
<b>RCT</b>	Randomised Clinical Trial
<b>REC</b>	Research Ethics Committee
<b>RNL</b>	Reintegration to Normal Living Index
<b>RoC</b>	Receiver Operating Characteristics
<b>SEM</b>	Structural Equation Modelling
<b>SF</b>	Short Form
<b>SIP</b>	Sickness Impact Profile
<b>STROBE</b>	Strengthening the Reporting of Observational Studies in Epidemiology
<b>TIMPs</b>	Tissue inhibitors of metalloproteinases
<b>TJR</b>	Tayside Joint Replacement
<b>UK</b>	United Kingdom
<b>VAS</b>	Visual Analogue Scale
<b>VLAS</b>	Valued Life Activities Scale
<b>WHO</b>	World Health Organisation
<b>WOMAC</b>	Western Ontario and McMaster Universities Osteoarthritis Index




# **Chapter 1**

## **Thesis statement and overview of research studies**

### **1.1 Introduction**

Life expectancy is increasing with older adults in the United Kingdom (UK) expected to live until they are 80 years old (Office for National Statistics (ONS), 2010); eight years longer than in the 1970s (Leon, 2007). The number of adults aged 50 years and over has increased over the last two decades, particularly those aged 80 and over, which now represents 4.5 percent of the UK population (ONS, 2010). The proportion and number of adults aged 50 and over is expected to increase even further over the forthcoming years (Figure 1.1). This will have considerable impact on future health and social care provision (Medical Research Council (MRC) 2010) because with increasing life expectancy comes an increased prevalence of chronic health conditions (Denton and Spencer, 2010), disability (Berlov et al., 2012; Martin et al., 2010; MRC 2010; Nusselder et al., 2006) and a reduction in Health Related Quality of Life (HRQoL) (e.g. HRQoL is four times lower in adults aged 75 years and over compared to those aged 30-44 years (Saarni et al., 2007)).

**Figure 1.1 Expected population change in the United Kingdom, 2005-2051**



**Source:** ONS, Population estimates 2008

Musculoskeletal conditions (such as osteoarthritis and back pain) are the most frequent morbidity in older adults (World Health Organisation (WHO), 2003). In the UK, pain, the most common symptom of musculoskeletal conditions, affects two thirds of adults aged 50 years and over (Thomas et al., 2004:a) and is often not controlled by medical treatment (British Pain Society, 2004). As a consequence, musculoskeletal pain often becomes chronic (i.e. lasts for more than three months), significantly reduces people's ability to participate in daily activities, and is strongly related to further comorbidities (e.g. anxiety and depression) (Cherubino et al., 2012; Breivik et al., 2006). However the consequences of musculoskeletal conditions are extensive and not completely understood (Woolf and Pfleger, 2003). Many studies have focused on physical function, but greater acknowledgement of the wider impact of musculoskeletal conditions (for example,

the impact on social roles and essential activities of daily life) is required to guide strategies and management to reduce the impact of musculoskeletal conditions in the growing population of older adults. To facilitate this there is a need for epidemiological studies to enhance the understanding of the frequency, incidence, causes and progression of the different forms of impact (Woolf and Pfleger, 2003).

## **1.2 Self-care**

The daily lives of adults are built upon routines and habits that reflect their roles, needs, motivators and responsibilities (Turner et al., 2002). The ability to participate in self-care activities such as bathing and toileting is important to older people, as they are an essential part of daily life. Losing the ability to carry out self-care tasks is directly associated with losing independence (Jagger et al., 2001). The ability to function independently is an important aspect of quality of life (Harwood et al., 2004; Spector et al., 1987). Being dependent on others later in life is one of the biggest fears of older people (Gignac et al., 2000) and is associated with poor health and wellbeing (Covinsky et al., 2008; Dawson et al., 2004; Dunlop et al., 2001; Hosie and Dickson, 2000; Gill and Feinstein, 1994). Limitations in self-care activities predict further morbidity and mortality in older adults (Gill et al., 2004 & 2002; Naik et al., 2004; Jagger et al., 2001; Czaja et al., 1993).

### **1.2.1 Epidemiology of self-care restriction**

Studies reporting the frequency of self-care have tended to measure specific tasks as part of a wider measure of activities of daily living (ADL) or instrumental activities of daily life (IADL). Although ADL limitations are commonly assessed in clinical research, measurement of ADL varies across studies and can include items referring to mobility, transfers (e.g. getting up from a chair) and self-care (e.g. washing, dressing, bathing, toileting). IADL involves more complex tasks such as meal preparation, grocery shopping and using the telephone.

Even though the application of different forms and definitions of self-care restriction in the literature results in varied estimates, there is some consensus that self-care restriction is frequent in older adults. Whilst mobility limitation is the most commonly reported disability amongst the older people, 70% of adults with mobility problems are also restricted in specific self-care tasks (i.e. they were restricted in at least one of the following tasks; dressing, bathing or showering, eating, getting in and out of bed or using the toilet; Melzer et al, 2005). The prevalence of ADL disability in older people ranges from 5-40% depending on how it is defined and measured (Balzi et al., 2010; Gill et al., 2004:a-b and 2002:a; Jagger et al., 2001; Dunlop et al., 1997). The most common ADL disability is bathing disability and is estimated to affect 24% to 34% of older adults (Gill et al., 2006: a; Naik et al., 2004; Jagger et al., 2001). Gill et al. (2002:a) suggested that the occurrence of ADL disability is substantially underestimated because it fluctuates; frequent assessments could provide an improved understanding of the course of disability in community-living older persons (Gill, et al., 2002:a and

2004). In a longitudinal study of 754 community-dwelling older adults aged >70, it was reported that the burden of bathing disability increased with age, was greater in women and in participants who were physically frail (Gill et al, 2006). This study also reported that the onset of bathing disability was associated with a fivefold increase in the likelihood of developing disability in the other essential ADL in the following month (Gill et al, 2006).

Importantly self-care restriction has been found to predict future morbidity and mortality in older adults (Czaja et al., 1993; Spector et al., 1987). Jagger et al. (2001) investigated the patterns of the onset of different types of disability and the association with age and gender. Disability was measured by self-report of performance in mobility around home, getting to and from the toilet, transfer from bed, feeding, dressing and bathing, and defined as being independent but having difficulty, using aids or help, or being unable to perform in any one of these ADL. The overall prevalence of ADL disability was 48% and it occurred in the following order; bathing, mobility, toileting, dressing, transfer from bed or chair and feeding. Women were identified at a higher risk of disability in bathing (relative risk (RR) 1.6; 95% Confidence Interval (CI) 1.3, 1.9;  $p < 0.001$ ) and toileting (RR 1.7; 95% CI 1.2, 2.5;  $p = 0.003$ ) and for all ADL there was a significant increase for risk of disability with increasing age. However the order of onset of disability for ADL was invariant across age and genders (Jagger et al., 2001). Previously, Dunlop and colleagues (1997) reported a similar order of disability to Jagger (i.e. based on both the ordering of median ages to disability onset and patterns of incident disability, dependency in ADLs were ordered as walking, bathing, transferring,

dressing, toileting, feeding). This study also found gender differences in disability incidence rates, suggesting that although women live longer than men, they spend more time in a disabled state (Dunlop et al., 1997).

Previous research has characterised self-care in terms of an individual's capacity to do specific tasks such as bathing or dressing. However, self-care also involves environmental factors (e.g. bathing entails getting in and out the bath, operating taps and washing oneself; these tasks are influenced by pain, joint stiffness, comorbidities and environmental factors such as access to the bath (i.e. upstairs, downstairs, in a small bathroom difficult to conduct transfers), and the availability of aids and adaptations (i.e. does it have grab rails or is there a bath seat/board/non-slip mat in place; powered bath seats will lower and raise a person into and out of the bath). Investigating self-care restriction only with reference to an individual's capacity prevents a clear understanding of this form of disability, because it fails to consider important factors which determine its occurrence.

### **1.3 Thesis statement**

The International Classification of Functioning, Disability and Health (ICF) is a biopsychosocial model which offers the opportunity to develop an understanding of self-care through the proposal of a definition and a biopsychosocial model. This allows self-care to be viewed as the outcome of the interaction between the individual and their environment (WHO, 2001). This may better identify potential targets to reduce self-care restriction in older adults. This thesis describes an epidemiological study which investigates self-care restriction in line with the

definition and framework proposed in the ICF. In particular it will look at the links with osteoarthritis and focus on potential determinants within older adults who have the condition. Osteoarthritis is the most common musculoskeletal condition in older adults but previous studies have not provided a clear account of the link with self-care restriction (Covinsky et al., 2008; Donald and Foy, 2004; Dunlop et al., 1998; Davis et al., 1991; Yelin and Katz, 1990).

The overall aim of this study was to describe the population prevalence and distribution of self-care restriction, and associated factors, in adults aged 50 years and over with specific reference to joint pain. The impetus of this research was to identify factors that are potentially modifiable and amenable to interventions that can perhaps enhance self-care in older adults with joint pain.

## **1.4 Overview of research studies**

To achieve the overall aim, existing models of disability in the health literature were reviewed and critiqued, and a systematic review was conducted to identify what is already known about self-care restrictions in older people with joint pain to date. Following this, a number of studies with specific objectives were conducted and are described in the subsequent chapters.

### **1.4.1 Measuring the impact of musculoskeletal conditions in older adults: Models of disability [Chapter 2]**

This chapter describes and critiques the current models of disability which were proposed to guide an understanding of the consequences of health conditions,

including musculoskeletal disease, and provides the rationale for selecting the ICF as a framework to guide the analyses in this thesis (i.e. to investigate the wide range of factors associated with self-care disability in older adults with joint pain).

#### **1.4.2 The impact of osteoarthritis on self-care in older adults: A systematic review [Chapter 3]**

This chapter describes a systematic review conducted to explore the current literature on restricted self-care in community-dwelling adults aged 50 years and over in the general population and its links with joint pain/osteoarthritis.

The aim of the systematic review was to identify and evaluate literature which has examined links between self-care restriction and joint pain / osteoarthritis. The objectives were:

- i. To determine if self-care restriction is associated with joint pain in community-dwelling adults aged 50 years and over in the general population.
- ii. Explore the determinants of restricted self-care in community-dwelling adults aged 50 years and over with joint pain / osteoarthritis.

##### *Rationale*

Osteoarthritis is the commonest joint condition and a major cause of musculoskeletal pain and disability (Hosie and Dickson, 2000). The impact of osteoarthritis is broad, with many associated outcomes. Despite research suggesting links between joint pain and future self-care disability, the extent and mechanisms are unclear. There is a need to conduct a systematic review of the literature to identify studies which have specifically explored the links between self-



care restriction and joint pain and identified the determinants of this outcome in community-dwelling older adults with joint pain / osteoarthritis.

#### **1.4.3 North Staffordshire Osteoarthritis Project (NorStOP): The survey overview and sample analysis [Chapter 4]**

The data used for all analyses described in this thesis were collected in the North Staffordshire Osteoarthritis Project (NorStOP). The aim of this chapter was to critically appraise the potential of NorStOP to fulfil the objectives for this thesis.

To achieve this, the specific objectives were:

- i. To review the NorStOP study design, administration and questionnaire content for the capacity to describe the population prevalence and distribution of self-care restriction, and associated factors, in adults aged 50 years and over with specific reference to joint pain.
- ii. To evaluate the sample derivation including selection, recruitment, response and attrition over the 6 year period (baseline, 3 years and 6 years follow-up) and identify potential bias on findings.

#### *Rationale*

Each research study described in this thesis involved analysis of data from the North Staffordshire Osteoarthritis Project (NorStOP) - a population-based observational cohort of people aged 50 years and over. Notably, the Keele Assessment of Participation (KAP) was developed and applied in NorStOP to measure participation, in line with the conceptual framework of the International

Classification of Functioning (ICF); the conceptual model of this instrument considers self-care to be a form of participation. The suitability and quality of NorStOP data was critically evaluated for the capacity to meet the aims of this thesis. This was evaluated by reviewing the administration of the survey, the quality of data collection (i.e. validity of data) and potential bias related to sample selection and attrition.

#### **1.4.4 Person-perceived self-care restriction in middle and old age: prevalence, distribution and associated factors [Chapter 5]**

The aim of this chapter was to describe the extent of restricted self-care and its links with demographic, socio-economic and health factors in a general population of adults aged 50 years and over. In particular this analysis aimed to establish if joint pain was associated with restricted self-care in the general population.

##### *Specific objectives*

- i. To estimate the prevalence of person-perceived self-care restriction, and its distribution by age and gender.
- ii. To determine the direction and strength of association between restricted self-care and selected health conditions, impairments, activity limitation and contextual factors.
- iii. To derive and validate a parsimonious model of factors which are independently associated with restricted self-care.
- iv. Examine the potential for other factors to 'explain' the relationship between joint pain and self-care restriction.

### *Rationale*

Describing the frequency and distribution of health-related states and events in the general population often provides essential information on their extent and may help to generate hypotheses on possible determinants. Thus, quantification of the prevalence and distribution of self-care restriction and its relation to other health, socio-demographic and contextual factors in a general population of older adults is an essential part of understanding the problem and developing hypotheses around factors that may be associated with restricted self-care. The specific focus on the links between self-care restrictions and joint pain is to establish if those with joint pain are more likely to report self-care restriction than those with no joint pain in older people.

#### **1.4.5 Person-perceived self-care restriction in middle and old age: onset and persistence over a 3 year period [Chapter 6]**

The aim of this chapter was to describe the longitudinal course of self-care restriction in older people in the general population over a three year period.

#### *Specific objectives*

- i. To estimate the extent of the onset and persistence of self-care restriction at 3 years.
- ii. To describe the relationship between onset and persistence of self-care restriction at 3 years and baseline demographic, socio-economic and health characteristics.

- iii. To explore the direction and strength of association between the onset and persistence of self-care restriction at 3 years and the site, number, and pattern of peripheral joint pains at baseline.

### *Rationale*

There are no data on the onset and persistence of participation restriction in self-care. Reported estimates of the onset and persistence of limitations in specific self-care activities measured as part of Activities of Daily Living instruments (ADL) differ between studies, due to a lack of consensus in definitions. It is important to have an estimation of the onset and persistence rates of self-care restriction to understand how the course of self-care restriction develops and persists over time in older people in the general population. Measuring the course of self-care restriction in older people over a 3 years period may help to determine the link between joint pain and future onset or persistence. Examining the demographic, socio-economic and health characteristics of the study cohort provides the estimation of biopsychosocial risk factors associated with the onset and persistence of self-care restrictions in older people in the general population.

#### **1.4.6 Potential predictors of the onset and persistence of restricted self-care in older adults with joint pain [Chapter 7]**

Having established a link between self-care restriction and joint pain, the rest of the thesis will focus on older adults with joint pain. In this chapter, the aim was to identify the potential predictors of the onset and persistence of restricted self-care in older adults with joint pain; this focused on investigation of the association

between onset and persistence and (i) joint specific characteristics (e.g. pain severity, stiffness and chronicity) and (ii) the factors that constitute the parsimonious model of factors established in the general population at baseline (this was to determine whether the risk factors for those with joint pain were different to those for the general population).

### *Specific objectives*

- i. To examine the association between the onset and persistence of self-care restriction at 3 years and joint specific characteristics at baseline.
- ii. To investigate links between the onset and persistence of self-care restriction at 3 years and individual factors that was independently associated with restricted self-care in the cross-sectional analyses at baseline ('the parsimonious model').
- iii. To test the extent to which the parsimonious model predicts the onset and persistence of self-care restriction at 3 years and to determine whether joint-specific characteristics make an additional independent contribution to the prediction of future self-care restriction.

### *Rationale*

Previous studies identified joint specific characteristics such as pain severity as an important risk factor that increase poor physical functioning, and predict future disability in ADLs in older people. Psychosocial factors (e.g. depression) may also influence the relationship between impairments and participation restriction. However, the factors associated with the risk of onset or persisting self-care

restriction in those with joint pain is unknown. It is important to determine if (i) factors particular to osteoarthritis (such as frequent or severe pain) and (ii) demographic, comorbid and socio-economic factors are linked to the onset and persistence of self-care restriction to identify potential targets to reduce this burden.

#### **1.4.7 The concordance between activities and participation and the impact of environmental factors on self-care [Chapter 8]**

The aim of this chapter was to examine (i) empirical differences between task-specific activity limitation and self-care restriction, and (ii) the link between environmental factors, such as receiving help and assistance and using aids and assistance, and self-care restriction in community-dwelling older people with joint pain.

##### *Specific objectives*

- i. To compare the frequency and age and gender distribution of task-specific activity limitations, multiple activity limitation, and relative severity of each limitation in relation to the age and gender distribution of person perceived self-care restriction.
- ii. To estimate the level of agreement between estimates of person-perceived self-care restriction and task-specific activity limitation.
- iii. To determine and compare the risk profiles of person-perceived self-care restriction and task-specific activity limitations.

- iv. To determine whether environmental factors moderate the relationship between restricted self-care and task-specific activity limitation.
- v. To investigate the associations between task-specific activity limitation and person-perceived participation restriction by the need for, and use of, environmental facilitators.
- vi. To examine the socio-demographic, health and joint specific characteristics of the study sample to elucidate the wider factors associated with the need for, and use of, environmental facilitators in older adults with joint pain.

### *Rationale*

In contrast to measuring restriction in self-care as a form of participation and from the perspective of the individual (i.e. measuring whether ones' needs were met as and when they have wanted) limitation in specific self-care tasks was traditionally measured by *difficulty* (none, some, a lot, unable) or *dependence* (requiring help from another person) in executing these tasks (i.e. measuring the individual's capacity (task-specific activity limitation)). Although conceptually different, these two approaches may identify similar populations and there is a need to empirically test if this is true. According to the ICF (WHO, 2001), social and environmental factors impact on all components of functioning and restriction, thus it is important to understand to what extent the relationship between task-specific activity limitation and person-perceived self-care restriction is moderated by environmental factors to identify target populations.

#### **1.4.8 Summary and discussion [Chapter 9]**

This chapter summarises the findings from the analyses, critically appraises strengths and weaknesses, and outlines the implication with relation to the research topic, management of self-care restriction in older adults with joint pain, and future research.



## Chapter 2

# Measuring the impact of musculoskeletal conditions in older adults: Models of disability

## 2.1 Introduction

Disability models have been proposed to guide an understanding of the consequences of health conditions, including musculoskeletal disease. Disability models offer frameworks that allow information to be organised and guide an understanding of disablement (Bickenbach et al., 1999). For research purposes this allows questions to be framed and different disciplines to discuss how factors interact, which improves the understanding of how disability occurs. However the epidemiology of disability is hindered by different definitions and models (Mont, 2007). This prevents a common approach to research and consensus on the frequency of disability and potential causal mechanisms. There is a need for a standard method for defining and characterising disability to facilitate the development of an epidemiological base.

Despite the lack of consensus, all disability models draw on the medical and social models of disability to a greater or lesser extent. In the medical model disability is defined as “*an observable deviation from biomedical norms of structure or function that directly results from a disease, trauma or other health condition*” (Boorse, 1975: p.19). This places disability as a consequence of a health condition and fails to take account of social, psychological or environmental factors which may help to reduce or enhance the effects of health conditions (Albrecht, 2010; Jette, 2006; WHO, 2001). This affects the potential to reduce disability for those with chronic

health conditions which persist despite medical treatment. In contrast, the social model views disability as a consequence of the environment (e.g. the lack of public transport prevents people with leg pain and problems walking from getting to work or shops) (Albrecht, 2010). Disciplines that focus on the social model often underplay the importance of the signs and symptoms of health conditions and don't link these with the environmental and social factors which lead to disability (Bury, 2000). The complexity of the consequences of health conditions suggests that integrated models, which consider both the medical and social models, may provide the best opportunity to characterise disability and facilitate a greater understanding.


This chapter aims to describe and critique key models which have guided strategies to measure disability in epidemiological studies such as the Nagi model (Nagi, 1965), which has subsequently been revised, two models proposed by the WHO (The International Classification of Impairment, Disability and Handicap and the International Classification of Functioning, Disability and Health) and models of participation developed in Occupational Therapy to provide the rationale behind the chosen framework to guide the research studies undertaken in this thesis.

## 2.2 An overview of models of disability

### 2.2.1 The Nagi model of disablement (Nagi, 1965)

The Nagi model of disablement (Nagi, 1965) is a theoretical model of health, functioning and disability and was one of the first attempts to structure the consequences of health problems into different levels. It has been subsequently revised and provides the basis for other models, discussed below. In this model, disablement was explained through four concepts; pathology, impairment, functional limitation, and disability (Nagi, 1965). In this, *active pathology* referred to a state of the body's defences and coping mechanisms caused by infections, trauma, metabolic imbalance, degenerative disease processes, or other pathologies; *impairment* referred to a loss or abnormality at the tissue, organ, and body system level; functional limitation referred to the individual's ability to perform the tasks and obligations of his usual roles and normal daily activities (Pope and Tarlov 1991); and *disability* was defined as an 'expression of a physical or a mental limitation in a social context'. Thus, this model suggested that impairment is associated with functional limitation (i.e. limitation in performing basic physical (i.e. walking, lifting objects, climbing stairs, reading standard-sized print and hearing other people) and mental actions (short-term memory, intelligible speech, alertness, orientation), that are involved in daily activities. However this is not directional and will depend on personal and social factors (i.e. loss of a finger could be severely limiting to a pianist but might not be limiting at all to a teacher) (Nagi 1965, p.102) (Figure 2.1).

**Figure 2.1     Nagi Model (Nagi, 1965)**



**Source:** Albrecht et al, 2010


This model was revised in 1991 and the *disability* component was removed due to the view that disability is not inherent within the person. The revised model proposed that the patterns of an individual's impairments are associated with functional limitations and the interaction of the individual and the environment (Nagi, 1991). In this revised model functional limitation may not be a direct result of impairment, but could be also due to personal, social and/or environmental factors, emphasising that disability is the result of the gap in an individual's own capabilities and their social and physical environment (Jette, 2006). However, the revised Nagi model still lacked clarity and necessary detail to study the specific aspects of disability. Thus, following Nagi's own revision, other conceptual models were developed based on the revised model, with the main inclusion being the role of social and physical environment in the disablement process. These models are summarised and discussed below.

### *Institute of Medicine Model*

The development of the Institute of Medicine's (IOM) Model (The enabling-disabling process) (Pope and Tarlov, 1991) involved modifying the Nagi model, to propose a prevention based rehabilitative model (Brandt and Pope, 1997). This model proposed a new conceptual approach by analysing the components of disability, and describing the importance of different risk factors. Nonetheless, over time, the original IOM model (1991) has been criticised for proposing a disablement process that is unidirectional, progressive toward disability without the possibility of its reversal (due to the arrows in the model that pointed toward disability), limited classification of the environment, and its' interaction with the individual (Brandt and Pope, 1997). Even though the significance of the environment is discussed in this model, it was not clearly represented in the model.

The IOM model was revised in 1997 and the new model was designed to provide clarity in describing disability as the interaction of the person with the environment and also to show the possibility of movement in the direction of rehabilitation (Brandt and Pope, 1997). This model defined disability as "a function of the interaction of the person with the environment" and focused on the environmental factors, suggesting that physical and social environmental factors act as risk factors in the disablement process (Masala and Petretto, 2012). This highlighted the role of environmental factors in either leading to or preventing disability. The IOM model also included bidirectional arrows and a state of "no disabling condition" that depicted disability as a reversible state (Figure 2.2).

**Figure 2.2 The IOM Enabling-Disabling Process**




**Source:** Brandt and Pope, 1997.

*The National Centre for Medical Rehabilitation Research Model*

The National Centre for Medical Rehabilitation Research (NCMRR) (National Institute of Health (NIH), 1993) model was also developed using Nagi's model, and included consideration of the environment in the dimension called "societal limitation". Societal limitations were defined as barriers to full participation in society that result from attitudes, architectural barriers and social policies. This

model suggested that personal factors may influence an individual's response to a specific situation, thus the interaction among organic, psycho-social, personal and environmental factors would produce the individual's response to impairment (NIH, 1993). The NCMRR model highlighted the fact that the barriers between individuals and the environment could be potentially reduced or eliminated by public policies and services (Figure 2.3).

**Figure 2.3     The NCMRR Model**



**Source:** The NCMRR Model, 1993.

#### *The Disablement Process Model*

Verbrugge and Jette (1994) extended Nagi's model further and developed "The Disablement Process Model" with the aim of facilitating health research. The term 'disablement' referred to the effect of acute and chronic conditions on the functioning of body systems, and individual's interactions with the environment. The term 'process' considered the impact of changes in functional consequences on individuals over time. This model was based on a "socio-medical" approach (i.e. one that combines the medical and social models of disability) and defined functional limitation and disability as separate behaviours as opposed to different features of measuring the same behaviour. In this model *functional limitations*

were defined as *restrictions* in performing fundamental physical and mental actions used in daily life by one's age-sex group, and *disability* was defined as 'experienced difficulty in doing activities in any domain of life due to a health or physical problem' (Verbrugge and Jette, 1994).


The disablement process model highlighted the importance of social and environmental factors by emphasising that limitations in activities were not just a result of poor physical functioning because they involved interaction with the environment (e.g. washing oneself would require to undress and get in and out of bath/shower, dry oneself). This model elaborated further on the importance of 'intrinsic' (i.e. capacity: e.g. having difficulty in carrying out a task such as washing without a personal or equipment assistance) and 'actual' (i.e. performance: e.g. having difficulty with carrying out a specific task with personal and equipment assistance) disability, underlining the implications on future health-care policies and design and development of aids and adaptations tools. This division had previously been criticised by researchers, health-care practitioners and civil rights advocates because many people use external sources, such as help and assistance, to enable them to function adequately. Therefore it may be better to measure actual disability rather than intrinsic disability (i.e. it is better measure performance rather than the capacity to perform) (Verbrugge and Jette, 1994).



### 2.2.2 The International Classification of Impairments, Disabilities, and Handicaps (ICIDH)

The ICIDH, published in 1980, was the first attempt by the WHO to provide a common language to describe the consequences of health and disease, and analyse health related issues. The ICIDH linked with the International Classification of Diseases (ICD), which was established to provide a standard diagnostic tool for epidemiology to monitor the incidence and prevalence of diseases and other health problems by WHO in 1946 with publication of ICD-6. Notably its origin can be traced back into the Bertillon Classification, which was first published in 1893. The ICIDH defined the consequences of health in three sequential levels; body (impairment), individual (disability) and societal (handicap) (Figure 2.4). Impairment was defined as *“any loss or abnormality of psychological, physiological or anatomical structure or function”*. Disability was defined as *“any restriction or lack (resulting from an impairment) of ability to perform an activity in the manner or within the range considered normal for a human being”*. Handicap was defined as *“disadvantage, resulting from impairment or a disability that limits or prevents the fulfilment of a role that is normal (depending on age, sex, social and cultural factors) for that individual”* (WHO, 1980).

**Figure 2.4 The ICIDH Model**



**Source:** WHO, 1980

In the development of the ICIDH, the WHO attempted to integrate the social and medical models of disability by recognising the social consequences of health conditions (i.e. through the inclusion of the handicap concept). Although the ICIDH in general was welcomed in health research, it was criticised for its alignment with the medical model, because its' linear approach (i.e. impairment causes disability which then causes handicap) implied that disability is only a consequence of health conditions (Whiteneck, 2006). Although it acknowledged that health conditions interact with social and environmental factors through the inclusion of handicap, it did not implicitly outline the role of these factors on an individual's function (Imrie, 2004; Bornman, 2004; Simeonsson et al., 2000; Chapiereu and Colvez, 1998). The model implied that handicap was an outcome of an individual's impairment, isolated from the societal constraints and choices which were offered to the individual and did not reflect the real experiences of those people living with the health conditions (Simeonsson et al., 2000). The linear approach also failed to show that the body, individual and societal levels were distinctive. It was not possible to assess the degree of severity of disabilities or handicaps using the ICIDH (WHO and Economic and Social Commission for Asia and the Pacific (ESCAP), 2008). Therefore, the ICIDH was seen as inconsistent in general, and its use was not commonly adopted by disability researchers and advocates of the social model of disability (WHO and ESCAP, 2008).


### **2.2.3 The International Classification of Functioning, Disability and Health (ICF)**

As a result of the criticisms of the ICIDH, the WHO collaborated with international researchers to put together a framework which married the medical and social models of disability and incorporated environmental factors. Following seven years of expert partnership, process and validation in over 70 countries, the ICF was officially endorsed by the WHO in 2001.

The ICF was conceptually different to the ICIDH as its rationale was to capture the components of health rather than classify the consequences of disease (Ustun et al., 2003). This is why the ICF was named as it is, with an emphasis on health and functioning, rather than on disability (Jette, 2002). This acknowledges that (i) people can continue to function positively despite having health problems and (ii) that everyone can experience some form of disability, independent of their health status.

The ICF framework was based on a biopsychosocial model of health, which was previously introduced by George Engel (1977; 1980) (Figure 2.5). The biopsychosocial model was a departure from the previous linear, cause–effect based approach to health and included the psychosocial dimensions (personal, emotional, family, community) as well as the biological aspects of diseases.


**Figure 2.5     The biopsychosocial model of health and illness**



**Source:** Engel, 1977

The ICF defined functioning and disability as multi-dimensional concepts, relating to (i) body functions and structure, (ii) the activities people do and the life areas in which they participate, and (iii) the factors in their environment that affect these experiences. Within the ICF, a hierarchy of classifications and codes were provided for each of these components. In this model, an individual's functioning or disability was conceived as a dynamic interaction between health conditions and personal and environmental factors (WHO, 2001) (Figure 2.6).

**Figure 2.6**      **Diagram of interactions between the components of ICF**



**Source:** WHO, 2001

The ICF organised information in two sections, (i) function and disability, and (ii) contextual factors, to encourage and facilitate a broader understanding of human functioning (WHO, 2001).

#### (i) Functioning and disability

The ICF domains were summarised at three levels (body, individual and societal) as body structures and functions, activities and participation, which also indicated positive functioning (Table 2.1). In the ICF, disability is an umbrella term for any negative functioning in these three levels which are referred to as impairments, activity limitation and participation restriction (WHO, 2001). Impairment was defined as problems in body function and structure, such as a significant deviation or loss (WHO, 2001). Activity limitations were defined as difficulties that an

individual may have in executing tasks (WHO, 2001). These tasks could be any activities related to the individual's life and encompasses the use of purposeful body functions. Participation restrictions were defined as problems an individual may experience in life situations when interacting with society and/or environment which may enable or restrict participation (WHO 2001). Thus, participation restriction is the result of a complex interaction between the individual (including the consequences of a health condition) and the environment (Stucki et al., 2002; Grimby et al., 2001).

The ICF indicates that both concepts (activity limitation and participation restriction) can be measured with reference to either capacity or performance and judged against accepted population standards (i.e. what is normally expected from someone of a specific age and gender) (WHO 2001). In the ICF, *capacity* is defined as a construct which describes an individual's ability to execute a task or an action in a standardised environment, and it is useful to indicate the highest probable level of functioning that a person can reach in the domain that is being evaluated (i.e. actual ability). To evaluate the individual's full capacity, a standardised environment is necessary to remove the impact of environmental factors. Thus, capacity reflects the individual's ability adjusted to the environment. On the other hand, *performance* refers to what individuals do in their current environment (Jette, 2002). Therefore, *"the gap between capacity and performance reflects the difference between the impacts of current and uniform environments, and thus provides a useful guide as to what can be done to the environment of the individual to improve performance"* (WHO, 2001:p.15).

The ICF recommends judging the capacity and performance of individuals against population standards (i.e. what is expected of a similar individual without the similar impairment). However participation restriction cannot be the same for each person due to changing roles, environments and personal factors, thus judging against a population standard does not reflect the subjective dimension of functioning and disability (Ueda and Okawa 2003; Harwood et al., 1994). However, taking the participant's perception is a highly relevant dimension for clinical practice as perceived disability may identify care needs. It is also difficult to define a population standard for judging participation, because different needs direct whether participation will occur at all in some areas of life (e.g. older adults who have sufficient financial means to allow them to retire may no longer choose to work) (Peronboom and Chorus, 2003).

**Table 2.1      The ICF components and definitions**

<b>Level</b>	<b>Functioning</b>	<b>Definition</b>	<b>Disability</b>	<b>Definition</b>
Body	Body structures and functions	Anatomical parts of the body such as organs, limbs and their components. Physiological components of body (including psychological functions)	Impairments	Problems in body function or structure such as significant deviation or loss
Individual	Activities	Execution of a task or action by an individual	Activity limitation	Difficulties an individual may have in executing activities
Societal	Participation	Involvement in life situations	Participation restriction	Problems an individual may experience in involvement of life situations

**Source:** ICF (WHO, 2001)

To further aid a common language and approach to research, different domains of disability were also outlined in the ICF; this was a single list of domains and sub-domains that are proposed to cover the full range of life areas. However, the ICF does not direct whether these domains refer to activity limitation or participation restriction (Table 2.2).

**Table 2.2 The ICF concepts in context**

<b>Body</b>	
<b>Function</b>	<b>Structure</b>
Mental Functions	Structure of the Nervous System
Sensory Functions and Pain	The Eye, Ear and Related Structures
Voice and Speech Functions	Structures Involved in Voice and Speech
Functions of the Cardiovascular, Haematological, Immunological and Respiratory Systems	Structure of the Cardiovascular, Immunological and Respiratory Systems
Functions of the Digestive, Metabolic, Endocrine Systems	Structures Related to the Digestive, Metabolic and Endocrine Systems
Genitourinary and Reproductive Functions	Structure Related to Genitourinary and Reproductive Systems
Neuromusculoskeletal and Movement-Related Functions	Structure Related to Movement
Functions of the Skin and Related Structures	Skin and Related Structures
<b>Activities and Participation</b>	
Learning and Applying Knowledge	
General Tasks and Demands	
Communication	
Mobility	
Self Care	
Domestic Life	
Interpersonal Interactions and Relationships	
Major Life Areas	
Community, Social and Civic Life	
<b>Environmental Factors</b>	
Products and Technology	
Natural Environment and Human-Made Changes to Environment	
Support and Relationships	
Attitudes	
Services, Systems and Policies	

**Source:** ICF (WHO, 2001)



## (ii) Contextual factors

The ICF includes personal and environmental factors to set an individual's function in the context of their life (WHO, 2001); these contextual factors “*represent the complete background of an individual's life and living*” (Murray, 2002). Personal factors (i.e. characteristics that exist independently from the presence or absence of disease or the functional state) are not classified in the ICF framework, and refer to attributes such as age, sex, educational background, social class, culture, past experiences, personal character traits, lifestyle, coping styles and occupation (Jette, 2002). Environmental factors encompass the physical, social and attitudinal environment in which people live and conduct their lives (Table 1.2) (WHO, 2001). The model proposes five categories; ‘Products and Technology’ (e.g. any product, instrument, equipment or technical system used by a disabled person, especially produced or generally available, preventing, compensating, monitoring, relieving or neutralizing disability), ‘Natural Environment and Human-Made Changes to Environment’ (e.g. animate and inanimate elements of the natural environment, and components of that have been modified by people, as well as characteristics of human populations within that environment), ‘Support and Relationships’ (e.g. people or animals that provide practical physical or emotional support, nurturing, protection, assistance and relationships to other persons, in their home, place of work, school or at play or in other aspects of their daily activities), and ‘Services, Systems and Policies’ (e.g. systems and services that provide benefits, and policies govern and regulate the systems that organize, control and monitor services, structured programmes and operations in various sectors of society). Environmental factors can impact the individual's life positively or negatively during

life situations. The environment can be viewed as immediate (e.g. physical features of the environment, social environment) or societal (e.g. formal and informal social structures, services and systems). Particular environments may have a specific impact on an individual, regardless of their health condition (WHO, 2001).

### *Strengths and limitations of the ICF*

The ICF framework is based on a biopsychosocial model which integrates the medical and social models and allows users to record the impact of health conditions on human functioning at biological, individual, and societal levels (Reed et al., 2005). In addition, the ICF also includes personal and environmental factors to encourage acknowledgement of their influence on an individual's function and disability. This is an important addition as both personal and environmental factors may act as barriers or facilitators to participation (Allan et al., 2006).

The ICF also provides a clear definition of the three levels of disability, offering “*a unified and standard language and framework for the description of health and health-related states*” (WHO, 2001:p. 3). This terminology allows the communication of health information across professional disciplines; the presence of a common framework and language across disciplines facilitates inter-professional collaboration (Giacomini, 2004).

The neutrality of the ICF definitions allows the framework to be applied to all populations, allowing comparisons across health conditions. However the flexible

nature of this framework has also been criticised, as the definitions offered may not be sufficiently clear to lead the standardisation of such information (Bornman, 2004; Stucki et al, 2002). It is also being suggested that the lack of clarity between the concepts of activity limitation and participation restriction prevents clear application of this framework (Fougeyrollas and Beauregard, 2001).

#### **2.2.4 Models of participation developed in Occupational Therapy**

The author of this thesis is an occupational therapist and this profession has developed models of participation and function to direct practice and research. Occupational therapy (OT) is an allied health-care profession that aims to promote health by enabling individuals to perform meaningful and purposeful activities to develop, recover, or maintain their daily living (World Federation of Occupational Therapy (WFOT), 2011). The theoretical models of disability relate closely to the OT practice of supporting health and participation in life through engagement in occupation (American Occupational Therapy Association (AOAT), 2002). The Canadian Model of Occupational Performance (CMOP) was developed by the Canadian Association of Occupational Therapists, and illustrates the relationship between person, occupation (activities of daily living) and environment (Fearing et al., 1997). This model suggests that all these three levels (person, occupation and environment) are interdependent and the change in one area affects all others (Ward et al., 1996). The model is applied in practice by the accompanying Canadian Occupational Performance Process Model (COPM), which describes the therapeutic process from a client's perspective (Clarke, 2003). Model of Human Occupation (MOHO) is another occupational therapy model developed in 1985.

MOHO is a conceptual model of practice that evolved from Reilly's *Occupational Behavior Model* and *General System Theory* (Reilly, 1974). This model aimed to explore what motivates individuals in establishing everyday routines and how the environment influences these behaviours using concepts such as volition, habituation, performance capacity and environmental context (Kielhofner et al, 2002). Its application in clinical practice aids the therapist to understand that there are multiple factors within the person and the environment that influence each other. Thus a change in any one factor will spontaneously result in a change in one's motivation, behavior, and/or performance (Cole, 2008). MOHO has both structured and unstructured assessments. The Model of Human Occupation Screening Tool (MOHOST) is an assessment that addresses the majority of MOHO concepts, and helps the therapist to gain an overview of the client's occupational functioning. Criticisms around the use of MOHO, are the lack of consideration around the influence of the environment on human behaviour, and appreciation of the dialectic process between the human and the environment (Haglund and Kjellberd, 1999). However the MOHOST assessment went through a number of revisions by Kielhofner and colleagues periodically and subsequently its validity was supported by using both classical and modern test theories (Ay-Woan Pan et al., 2010).

These models use terms specific to the OT profession and require specialist training in application for clinical and research practice. This creates a barrier for other health-care and research practitioners to interpret and compare information, and isolate the use of these models to OTs only. OTs involvement in inter-agency

and inter-disciplinary work is increasing, and the use of generic models and frameworks that complement OT philosophy and framework could prove beneficial for the future of multidisciplinary working (College of Occupational Therapists (COT), 2004; Baum, 2002). The biopsychosocial approach taken in the ICF framework complements the person-centred approach that is central to OT philosophy and beliefs (Brintnell, 2002; College of Occupational Therapists, 2002; Law and Baum, 2001). Current OT textbooks include chapters that compare OT models to the ICF (Atwal and Jones, 2009; Kielhofner, 2008; Davis, 2006; Christiansen et al., 2005), and the WFOT has incorporated the ICF into the definition and description of OT (WFOT, 2011).

### **2.2.5 Overview of models of disability**

The disability models discussed in this section were developed over time, utilising criticisms and constantly evolving social, cultural and environmental demands to create a better model. The Nagi model brought together a way to classify the impact of health conditions into different levels, and subsequent revisions have included environmental factors. Nevertheless, these models were not able to offer a framework in which the consequences of health conditions could be classified, or provide domains of disability where an individual's participation in life situations could be assessed. Thus, several models of disability were developed over the years, but none of these were universally adopted. The ICF framework was launched globally in 2001 by the WHO with this intention.

The ICF framework is the most recently developed model, developed to organise information on the disability process, and consist of, body functions / impairments, activity, participation, personal and environmental factors. The use of the ICF framework is promoted in OT to encourage inter-disciplinary collaboration through use of common terminology, and supports the design and organisation of epidemiological studies, which can provide evidence for the holistic OT philosophy and patient-centred interventions.

One of the main advantages of the ICF framework is its capacity to provide a standardised method for collecting and presenting health data (Wilkie et al., 2005). This could address the complications created by the use of different terminologies for the same concept within the literature and/or the same terminology referring to different concepts respectively (e.g. the difference between functional limitation and activity limitation; self-care and self-management). These differences could be observed in different studies as well as government legislation and policies (Simeonsson et al, 2000).

As the philosophy of epidemiology is based on the approach that the frequency of occurrence of a health outcome in a population is governed by the interaction of a large number of different factors or determinants, the use of clear terminology to differentiate between different concepts is an important step in building the epidemiology of self-care disability in older people. In the context of this thesis, being able to clearly differentiate between self-care disability and other disabilities, such as locomotor disability, is of great importance. This allows a clear construct

to be operationalized to allow description of the course of self-care restrictions in community-dwelling older people and its links with joint pain, health, socio-demographic and environmental factors which can help guide the development of future interventions, services and policies to better older people's lives.

Despite criticisms, the ICF is an accepted tool internationally. There are growing numbers of implementations of the framework around the globe, in health information systems and surveys (Madden et al., 2005). The ICF provides a framework to direct data collection in population surveys by offering clear definitions for constructs. Consistent adherence to these definitions will allow comparisons between studies, and build the epidemiology of the health outcome of interest (e.g. self-care restriction). The different domains offered in the ICF permit different types of disability such as self-care to be explored using a standard definition. The ICF framework also allows data to be organised into relevant sections such as impairments, activities, personal and environmental factors, providing a framework for investigating associations and potential pathways to the consequences of health conditions. In the ICF, the locus of problem is not only the individual, and includes a wide range of factors that could be linked to self-care disability in older adults. This could help to expand the knowledge of potential strategies to manage self-care in populations (Table 2.3).

**Table 2.3      A critical review of the models of participation and disability**

<b>The conceptual models of participation and disability</b>	
<b>Nagi Model</b> (Nagi, 1965)	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• It was the first disablement model to recognise that environment, family, society, and community factors may influence disability.</li> <li>• It has changed the perception of disability away from a focus on physical limitations to interaction between the individual and the environment.</li> <li>• It provided a basis for future models.</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• It was based on a uni-dimensional (pathology-based) model of disability.</li> <li>• It does not account for impairments and functional limitations not due to pathology (e.g. obesity, or sedentary life style) (McKeough 2009).</li> </ul>	
<b>Institute of Medicine (IOM) Model</b> (Brandt and Pope, 1997; Pope and Tarlow, 1991)	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• Proposed three interacting risk factors as an addition to the Nagi model; biological, environmental (social and physical), and lifestyle/behavioural that affects each component of the disabling process: pathology, impairment, functional limitation, and disability.</li> <li>• Included the impact on quality of life as a new component in disabling process.</li> <li>• Focused on the effects of the environment in producing or reducing disability.</li> <li>• The revised model (1997) provided clarity in describing disability and included bidirectional arrows to show interaction between different levels.</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• Disability was depicted in the 1991 model as a linear process.</li> <li>• The earlier model (1991) gave a “limited characterisation of the environment and the interaction of the individual with the environment.</li> </ul>	
<b>The National Centre for Medical Rehabilitation Research (NCMMR) Model</b> (1993)	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• Added ‘societal limitation’ to the disablement process</li> <li>• Included personal factors as a potential influence on an individual’s response to a life situation.</li> <li>• Highlighted that the barriers between individuals and environment could be reduced or eliminated by public policies and services.</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• The NCMRR model was a linear model of disablement.</li> <li>• It did not account for the social or physical environment.</li> <li>• It did not offer an adequate working model for practice and research on its own due to use of broad definitions when describing different levels in the proposed disablement process.</li> </ul>	



<b>The conceptual models of participation and disability</b>	
<b>The Disablement Process Model</b> (Verbrugge and Jette, 1994)	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• It described how chronic and acute conditions affect functioning in specific body parts, mental function and daily life, and the role of personal and environmental factors in disablement.</li> <li>• Encouraged research into the impact of risk factors (intra and extra-individual factors) on impairment, functional limitation, disability and the disablement process.</li> <li>• Suggested that it may be better to measure 'actual disability' rather than intrinsic disability (without personal or equipment assistance).</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• Although it is proposed as a socio-medical model of disability, it has been criticised for ignoring 'impairment' by emphasising the effects of social and environmental factors.</li> </ul>	
<b>The (ICIDH) Model</b> (WHO, 1980)	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• It was the first attempt by WHO to provide a common language to describe the consequences of health and disease.</li> <li>• It was linked with the International Classification of Diseases (ICD), which aimed to provide a standard diagnostic tool for epidemiology to monitor the incidence and prevalence of diseases and other health problems.</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• It was criticised for being a medical model; it implied a linear relationship (i.e. impairment causes disability which then causes handicap).</li> <li>• It did not implicitly outline the role of social and environmental factors on an individual's function.</li> <li>• Its linear approach failed to show that the body, individual and societal levels were distinctive.</li> <li>• Introduced the use of the term handicap, which resulted in people being labelled as 'handicapped'.</li> </ul>	
<b>The International Classification of Functioning (ICF) Model</b> (2001)	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• The emphasis was on health and functioning, not on disability. This allowed researchers to explore pathways.</li> <li>• Based on a biopsychosocial model of health, it was a departure from previous linear, cause-effect based approaches to the consequences of health.</li> <li>• It was the first model to categorise environmental factors, which allowed the consideration of societal consequences on disability in a greater clarity.</li> <li>• The framework included the psychosocial dimensions as well as the biological aspects of health conditions.</li> <li>• Shifted the focus from the pathological processes to the level of health and participation.</li> <li>• Provided clear definitions for different domains, offering an international language for use in research and clinical practice.</li> </ul>	

<b>The conceptual models of participation and disability</b>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• Its emphasis on health rather than disability meant that the model does not provide a clear pathway to disability, or offer a model of disablement process, instead promote a new language to define the consequences of health conditions which might be interpreted as introducing further complexity to disability literature.</li> <li>• The interpretation and categorisation of the concepts of activity and participation had been widely criticised for the difficulty in differentiating the two concepts. The ICF describes them conceptually different but provides a single combined list of life areas that are not necessarily linked to one or another.</li> <li>• Personal factors were not clearly defined due to its wide variability.</li> </ul>	
<b>Canadian Model of Occupational Performance (CMOP) (Fearing et al., 1997)</b>	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• Illustrated the relationship between person, occupation (ADL) and environment with a three dimensional model; suggesting that a change in one area automatically affects the others.</li> <li>• With the person at its centre, the model conveys Occupational Therapy's (OT) client centred perspective.</li> <li>• Designed to demonstrate change in a client's self-perception of occupational performance over the course of occupational therapy, thus, aiding the therapy in the rehabilitation process.</li> <li>• Applied in practice through an accompanying outcome measure; Canadian Occupational Performance Measure (COPM) designed to help clients identify, prioritise, and evaluate important issues they encounter in occupational performance.</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• Through its use of terminology specific to OT and the need for specialist training in application to the clinical and research practice, it creates a barrier for other health-care/ research professionals to interpret the results of this assessment and compare information with other studies.</li> </ul>	
<b>The Model of Human Occupation (MOHO) (Kielhofner et al., 2002)</b>	
<b>Strengths</b>	
<ul style="list-style-type: none"> <li>• MOHO emphasised that to understand human occupation, it is important to understand the physical and social environments in which the occupation is performed and conceptualised human occupation as a product of three interrelated components; volition, habituation, and performance capacity, taking a person centred stance.</li> <li>• It is applied in clinical practice and research by both structured and unstructured assessments such as the Model of Human Occupation Screening Tool (MOHOST) to provide the therapist an overview of the client's functioning.</li> </ul>	
<b>Limitations</b>	
<ul style="list-style-type: none"> <li>• It has been criticised for its lack of consideration around the role of environmental factors on human behaviour and impact of this on the occupational performance.</li> <li>• Similar to CMOP, this model is not openly accessible to those who are not occupational therapists.</li> </ul>	

### 2.3 The ICF model of self-care

Differences in definitions and measurement of self-care across studies highlight the need for use of a common language and a consistent approach to define and quantify self-care restrictions in populations. The ICF offers a definition and a model of self-care to encourage a consistent approach. In this framework self-care is presented as one domain of functioning that warrants particular attention to understand how common it is and the factors linked with restriction (WHO, 2003).

The ICF defines self-care as *“caring for oneself, washing and drying oneself, caring for one's body and body parts, dressing, eating and drinking, and looking after one's health”* (WHO, 2001). Each of the sub-domains of self-care is also defined within the ICF framework to avoid ambiguity and provide guidance for analysis (D510; Washing oneself, D520; Caring for body parts, D530; Toileting, D540; Dressing, D550; Eating, D560; Drinking, D570; Looking after one's health) (Table 2.4). Having clear definitions of self-care allows differentiation from other forms of disability such as locomotor disability and other constructs such as self-maintenance (i.e. management of health). The ICF offers a biopsychosocial model of self-care that accounts for the context of an individual's life when assessing disability.

The ICF does not propose whether self-care should be considered as “Activities” or “Participation”, or in other words, whether it should be considered as functioning at the level of the individual or society. One of the clear strengths of the ICF framework is that it encourages a biopsychosocial approach to understanding “disability”. The framework encourages a wider approach to investigating factors

that are associated with self-care restriction and in particular the role of environmental factors. For example an older person with joint pain may experience difficulty bending to wash, dress and pick things up. In this case access to environmental facilitators such as a grab-rail by a bath or shower cubicle can assist transfers, whilst dressing aids such as button hooks can enable those with grip problems to put on and take off shirts and trousers. Similarly, those who are in receipt of personal help and assistance for personal care activities such as washing and dressing or caring for body parts may feel that their needs in self-care have been met, hence, may not be disabled by their limitations.

The model of self-care provided by the ICF offers opportunities to develop the epidemiology of self-care using a common language and a consistent approach. Therefore, this model was adopted as a conceptual framework in this thesis to investigate the course of self-care restriction and its links with joint pain in community-dwelling older people.

**Table 2.4 The self-care and its components in the ICF framework**

<b>D5- Self-care Domain</b>	
Sub-domain	Definition
D510 - Washing one-self	Washing and drying one's whole body, or body parts, using water and appropriate cleaning and drying materials or methods, such as bathing, showering, washing hands and feet, face and hair, and drying with a towel.
D520 - Caring for body parts	Looking after those parts of the body, such as skin, face, teeth, scalp, nails and genitals that require more than washing and drying.
D530 - Toileting	Planning and carrying out the elimination of human waste (menstruation, urination and defecation), and cleaning oneself afterwards.
D540 - Dressing	Carrying out the coordinated actions and tasks of putting on and taking off clothes and footwear in sequence and in keeping with climatic and social conditions, such as by putting on, adjusting and removing shirts, skirts, blouses, pants, undergarments, saris, kimono, tights, hats, gloves, coats, shoes, boots, sandals and slippers.
D550 - Eating	Carrying out the coordinated tasks and actions of eating food that has been served, bringing it to the mouth and consuming it in culturally acceptable ways, cutting or breaking food into pieces, opening bottles and cans, using eating implements, having meals, feasting or dining.
D560 -Drinking	Taking hold of a drink, bringing it to the mouth, and consuming the drink in culturally acceptable ways, mixing, stirring and pouring liquids for drinking, opening bottles and cans, drinking through a straw or drinking running water such as from a tap or a spring; feeding from the breast.
D570 - Looking after one's health	Ensuring physical comfort, health and physical and mental well-being, such as by maintaining a balanced diet, and an appropriate level of physical activity, keeping warm or cool, avoiding harms to health, following safe sex practices, including using condoms, getting immunizations and regular physical examinations.

**Source:** International Classification of Functioning (WHO, 2001)

## **Chapter 3**

# **The impact of osteoarthritis on self-care in older adults: a systematic review**

### **3.1 The burden of musculoskeletal conditions in older adults**

The prevalence of musculoskeletal conditions is high and they are the most common cause of disability in older adults (Woolf and Pfleger, 2003). Musculoskeletal conditions are rarely fatal and prevalence rises with age (Verbrugge, 1995). With population ageing there will be an increasing number of older adults with these conditions. Despite the large and growing impact, identifying and managing musculoskeletal conditions has not been a health care priority in the United Kingdom (UK) or other western countries (i.e. in Europe or the United States). Initiatives, such as the Bone and Joint Decade (1998), have raised awareness of musculoskeletal diseases, but there is still a need to better understand the full burden of these conditions on the individual and society (Woolf and Pfleger, 2003). This includes recognising the frequency, incidence, causes, progression and consequences to inform preventative and management strategies.

### **3.2 Osteoarthritis**

Osteoarthritis is the most common musculoskeletal condition in older adults (Arthritis Care, 2004; Arthritis Research Campaign, 2002; Scott et al., 1998). In the UK in 2000 it was estimated that there were just over three million consultations to general practice and over one hundred thousand admissions to hospital for

osteoarthritis (Arthritis Research Campaign, 2002). It is strongly associated with age and is expected to have an increasing impact on health and social care needs (Woolf and Pfleger, 2003).

### **3.2.1 Definition of osteoarthritis**

*“The name osteoarthritis was adopted from three Greek words meaning bone, joint, and inflammation”* (Mosby's Medical Dictionary, 2009: osteoarthritis). It can be defined by two different but overlapping syndromes; the disease and the clinical syndrome of joint pain and disability (Moskowitz, 2009). As a disease, osteoarthritis is defined as a non-inflammatory disorder of synovial joints that is characterised by focal areas of damage to the articular cartilage, remodelling of underlying bone and the formation of osteophytes (i.e. new bone at joint margins), and mild synovitis (Dequeker and Luyten, 2008; National Collaborating Centre for Chronic Conditions, 2008). In osteoarthritis the disease processes do not only affect the articular cartilage, but involve the entire joint, including the subchondral bone, ligaments, capsule, synovial membrane, and periarticular muscles (Brandt et al., 2008).

The articular cartilage is whitish in colour, elastic and very strong due to its components; which consist mostly of water (nearly 65%), proteoglycans, and collagens. Cells embedded throughout the cartilage called chondrocytes are responsible for synthesising proteins that make up cartilage, as well as the enzymes that break down the cartilage maintaining the balance of repair and degradation. A number of enzymes are produced by chondrocytes called

metalloproteinases (Lane and Wallace 2002). Normally, these enzymes are found in very low concentrations in cartilage with normal levels of their inhibitors (tissue inhibitors of metalloproteinases (TIMPs)) (i.e. *“naturally occurring small proteins which inhibit metalloproteinases so that they do not break down too much cartilage”* (Lane and Wallace 2002:p.32). However, in osteoarthritis, there is a change in this metabolic balance, so the levels of these enzymes increases and TIMPs decrease, breaking down the cartilage faster than the chondrocyte can repair, resulting in a loss of cartilage (Lane and Wallace 2002). Therefore, the disease process in osteoarthritis usually starts with inflamed and swollen cartilage when the joint loses proteoglycan molecules, other tissues and water. Cartilage loses its elasticity and fluid, and cracks starts to appear on the surface. In this process, the cartilage hardens and becomes more prone to further damage with continuous usage and loading. Over time, large amounts of the cartilage can be lost, leaving the ends of the bone within the joint unprotected. As the body reacts to the loss of cartilage, clusters of damaged cells or fluid-filled cysts form on the bony areas or near the splits in the cartilage. This may also happen within the bone marrow itself, which results in swelling. The marrow is rich in nerve fibres. These changes may cause pain in those with osteoarthritis. Bone cells may respond to damage by multiplying, growing, and forming dense, misshapen plates around exposed areas (i.e. osteophytes) causing limitation in joint movements and pain (Simon, 2009).

Definition and diagnosis of osteoarthritis has proven to be challenging (Peat et al. 2005). Due to the complex nature of OA, the clinical presentation and outcomes



can be extremely variable, both between individuals and at different joint sites (Doherty et al., 2001). Osteoarthritis can present itself as a symptom free but mechanically altered joint, or symptoms may be experienced but without radiographic change (Dahaghin et al., 2005). More often than not, the natural repair process that takes place in the joint fails to compensate for the damage, and this results in joint failure (NICE, 2008). Therefore, radiographic change only identifies those with severe osteoarthritis and not those present with the clinical symptoms of pain and stiffness (Wenham and Conaghan, 2009). The National Institute for Health and Clinical Excellence (NICE) guidelines state that the presence of radiological change is not required for the diagnosis of osteoarthritis, and the complaint of joint pain and functional limitation is sufficient (Table 3.1). Joint pain is a recommended starting point for population-based studies of osteoarthritis and its impact (Peat et al., 2001). The most commonly affected joints are the knees, hips, and the small joints of the hand (WHO, 2003).

**Table 3.1      The GDP diagnostic criteria for osteoarthritis**

i.	persistent joint pain which is worse with use
ii.	aged 45 years and over
iii.	morning stiffness lasting no more than half hour

**Source:** NICE, 2008

### **3.2.2 Prevalence of osteoarthritis**

The reported prevalence of osteoarthritis in the literature varies due to different definitions of osteoarthritis (e.g. joint pain or radiographic osteoarthritis (Jacobson et al., 2004) (Table 3.2). However, osteoarthritis at individual joint sites (e.g. knee, hip and hand), no matter how it is defined, increases with age (Arthritis and Musculoskeletal Alliance 2005). In population studies, more people report multiple joint pains than pain at a single site (Conaghan and Nelson, 2009).

The prevalence of radiographic osteoarthritis is reported to be higher in women than men, particularly in those aged 50 years and over with hand and/or knee osteoarthritis (WHO, 2003). Radiographic osteoarthritis of the knee affects about a quarter of adults aged 50 years and over in the community (Peat et al., 2001). However the prevalence estimates vary according to the diagnostic criteria and the age of the sample in each study (Table 3.2). Evidence suggests that radiographic osteoarthritis is more common in those who have had the condition longer and have more persistent symptoms. However this also depends on the applied definition (Bedson and Croft, 2008). Duncan et al. (2007) reported that there is a steady association between severity of pain, stiffness and physical function, and the presence of radiographic knee osteoarthritis. Peat et al (2006) suggest that concordance between symptoms and radiographic osteoarthritis seems greater with more advanced structural damage.

Joint pain is common in adults aged 50 years and over in the general population (Thomas et al., 2004:a). The most common site of peripheral joint pain is the knee

(19%; 95% CI 18, 21), and the prevalence of knee pain among women aged 75 and over is estimated to be 35% (defined as pain in the past month lasting for more than a week) (Urwin et al., 1998). However the reported frequencies of joint pain are varied, due to differences in definitions used to measure joint pain (Table 3.2).

**Table 3.2 Standardised prevalence rates of knee osteoarthritis**

Source	Diagnosis	Age	Prevalence
Steven et al 1992	Symptomatic	16-99	6.50
Badley et al 1992	Self-reported	16-99	9.84
Forsberg et al 1992	Radiographic	67-92	53.87
Bagge et al 1992	Radiographic	67-92	43.01
Mannoni et al 2003	Symptomatic	65-99	29.80
Picavet et al 2003	Self-reported	25-99	11.90
Andrianakos et al 2003	Symptomatic/ Radiographic	19-99	6.55
Salaffi et al 2005	Symptomatic	18-91	5.39
Horvath et al 2006	Self-reported pain	14-69	28.30
Tamm et al 2008	Radiographic	34-55	3.74
Fernandez-Lopez et al 2008	Symptomatic	20-99	11.72
Quintana et al 2008:a	Symptomatic/ Radiographic	60-89	71.10
Quintana et al 2008:b	Symptomatic	60-90	40.39
Quintana et al 2008:b	Self-reported	60-90	35.12

**Source:** Data extracted from the Musculoskeletal Health in Europe Report (2012)

Hand pain is also common in older adults in the general population (Dziedzic et al., 2007; Arthritis and Musculoskeletal Alliance (ARMA), 2005; Arthritis Care, 2004; Hochberg et al., 2000; Urwin et al., 1998). One study estimated the prevalence of hand pain as 12% in adults aged 45 years and over (defined as pain in the past month lasting for more than a week) (Urwin et al., 1998). Another study

estimated the prevalence of hand pain and as ~30% in those aged 50 years and over in the general population (participants were asked whether they had experienced any recent pain, specifically *“in the past month any ache or pain that lasted for one day or longer in your hands”*) (Dziedzic et al., 2007). It was suggested that prevalence of hand pain was higher in women (Dziedzic et al., 2007; Dahaghin et al., 2005).

### **3.2.3 The impact of osteoarthritis on self-care in older people**

The impact of osteoarthritis is broad and is associated with many outcomes. The extent of pain severity is associated with increasing mobility problems (Wilkie et al., 2006) and is a predictor of future disability in activities of daily living (ADL) (Davis et al., 1991). According to WHO (2003), about 80% of those with osteoarthritis have limitation in movement and a quarter have limitations in ADL. Studies suggests that joint pain can affect mobility, participation and the psychosocial functioning of the individual, leading to increased dependency in carrying out ADL (Verbrugge and Juarez, 2001; Dunlop et al., 2001; Hochberg et al., 1995). Knee pain was found to be associated with severe difficulties in conducting ADL in those aged 50 years and over (Jinks et al., 2002), and older adults with joint pain are more likely to have participation restriction, than those without (Wilkie et al 2007:a). Reduced quality of life is linked to joint pain through its impact on self-esteem, independence and its considerable effect on personal relationships (Conaghan and Nelson, 2012). Depression is also common amongst those with osteoarthritis, and is linked to activity limitation and participation restriction (Wilkie e al., 2005; Machado et al., 2008).

Studies suggest that personal factors such as psycho-social dynamics and mental health status should be taken into account when attempting to understand the predictors of functional decrease in those with joint pain (Gignac et al., 2000; Yelin et al., 1987). Environmental factors such as social support and exercise are positively associated with protection against poor functional outcome (Sharma et al., 2003). However many studies focus on the individual (i.e. measuring activity limitation) and not the individual's interaction with the environment.

Despite the research indicating links between joint pain and future self-care disability, the extent and mechanisms of restriction in self-care in those with joint pain is fragmented. Differences in the terminologies used to define self-care and joint pain further adds to the variation in research findings. There is a need to conduct a systematic review of the literature to identify whether there are any studies which have specifically explored the links between self-care restriction and joint pain and identified the determinants of self-care restrictions in community-dwelling older adults with joint pain/ osteoarthritis.

### **3.3 Aims and objectives**

The aim of the systematic review was to explore the current literature on restricted self-care in community-dwelling adults aged 50 years and over in the general population and its links with joint pain/ osteoarthritis.

The objectives of the review were to (i) determine if self-care restriction is associated with joint pain in community-dwelling adults aged 50 years and over in

the general population, and (ii) identify the determinants of restricted self-care in community-dwelling adults aged 50 years and over with joint pain/osteoarthritis.

### **3.4 Methods**

#### **3.4.1 Search strategy**

Systematic reviews are conducted to identify relevant studies, synthesise information, and to present an objective summary of the results with due consideration of any limitations in the evidence (Davies and Crombie, 2001). Systematic reviews are formal processes which use explicit methods to perform a thorough literature search and critical appraisal of individual studies to identify the valid and applicable evidence (Moher et al., 2007).

The search strategy was designed to be comprehensive and thorough with the aim of identifying all relevant studies for the research question. Through an eliminatory literature search, keywords were identified to capture the International Classification of Function's (ICF) (WHO, 2001) definition of self-care. A similar approach was taken to identify keywords for 'restriction', such as limitation and disability. In order to include all studies that capture the population aged 50 and over, keywords to represent age groups in numbers and words were used. To ensure the in-depth search of the specialised literature, the search strategy was divided into two stages to meet the reviews objectives. (i) To determine whether self-care restriction was associated with joint pain in community-dwelling adults aged 50 years and over, general population studies were identified. (ii) To explore the determinants of restricted self-care in community-dwelling adults aged 50

years and over with joint pain/osteoarthritis, studies in joint pain/ osteoarthritis populations were identified. For the general population, related terms were also identified through the literature review. For the joint pain population, both medical terms and clinical references were used to gather the studies of joint pain and osteoarthritis. Where necessary, keywords were truncated to yield all results covering varied forms of the word. For example, when a search using the word 'limit' was truncated, this yielded results for the variation of this term such as 'limited' and 'limitation' to ensure that the keywords used in the search were comprehensive.

To capture all studies published to date, titles were retrieved by a two stage computerised search of the Cochrane Library, Medical Literature Analysis and Retrieval System Online (Medline), Excerpta Medica Database (Embase), Cumulative Index to Nursing and Allied Health Literature (CINAHL), King's Fund (Her Majesty's Inspectorate of Constabulary (HMIC)), Allied and Alternative Medicine (Amed), Psychological Information Database (PsychINFO), Applied Social Sciences Index and Abstracts (ASSIA), Ageline-CSA, Social Services Abstracts and British Nursing Index from earliest (1900) to January 2009. Keywords with Medical Subject Headings (MeSH) and text words were used in following order to retrieve the target literature (Table 3.3).

**Table 3.3 Search strategy**

<b>Stage (i) General population</b>	
<b>1</b>	("self care" OR "self-care" OR "self maintenance" OR "self-maintenance" OR "self reliance" OR "self-reliance" OR "personal care" OR "personal-care" OR "activities of daily living" or "daily living activities" or ADL or IADL or PADL or ADLS AND restrict* or limit* or disab*).ti,ab
<b>2</b>	(wash* OR dress* OR "putting on clothes" OR "taking off clothes" OR "putting on footwear" OR "taking off footwear" OR bath* OR shower* OR grooming OR drying OR combing OR shaving OR brushing OR clean* OR walk* OR toilet* OR feeding OR drinking OR "meal preparation" OR "meal-preparation" OR cooking) AND (restrict* or limit* or disab*).ti,ab
<b>3</b>	("older adults" OR "over 50" OR "over fifty" OR "over 65" OR "over sixty five" OR eld* OR age* OR geriatric OR middle age*).ti,ab
<b>4</b>	("community dwelling" OR "community-dwelling" OR "lives at home" OR "lives at-home" OR "lives alone" OR "lives-alone" OR "general population") .ti,ab
<b>5</b>	("joint pain" OR arthrit* OR osteoarthritis* OR osteoarthritis OR OA OR gonarthrosis OR coxarthrosis) AND (pain OR complain*).ti,ab
<b>6</b>	(hip OR knee OR hand OR thumb OR foot) AND (pain) .ti,ab
<b>7</b>	Combine 1 OR 2
<b>8</b>	Combine 3 AND 4
<b>9</b>	Combine 7 AND 8
<b>Stage (ii) Joint pain population</b>	
<b>10</b>	Combine 5 OR 6
<b>11</b>	Combine 3 AND 7
<b>12</b>	Combine 10 AND 11

### 3.4.2 Selection criteria

First of all, the author reviewed the titles of all identified publications for potential relevance to self-care and osteoarthritis. Two observers (Yeliz Prior (YP) and Ross Wilkie (RW)) then independently reviewed the titles of all identified publications for potential relevance to the ICF's definition of self-care used in this thesis. This process was repeated for both stages of the search. Selection criteria were developed to select publications that were relevant to the objectives of this review. Inclusion criteria were:



- i. Method: Epidemiological studies (i.e. cross-sectional, case-control or prospective/cohort)
- ii. Population of interest: Community-dwelling adults aged 50 and over
- iii. Exposure: Joint pain/ Osteoarthritis
- iv. Comparison: General population
- v. Outcome: Limitations/ Restrictions in self-care (i.e. washing oneself, caring for body parts, dressing, eating and drinking and toileting)
- vi. Fully published papers in the English language

Following their review of study titles, the two reviewers (YG and RW) met to discuss which papers were relevant to the objectives of the systematic review. Disagreements were resolved during this consensus meeting. Once agreement on the titles was reached, corresponding abstracts were retrieved and both reviewers independently applied the selection criteria to abstracts, or to full papers when the information provided on abstracts were not sufficient.

Following the review of abstracts of full papers, full articles of all relevant studies were retrieved and the inclusion criteria were applied to these studies independently to ensure they met the systematic review objectives. A further meeting was held by reviewers to discuss the inclusion of these studies and finalise the selection process.

### 3.4.3 Quality assessment and data extraction

Differences in methodological quality across studies can indicate that the results of some studies are more likely to be affected by bias than others. Bias tends to be a much greater problem in observational studies (Bonita et al., 2006). Thus, assessing quality and susceptibility to bias is essential when interpreting research findings and conducting systematic reviews (Sanderson et al., 2007). Bias is defined in epidemiology as *“an error in design or execution of a study, which produce results that are consistently distorted in one direction because of non-random factors”* (Bayona and Olsen, 2004: p15).

There are three major biases in observational studies; confounding, selection and information bias (Hennekens and Buring, 1987). Confounding bias occurs when *“all or part of the apparent association between the exposure and outcome is in fact accounted for by other variables that affect the outcome and are not themselves affected by the exposure”* (Porta, 2008 p:49). Selection bias is a distortion in the estimate of association between the risk factor and disease that results from the methods used during the selection of participants. Selection bias could occur either because the sampling frame is sufficiently different from the target population, or because the sampling procedure cannot be expected to deliver a sample that is a mirror image of the sampling frame (Silman and Macfarlane, 2002). Information bias is an estimate arising from systematic measurement error or misclassification of subjects on one or more variables, risk factor or disease status (Bayona and Olsen, 2004).

Although tools for assessing quality in clinical trials are well-described, notably less attention has been given to tools assessing the quality of observational epidemiological studies (Sanderson et al., 2007). A recent review, considered 121 critical appraisal tools for allied health research, and concluded that there is no agreed 'gold standard'. At the time of this review, no tool has been adopted for widespread use within systematic reviews. Quality assessments in addition to assessing the susceptibility to bias, can also consider issues around sample size, power calculations and ethical approval. However it is important to differentiate between the quality of reporting and the quality of the design, conduct and analysis of a given study (Huwiler-Muntener et al., 2002). A number of consensus statements have encouraged higher quality of reporting, including recommendations for reporting systematic reviews (The Quality of Reporting of Meta-analyses (QUOROM)) (Moher et al., 1999) and observational epidemiological studies (Strengthening the Reporting of Observational studies in Epidemiology (STROBE)) (Von Elm et al., 2007). These tools aimed to assist the author of reports, not the reader who would like to assess the validity of the research (Sanderson et al., 2007). The following assessment tools have been proposed to assess the quality of research studies in peer-reviewed journals.

### Critical Appraisal Skills Programme (CASP)

The Critical Appraisal Skills Programme (CASP) was developed by the Public Health Resource Unit of the National Health Service (NHS) (2006). This tool considered three broad issues; i) are the results of the study valid? ii) what are the results? and iii) will the results help locally? (Bradley and Burls, 1999). The primary

aim was to help the reviewer to think systematically when appraising systematic reviews, thus this tool was not designed to assess individual studies. There are 10 questions in total, of which two are screening questions, advising the reviewer to complete the remaining questions if the answer to both these questions are 'yes'. Reviewers are asked to record a "yes", "no" or "can't tell" to most of the questions, with prompts after each question. Screening questions ask whether there is a clearly focused question; include the right type of study; identify all relevant studies; assessed the quality of the included studies; if the results of the studies combined, was it reasonable to do so; how are the results are presented and what is the main result; how precise are the results; can the results be applied to the local population; were all important outcomes considered, and finally should policy or practice change as a result of the evidence contained in this review? (Public Health Resource Unit, 2008).

#### *Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)*

The STROBE statement was developed by a group of methodologists, researchers, and editors to improve the quality in reporting of observational studies by facilitating critical appraisal and interpretation of studies by reviewers (Malta et al., 2010).

This statement restricts their recommendations mainly to three study designs (cohort, case-control and cross-sectional studies) and provides a checklist of items that should be included in reports of observational studies. This consists of 22 points to summarise the title and abstract, introduction, methods, results and

discussion sections of the studies (Vandenbroucke et al, 2007). The statement makes recommendations for each section. For example, it recommends that the methods of a study should present the key elements of the study design early in the paper and describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection. It provides clear instructions on the need to describe the participants (i.e. for a cross-sectional study, the eligibility criteria, and the sources and methods of selection of participants should be presented); to clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers; give sources of data and details of methods of assessment for each variable of interest; to describe any efforts to address potential sources of bias; to explain how the study size was arrived at; to explain how quantitative variables were handled in the analyses and how the statistical methods followed should be summarised. In the results section, it again provides detailed instructions on how to report the participants at each stage of the study; to give characteristics of study participants (e.g. demographic, clinical, social) and information on exposures and potential confounders; to report numbers of outcome events or summary measures; to give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (e.g., 95% confidence interval), and report other analyses done (e.g. analyses of subgroups and interactions, and sensitivity analyses). The Strobe Statement also gives pointers for how to report the discussion and any other information such as funding (Sanderson et al., 2007). The STROBE group stress that the aim is to provide guidance on how to report observational research well, and that the recommendations are not prescriptions for designing or conducting studies. They

also add that the checklist is not an instrument to evaluate the quality of observational research (The Strobe Initiative, 2009) (Table 3.4).

**Table 3.4 STROBE Statement—Checklist of items that should be included in reports of cross-sectional studies (an extract)**

Item no		
<b>Title and abstract</b>	<b>1</b>	<b>Recommendation</b> a) Indicate the study's design with a commonly used term in the title or the abstract b) Provide in the abstract an informative and balanced summary of what was done and what was found
<b>Introduction</b>		
Background and Rationale	<b>2</b>	Explain the scientific background and rationale for the investigation being reported
Objectives	<b>3</b>	State specific objectives, including any pre-specified hypotheses
<b>Methods</b>		
Study design	<b>4</b>	Present key elements of study design early in the paper
Setting	<b>5</b>	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	<b>6</b>	Give the eligibility criteria, and the sources and methods of selection of Participants

**Source:** The Strobe Statement, 2009

#### Quality in Prognosis Studies (QUIPS-LBP) tool

The QUIPS-LBP tool was designed for systematic reviews of prognostic studies through international expert consensus (Hayden et al., 2006). Originally developed to provide criteria and guidance for assessing the risk of potential biases in low back pain prognosis studies, this tool considers bias related to study participation,

study attrition, prognostic factor measurement, outcome measurement, confounding measurement and account bias related to analysis.

The QUIPS-LBP does not offer a scoring system to appraise studies. Instead, it allows the overall quality of the research to be comprehensively evaluated by recognising flaws in the study which may have resulted in bias (Wynne-Jones et al., 2010; Hayden et al., 2008). The reviewer is asked to consider if the study methods satisfy each general statement for the six biases assessed by the tool and explain why they chose their rating under 'comments' (Table 3.5). There are six sections and 36 questions with five possible ratings as: "Yes", "Partly", "No", "Unclear" and "Not relevant" with a summary at the end of every section. Where the majority of domains are given a "no" rating this suggests that the methodological quality of that domain is poor. Where the majority of domains are given a rating of "unclear" or "partly", this indicates a moderate risk of bias, which is difficult to assess due to the lack of detail with regards to the methods reported in the study. Finally, the "yes" rating indicates a low risk of bias, suggesting that the reporting of the methodological quality is sufficient to exclude the possibility of bias in the study (see appendix). Although originally developed for back pain studies, the QUIPS-LBP tool could be used to assess bias for other conditions (Hayden et al., 2007).

**Table 3.5      The layout of the QUIPS-LBP: Quality Appraisal Exercise (an extract)**

Study (First author, year):	
Bias related to Study Participation	
1. The source of population or population of interest is adequately described for key characteristics.	<b>Comments:</b> For example, what are the 'key characteristics; that need to be described for the source population?
<input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	

**Source:** Data from the original QUIPS Quality Appraisal Exercise (Hayden et al., 2006)

### The Quality Assessment Process

The most suitable tool to assess the quality of studies was the QUIPS-LBP tool (Hayden et al., 2007) (Table 3.5). This was both due to the use of expert consensus in development of this tool and the ease of practical application as it listed a structured series of questions to identify and score possible bias and confounding (Hayden et al., 2007). The QUIPS-LBP tool was adapted for use in this study by making minor adjustments to allow application to both cross-sectional and longitudinal studies (e.g. question 13 asks whether the study provides a clear description of the prognostic factor measured – including dose, level, duration of exposure, and clear specification of the method of measurement. This heading was changed to 'factors measured', and the section on the study attrition was not applied to cross-sectional studies).

Two reviewers (YG, RW) independently assessed the quality of each study using the QUIPS-LBP (Quality in Prognosis Studies) tool. Each study was given



individual scores at the end of the assessment in the overall summary section. A third reviewer was consulted to resolve any disagreements (Roger Beech (RB)). A descriptive summary of quality assessment was presented for each study according to the QUIPS quality appraisal exercise tool in the results section (Table 3.9).

#### Data extraction

Each study's detail on the study population (setting, sampling frame, sample size, response rate), outcome and the associations with other factors were extracted and presented as a narrative summary table in the results section (Table 3.8).

#### **3.4.4 Strength of the evidence**

To establish possible associations between joint pain and restricted self-care, important factors such as temporal relationship (i.e. does joint pain occur before activity limitations), plausibility, dose-response relationship (i.e. does increased pain results in increased restriction) and the consistency of possible associations across studies were considered. It was understood that the strength of these associations might be dependent on the relative prevalence of other possible causes (Bonita et al., 2006), thus the evidence acquired for or against the cause and effect relationship might be discordant. Nevertheless following criteria ensures that the systematic review maintains a methodical manner (Van Der Windt et al., 2000).

## 3.5 Results

### 3.5.1 Search Strategy

#### Search 1: Self-care and joint pain in the general population

The number of citations identified in each database was documented in “Reference Manager” (Table 3.6 and Table 3.7) and each reference was archived electronically. In total 9765 citations were identified and retrieved during the first stage of the search. 9731 citations did not meet the inclusion criteria as they were not relevant to the review (e.g. search terms yielded results from studies of brain injury, psychological disorders and other literature as well as studies of self-management of health conditions). Two reviewers read the remaining 34 abstracts and 19 abstracts were excluded for not meeting the inclusion criteria. The remaining 15 papers were retrieved and read by two reviewers and further 13 papers were excluded in this stage because they did not meet the inclusion criteria (i.e. either because of the study design (i.e. not an epidemiological study), definition of self-care (e.g. inclusion of mobility or IADL items) and joint pain (i.e. not all musculoskeletal pain is due to OA) or differences in sampling frames (i.e. not meeting the criteria of aged 50≥). Two papers were identified for review.


#### Search 2: Self-care in the joint pain population

During the second stage of the search, 988 citations were retrieved and 973 citations were excluded for not meeting the inclusion criteria. Of the remaining 15 abstracts, 14 abstracts were excluded for not meeting the inclusion criteria, leaving


one paper. The flow diagram in figure 3.1 describes the number of abstracts accepted and rejected by two reviewers during the selection procedure. A total of three studies were finally included in the review.

**Figure 3.1** Flow diagram of papers accepted and rejected in the review

**Search 1. General population**



**Search 2. Joint Pain population**



**Table 3.6 Search 1: General population**

Date of Search	Database	Host/System Used	Years Searched	Numbers of documents obtained	Number of Duplicates	Total after removal of duplicates
15.12.2008	Cochrane Library	Wiley-Interscience	All years	396	0	396
15.12.2008	ASSIA	CSA-Illumina	All years	172	0	568
15.12.2008	Ageline	CSA-Illumina	All years	1685	1	2253
16.12.2008	Social Services Abstracts	CSA-Illumina	All years	753	2	3004
17.12.2008	Embase	NHS Library	All years	2620	278	5346
17.12.2008	HMIC	NHS Library	All years	83	29	5400
08.01.2009	Medline	CSA-Illumina	All years	4364	2326	7438
07.01.2009	CINAHL	NHS Library	All years	2581	432	9587
07.01.2009	Amed	NHS Library	All years	233	154	9666
07.01.2009	Psychinfo	NHS Library	All years	979	92	9758
07.01.2009	British Nursing Index	EPSCO	All years	15	7	9765

**Table 3.7 Search 2: Joint pain population**

Date of Search	Database	Host/System Used	Years Searched	Numbers of documents obtained	Number of Duplicates	Total after removal of duplicates
26.01.2009	Cochrane Library	Wiley-Interscience	All years	291	0	291
15.12.2008	ASSIA	CSA-Illumina	All years	66	0	357
15.12.2008	Ageline	CSA-Illumina	All years	24	0	381
16.12.2008	Social Services Abstracts	CSA-Illumina	All years	319	1	699
17.12.2008	Embase	NHS Library	All years	94	3	790
17.12.2008	HMIC	NHS Library	All years	2	0	792
08.01.2009	Medline	CSA-Illumina	All years	156	37	911
07.01.2009	CINAHL	NHS Library	All years	80	28	963
07.01.2009	Amed	NHS Library	All years	16	12	967
07.01.2009	Psychinfo	NHS Library	All years	14	9	972
07.01.2009	British Nursing Index	EPSCO	All years	19	3	988

### 3.5.2 Articles Identified

The three articles selected for the review are summarised below;

(i) General population studies:

#### *Effect of Arthritis in Middle Age on Older-Age Functioning (Covinsky et al., 2008)*

This study aimed to examine whether symptomatic arthritis in middle age predicts early onset of functional difficulties that are associated with loss of independence in older adults (Covinsky et al., 2008). Participants were those who took part in the 1992 Health and Retirement Study (HRS), as they were considered to be an ideal cohort to address whether arthritis in mid-life leads to earlier onset of late-life disabilities. The study design was a prospective longitudinal study. 7543 participants aged 50 to 62 at baseline, reporting no difficulty in mobility or ADL function were enrolled in the study (Covinsky et al., 2008).

This study looked at the links with symptomatic arthritis defined by self-reported pain, stiffness or swelling in joints, use of medications or other treatments for their arthritis or rheumatism, or consultation with a doctor for arthritis in the previous 12 months. The primary outcome was time to persistent difficulty in one of five activities of daily life (ADLs) (bathing, dressing, transferring from a bed to a chair or out of chair, using a toilet, or eating) or mobility. Participants were classified as having difficulty with ADLs if they reported difficulty in one of these items. Difficulty with carrying out ADLs and mobility were assessed by interview every two years. Participants were classified as having persistent difficulty only if they reported a specific difficulty on two consecutive waves (participants who reported difficulty on one wave and died before the next wave was also classified as having persistent

difficulty). Kaplan-Meier curves were constructed to compare the time to persistent difficulty with mobility or ADL function in participants with or without arthritis at baseline. Proportional hazards survival analysis was used to calculate the unadjusted and adjusted hazard ratios for the association between arthritis and subsequent mobility or ADL difficulty. Analyses were adjusted for other comorbid conditions, body mass index, exercise, and demographic characteristics to determine independent associations between arthritis and mobility or ADL difficulty (Covinsky et al., 2008).

At baseline 29% of the participants had symptomatic arthritis. Participants with symptomatic arthritis were older, more likely to be female, of low socio-economic status and higher BMI than those who did not report arthritis. They were also more likely to have other comorbidities such as hypertension, diabetes mellitus, cancer, lung disease, and depressive symptoms. They were also more likely to have persistent difficulty in mobility and ADL function across the 10 year period than those who did not have symptomatic arthritis (Covinsky et al., 2008). The authors conclude that middle-aged adults with symptomatic arthritis had a greater risk for developing mobility and ADL difficulties that lead to loss of independence in late life (Covinsky et al., 2008).

#### *Arthritis Prevalence and Activity Limitations in Older Adults (Dunlop et al., 2001)*

This study aimed to evaluate the prevalence of arthritis and activity limitations among older Americans by assessing their demographic, ethnic, and economic characteristics. Data was utilised from the Asset and Health Dynamic Survey

Among the Oldest Old (AHEAD) study, a national probability sample of community-dwelling adults born before 1924, and analysed cross-sectionally (Dunlop et al., 2001).

The case definition of arthritis was defined as symptomatic arthritis that has been severe enough for participants to visit a doctor within the last 12 months prior to this study. The study compared the activity limitation experience of people with arthritis, with people reporting other conditions. Activity limitations included self-reported functional limitations in basic ADL or instrumental ADL (IADL) tasks and physical activity limitations. ADL limitations were having difficulty or receiving help in the tasks of dressing, toileting, bathing, or eating, and reports of using a device, having difficulty, or receiving help in the tasks of walking across a room or transferring in and out of bed. IADL limitations were defined as difficulties or receiving help with hot meal preparation, shopping, using the telephone, taking medication, or managing money. Functional limitations were categorised as no limitations, or limitations in only IADL, one to two ADL, or three or more ADL. Physical activity limitations were related to having difficulties in walking several blocks, climbing one flight of stairs, pulling / pushing heavy objects, carrying ten pounds, or picking up a dime (Dunlop et al., 2001).

All analyses were weighted and adjusted for the complex sampling design to provide appropriate statistical tests based on the sample size to reflect the probability sample of the national population. Logistic regression was used to calculate adjusted rates for selected population characteristics. The estimated logit



from these analyses was used to obtain estimates of adjusted rates and proportions. The standard error of the adjusted rate, used to calculate the associated confidence interval, was estimated by delta method, and statistical testing was done at a nominal 5% alpha significance level (Dunlop et al., 2001).

The prevalence of arthritis in older people ranged from 25% in non-Hispanic whites to 40% in non-Hispanic blacks and 44% in Hispanics. Increased prevalence of arthritis was associated with less education, lower income and less wealth. The prevalence of ADL limitations in those with arthritis was 29%, 30% and 37% respectively for non-Hispanic whites, non-Hispanic blacks and Hispanics, and increased to 48%, 57% and 56% in those with arthritis who also report other chronic conditions (Dunlop et al., 2001). The authors concluded that non-Hispanic blacks and Hispanic older adults reported higher rates of arthritis than non-Hispanic whites, and Hispanics reported higher frequency of ADL limitations than non-Hispanic whites (Dunlop et al., 2001).

#### (ii) Joint pain study

##### *Longitudinal Study of Joint Pain in Older People (Donald and Foy, 2004)*

This study aimed to describe the natural history of joint pain in older people. The study recruited participants as part of a randomised controlled trial (RCT) evaluating the benefit of targeted nurse visits. 4804 participants aged 75 years and over from 12 UK General Practices (GP) were enrolled in the study. Participants were followed for a year in 1998 (Donald and Foy, 2004).

The postal version of the Elderly At Risk Rating Scale (EARRS), which consists of 20 domains to include activity and mobility, personal ADLs, mental health, support at home and questions to include joint pain, were sent to all participants identified by nurses at the 12 GP practices. EARRS was originally designed for nurse interview, thus it required minor modifications. The presence of joint pain was measured by recording the participant's response to the question "regarding joint pains, do you i) have no trouble at all? ii) have only occasional pains? iii) have pain some of the time? iv) have pains a lot of the time? v) have pains virtually all of the time? (Donald and Foy, 2004). This was categorised to 'no pain' (no or occasional pain); "episodic" pain some of the time and "constant pain" for pain 'a lot' or 'all' of the time. Multiple regression modelling was used to examine for associations (Donald and Foy, 2004).

Joint pain was reported by 83% of the participants. The prevalence of constant pain was 26% and was higher in women and in those over the age of 85. Pain increased with age in women. Joint pain was associated with disability and dependency across a wide range of problems. Pain was associated with a threefold increase in likelihood of dependency for ADL. Over 1 year, 18% acquired or had increased frequency of pain and 14% had reduced frequency of pain. Resolution was associated with preserved indoor mobility, and functional recovery (Donald and Foy, 2004). The authors concluded that joint pain is very common in older people, fluctuates over time, and strongly associated with psychological factors and disability (Donald and Foy, 2004).

**Table 3.8 Data extraction from the identified articles**

<b>STUDIES</b>	<b>Effect of Arthritis in Middle Age on Older Age Functioning</b>	<b>Arthritis Prevalence and Activity Limitations in Older Adults</b>	<b>A Longitudinal Study of Joint Pain in Older People</b>
Publication Year and Journal	2008. 56:23-28. Journal of American Geriatric Society	Jan 2001. Vol. 44, No. 1, pp 12-221. American College of Arthritis & Rheumatology	June 2004. 43:1256-1260, Journal of Rheumatology
Authors	Covinsky, K.E., Lindquist, K., Dunlop, D.D., Gill, T.M., Yelin, E.	Dunlop, D.D., Manheim, L.M., Song, J., Rowland, W.C.	Donald, I.P., and Foy, C.
Methods	Prospective Longitudinal Study	Cross-sectional	Prospective Longitudinal Study
Definition of self-care	Participants were classified as having difficulty with ADLs if they reported difficulty with bathing, dressing, transferring from a bed to a chair or out of a chair, using a toilet, or eating.	ADL limitations included reports of having difficulty or receiving help in the tasks of dressing, toileting, bathing, or eating, and reports of using a device, having difficulty, or receiving help in the tasks of walking across a room or transferring in and out of bed.	Personal activities of daily living was assessed in personal hygiene and dressing dependency
Definition of arthritis	Participants were asked, "Have you ever had, or has a doctor ever told you that you have, arthritis or rheumatism?"	A person was considered to have arthritis if an affirmative response was given regarding seeing a doctor within 12 months for arthritis or rheumatism or if a person reported a joint replacement that was not associated with a hip fracture.	Cases were identified by the response to the question; Regarding joint pains, do you (i) have no trouble at all (ii) have only occasional pains (iii) have pains some of the time (iv) have pains a lot of the time (v) have pains virtually all of the time

<b>STUDIES</b>	<b>Effect of Arthritis in Middle Age on Older Age Functioning</b>	<b>Arthritis Prevalence and Activity Limitations in Older Adults</b>	<b>A Longitudinal Study of Joint Pain in Older People</b>
Setting	The Health and Retirement Study, a nationally representative sample of people aged 50 to 62 in the United States at baseline who were followed for 10 years.	Data were based on the AHEAD (Asset and Health Dynamic Survey Among the Oldest Old), a national probability sample of older Americans living in the community.	12 general practice populations in UK, with predominance of rural towns. All the practices had less than 5% ethnic minority on their list.
Sampling Frame	Participants enrolled in 1992 in the first wave of HRS who reported no difficulty in mobility or ADL function in baseline were used.	People born 1923 or earlier (70 years or older) who participated in 1993 AHEAD study.	All patients aged 75 and over on 1 May 1998 in 12 practices were identified and sent the postal version of Elderly At Risk Rating Scale (EARRS)
Sample Size	n= 7,543	n= 7,447	n= 4,804
Response Rate	80%	80%	80%
Mean Age	56 years	77 years	81 years
Female	51%	62%	58%
Prevalence of self-care restriction	Not reported	29%- Non-Hispanic Whites 30%- Non-Hispanic Blacks 37%- Hispanics (those with arthritis)	Not reported
Prevalence of Joint Pain/ OA	29%	24%- Non-Hispanic Whites 40%- Non-Hispanic Blacks 44%- Hispanics	83%

STUDIES	Effect of Arthritis in Middle Age on Older Age Functioning	Arthritis Prevalence and Activity Limitations in Older Adults	A Longitudinal Study of Joint Pain in Older People
Key Findings	<p>Participants with arthritis were more likely to reach the primary outcome measure of persistent difficulty with mobility or with ADL function (34% vs 18%). It was found that middle-aged persons with arthritis were at higher risk for developing mobility and ADL difficulties that lead to loss of independence in late life.</p>	<p>Ethnic differences were found in the arthritis prevalence rates among older adults. Arthritis prevalence rates among non-Hispanic blacks and Hispanics were more than 50% higher than among non-Hispanic whites. Across all ethnic groups, the rates of ADL limitations increased with the presence of arthritis and other conditions, with the highest rates of ADL limitations being among older people with arthritis and other conditions.</p>	<p>The prevalence of constant pain was 26% and higher in women. Disability, measured in EARRS questionnaire, was found to be far stronger predictor of adverse outcomes (what is the outcome) than joint pain.</p>

### 3.5.3 Methodological quality

The results of the quality appraisals are presented in the Table 3.9. This table shows an overall rating for each study to summarise the rating of individual items under the six main sections of the QUIPS-LBP assessment.

**Table 3.9 Overall summaries of the QUIPS-LBP ratings for each study**

QUALITY IN PROGNOSIS STUDIES (QUIPS-LBP): QUALITY APPRAISAL				
		Effect of Arthritis in Middle Age on Older Age Functioning	A Longitudinal Study of Joint Pain in Older People	Arthritis Prevalence and Activity Limitation in Older Adults
		OVERALL RATINGS FOR EACH SECTION		
1	Bias related to study participation	partly	unclear	partly
2	Bias related to study attrition	unclear	unclear	unclear
3	Bias related to prognostic factor measurement	yes	yes	partly
4	Bias related to outcome measurement	yes	yes	yes
5	Bias related to confounding measurement and account	yes	partly	partly
6	Bias related to analysis	partly	partly	partly

### *Effect of Arthritis in Middle Age on Older-Age Functioning*

The reviewers agreed that this study did not describe the sampling frame and recruitment clearly, expecting readers to be familiar with the HRS sampling strategy. Baseline participants were recruited on the basis of not reporting any difficulty in mobility or ADL function. However, environmental factors such as use of aids and adaptations were not measured. This could potentially have influenced the absence or presence of ADL difficulties reported. Participants were interviewed every two years to assess these difficulties and participants were classified as having persistent difficulty if they reported difficulty on two consecutive waves. This two-year gap between interviews and the definition of 'persistence' could be problematic due to participants' readiness to report symptoms of disability. There were no clear outlines of the interview process, which could be an important source of bias, raising questions about the validity and reliability of this outcome measure.

The study provides data for the first two years with regards to response rate and study attrition. However, the sample was followed for 10 years. This implies that the number of responders had not changed in the subsequent years, which may be questionable. Additionally, participants who were lost at follow-up were not clearly described. Discrepancies were noted with regards to censoring exercise, as whether they have used partial data on censored participants or not was not clearly stated.

### Arthritis Prevalence and Activity Limitations in Older Adults

Reviewers agreed that the inclusion and exclusion criteria for the study participation were not adequately described in this study. It was not clear in the paper that the study population was the correct representation of the target population. The term 'community-dwelling adults' was not clearly defined. It has been noted that some studies use this term for people living in supported housing schemes.

The study did not provide enough information for the characteristics of those, who were lost to follow-up and/or non-responders to judge whether the remaining responders were a representative sample of older adults in the general population. It was difficult to identify whether an adequate proportion of the sample had complete data for all relevant outcome periods, as there was no mention of missing data. In the statistical analysis, figures were weighted and adjusted, suggesting there may have been some imputation. However, there was no mention of this in the study. The conceptual framework within the Arthritis Prevalence and Activity Limitations in Older Adults study drove an examination of the differences in limitation with ethnicity but the results indicated that socio-economic factors may also be the predictor of the outcome.

### Longitudinal Study of Joint Pain in Older People

The reporting of the study design was complex due to references made to the randomised controlled trial (RCT) evaluating the benefit of targeted nurse visits. It was stated that all patients over 75 years of age in twelve practices were enrolled



in this study. However, the levels of joint pain or the problems that these participants were having were not described. Data were collected using EARRS (Donald, 1997) but the conceptual basis for the inclusion of this assessment was not established within the study. EARRS was not designed for a postal survey. The study suggested that the results for the nurse interview versus postal questionnaire were tested on feedback from 30 patients residing in a day care hospital, indicating good overall agreement. However this was a relatively small sample, derived from hospital patients, compared to 4804 general population sample, which subsequently carried out this postal questionnaire as a self-assessment. Again, it is questionable whether this small sample was representative of the general population of older adults living in the community.

The study stated that no specific instructions were included within the questionnaire regarding whether to include or exclude back pain, suggesting that participants were left to identify the difference between joint pain and back pain. Therefore, a potential assumption may be that this sample will include those with back pain, and this may have increased the prevalence of joint pain as well as strengthening associations with other factors.

Important confounders besides age and gender were not included in this study. The attempts to collect information on dropouts were not described. Results were presented without supporting data and, an important proportion of the information seems to be missing.

#### **3.5.4 The association between self-care restriction and joint pain in community-dwelling older people**

This systematic review involved a thorough search of the literature to identify observational studies that explored the relationship between limitation in self-care activities and joint pain in community-dwelling older adults in the general population. Despite finding a number of studies that have reported the prevalence, onset, persistence and other factors associated with ADL limitations in community-dwelling older adults in the general population (e.g. Gill et al., 2007:a-b, 2006:a-b, 2003, 2002:a-b, 1998; Hardy and Gill, 2004; Naik et al., 2004), and studies that have examined associations between joint pain and poor functional outcomes (e.g. Wood et al., 2008; Machado et al, 2008; Grothle et al., 2008; Dziedzic et al., 2007; Corti and Rigon, 2003; Peat et al, 2001), there were only two studies which specifically examined the relationship between limitation in self-care activities and joint pain in community-dwelling older adults in the general population.

As highlighted earlier in this chapter, the definition of osteoarthritis and self-care restrictions varies across studies, and this was observed whilst doing this study. This further evidenced in the two identified studies in which definitions of self-care problems (i.e. how they defined limitation / disability) and joint pain differed. Covinsky and colleagues (2008) study considered transferring from a bed or chair or out of chair as one of the five ADLs, and Donald and Foy (2004) considered dressing dependency and personal hygiene only. Whilst one study made no distinctions between OA and rheumatoid arthritis, and defined symptomatic arthritis as pain, stiffness and swelling of joints (Covinsky et al., 2008), the other study measured the frequency of joint pain as no pain, episodic, and constant pain

(Donald and Foy, 2004). Notably the study population in Donald and Foy's study was much older than the other ( $50 \geq CF 75 \geq$ ).

Donald and Foy's Longitudinal Study of Joint Pain (2004) found that joint pain fluctuates in frequency over time and is strongly related to disability as well as psychological factors. One in five adults with constant pain were housebound compared to 6% in those without pain. Pain was associated with a threefold increased likelihood of dependency for ADL. In examining the relapsing and remitting nature of joint pain, sadness and dressing dependency were found to be predictors of acquiring joint pain, which was suggested that these may be early markers of functional decline, which frequently accompanied new joint pain (Donald and Foy, 2004).

Although these studies adjusted the associations between self-care activities and joint pain for various other factors such as age and gender, psychological and other health and socio-demographic factors, there was no clear rationale or consistency behind choosing these individual factors across the two studies. These studies measured self-care as difficulty (i.e. at the level of the individual) and haven't considered self-care as a biopsychosocial construct. Thus, following this review, the question asked at the beginning (i.e. "are people with joint pain more likely to have self-care restriction?") was still not answered.

### **3.5.5 Determinants of self-care restrictions in older adults with joint pain**

The systematic review identified one study that specifically examined the determinants of self-care restrictions in community-dwelling older adults with joint pain. However, this study defined self-care as an ADL; it measured limitation in dressing dependency and personal hygiene only, which is at the level of an individual (i.e. environmental factors were not considered). Although this study offers some insight into the relationship between joint pain and self-care disability in older people, it does not clearly identify the determinants of self-care restrictions in older people. An approach that considers the many potential individual and contextual determinants may provide a better understanding of self-care restriction.

### **3.5.6 Summary of results**

Following a two stage comprehensive systematic search of the literature through eleven electronic databases to identify observational epidemiological studies investigating associations between self-care restrictions and joint pain in community-dwelling older adults in the general population and the determinants of self-care restrictions in older adults with joint pain, only three studies were included in the review. The overall methodological quality of these studies ranged from being moderate to low risk of bias.

The conclusion from the systematic review is that evidence of the extent and mechanisms of self-care restriction in community-dwelling older adults remains limited to date. Information on self-care in older adults in the general population is

fragmented. Limitation in specific self-care tasks (based on an individual's capacity) was associated with joint pain, but there were no studies which have focused on self-care as an outcome of the individual and environmental factors. There were no studies which identified the determinants of self-care restriction in older adults with joint pain.

### **3.6 Discussion**

Despite the comprehensive search of the literature, this review was only able to identify three relevant epidemiological studies, indicating that links between self-care restrictions and joint pain were an under-researched area. However, the limited number of studies identified could be due to the variety of terms used to describe self-care restrictions in the literature, and however carefully selected, the keywords used in this review to find these studies may have not been exhaustive. The inclusion and exclusion criteria were strictly applied to identify only observational studies that referred to self-care. This review found that measurement of self-care in studies varied with some self-care tasks being measured.

The main aim behind this review, which was to identify studies which could describe and explain the relationship between self-care and joint pain in community-dwelling older adults, may have potentially narrowed the results. The fact that only epidemiological studies were included in this review may have led to the exclusion of a substantial qualitative study, thereby impacting on the findings. Although qualitative studies cannot provide quantitative estimations of the

prevalence and patterns of such cases, they may identify important factors that contribute to the occurrence of self-care restrictions through exploration of the view and experiences of older adults with joint pain. Equally, there may have been studies conducted in languages other than English. As English was a selection criterion, this could have led to the exclusion of studies conducted in other languages. However there is little evidence that not including non-English language research results in biased estimates (Pham et al., 2005).

The search in this review included 11 bibliographic databases from the time of their inception until January 2009. However the search did not cover any grey literature or contact any local experts with regards to unpublished studies or thesis. At the time this thesis is written, there may be other studies of self-care restriction and joint pain conducted, and / or in the process of being peer reviewed for publishing, which may have had an impact on the results of this review.

This review utilised a standardised assessment tool to review studies to aid consistency of reviewers' observations and reduce the inter-rater bias. Inter-rater bias refers to the consistency in which multiple reviewers evaluate the same data using the same scoring criteria (Bailey, 1998) at a specified time (Stemler, 2004). This concept suggests that the behaviour of reviewers contains a degree of subjectivity, thus the judgements made will depend upon the reviewers' interpretation of the criteria. A good quality assessment aims to reduce this subjectivity by providing clear instructions (Tierney and Marielle 2004; Mertler, 2001). The quality assessments of these studies were individually and

independently carried out using the QUIPS-LBP tool (Hayden, 2007) by three reviewers (YG, RW and RB). Prior to the full review, one paper was selected to facilitate a consistent approach to the review and judgement of the different forms of bias. During the consensus meetings, reviewers 'explained' their choice of particular rating in the comments section; differences in opinions were explored, and agreement on ratings was achieved. The QUIPS-LBP tool was developed through international expert consensus, and aim to provide a criteria and guidance for assessing the risk of potential biases in low back pain prognosis studies (Hayden, 2007). However, the use of such a tool could be controversial and, in the absence of empirical evidence to support the 'quality' in observational studies, the assumptions behind equal weighting of scores could be deemed as questionable (Mallen et al., 2007).

The methods used in the three studies varied in terms of study setting, sample frames, outcome measures, and use of statistical analysis. Therefore, there isn't clear empirical evidence of an association between self-care restriction and joint pain or of the potential determinants of self-care restriction in older adults with joint pain (Table 3.10).

### **3.7 Conclusion**

There have been a number of studies that have described the impact of osteoarthritis in older adults. Despite this, the systematic search described in this chapter identified two papers that empirically examined the link between self-care and joint pain. Only one study was identified that explored self-care in older adults

with joint pain. Following this review, it is still unclear if self-care restriction is associated with joint pain. The frequency of self-care restriction and its potential determinants in older adults is also unclear.

A barrier to research on self-care is the inconsistent approach to definitions and measurement. The ICF offers an opportunity for a consistent approach to self-care research through the proposal of a standard definition and framework. It also offers a broader approach which views self-care within a biopsychosocial model where restriction is not only dependent on joint pain and physical function but also on psychosocial and environmental factors. As the conceptual model of restrictions in self-care restriction suggests that the potential determinants may be multi-factorial, the links with comorbidities, impairments, activity limitations, socio-economic and environmental factors should be taken into account.

There is a need to investigate the prevalence and patterns of restricted self-care in older adults in the general population, the links with joint pain and the potential determinants in older adults with joint pain. Using the ICF to define self-care and provide a framework to classify potential determinants, this thesis describes an observational study in community-dwelling older adults.



**Table 3.10     Summary of key findings**

- Although a number of studies described the impact of joint pain in older adults, only two studies examined the link between self-care and joint pain empirically.
- Only one study explored self-care restrictions in older adults with joint pain.
- Following this review, it is still unclear if self-care restriction is associated with joint pain. The frequency of self-care restriction and its potential determinants in older adults is also unclear.
- The inconsistent approach to definitions and measurement of self-care creates a barrier for research.
- The ICF offers an opportunity for a consistent approach to self-care research though the proposal of a standard definition and framework, providing a biopsychosocial model where restriction is not only dependent on joint pain and physical functions but also on psychosocial and environmental factors.
- There is a need to investigate the prevalence and patterns of restricted self-care in older adults in the general population, the links with joint pain and the potential determinants in older adults with joint pain.

## **Chapter 4**

### **North Staffordshire Osteoarthritis Project (NorStOP): The survey overview and sample analysis**

#### **4.1 Introduction**

Each study described in this thesis uses data collected in the North Staffordshire Osteoarthritis Project (NorStOP); a population based prospective cohort study which collected information using postal questionnaires. Cohort, cross sectional, and case-control studies are often referred to as observational studies as the data is collected through observations, and no interventions performed (Mann, 2003). Cohort studies are useful for investigating prevalence, incidence, cause and prognosis of health outcomes in populations observed over time (Mann, 2003, Schlesselman, 1982). Cohort studies can provide data on potential causes with data being collected prior to the outcome (Mann, 2003). However, a key limitation of cohort studies is loss to follow-up, particularly when the intervals between follow-ups is over a long periods of time, which can introduce bias to the study findings.

Secondary analysis involves the use of an existing dataset to address new research questions. It is a popular way to investigate research questions when resources (such as time and funding) restrict new primary data collections (Clarke and Cossette, 2000). It is regarded as an efficient way of producing data to support population health (Bibbs, 2007). This chapter will examine the suitability of the NorStOP dataset to meet the objectives of this thesis and answer the main

study questions by reviewing the study design, the psychometric properties of the data collected, and the sample derivation.

## **4.2 Study design**

### **4.2.1 Setting and sampling**

The aim of NorStOP was to study the impact of joint pain / osteoarthritis in adults aged 50 years and over in the general population (Thomas et al., 2004:b). The main objectives were to “(i) *determine the impact these syndromes have on activity and participation levels in older people, (ii) to determine factors which predict prognosis over time with respect to change in pain, activity and participation, and (iii) to determine frequency and predictors of health care use by sufferers of these syndromes through prospective linkage of self-completion survey instruments with primary care records*” (Thomas et al, 2004-b: p.7).

NorStOP recruitment was divided into three sub-cohorts (NorStOP<sup>1</sup>, NorStOP<sup>2</sup>, and NorStOP<sup>3</sup>). NorStOP<sup>1</sup> was set up as a population study only; NorStOP<sup>2</sup> and NorStOP<sup>3</sup> were organised to provide a sampling-base for a clinical epidemiology study of knee pain (CAS-K; Peat et al., 2004) and the hand pain and problems respectively (CAS-HA; Myers et al., 2007). This thesis uses data collected in NorStOP<sup>1</sup> only.

To obtain a representative sample of older adults, NorStOP recruited patients from general practice (GP) registers. In the UK, 98% of the British population are registered with a GP (Bowling, 1997). At the time of this study there were 101

general practices in North Staffordshire and 16 of these practices formed the General Practice Research Network. Those practices taking part in this study were fully computerised. All practice staff required to enter a code for each contact as part of READ classification of morbidity and event coding, which is widely implemented within the National Health Service. The READ classification was originally developed by Dr James Read (Read, 1991), and it was structured similarly to the International Classification of Diseases. The data kept in these practices were also audited annually to assess the quality and completeness of the data. Audits were undertaken by the Primary Care Sciences Research Centre Health Informatics team (Wilkie, 2005).

At baseline, three practices from the North Staffordshire Research Consortium were recruited to the study. These three practices were located in Stoke-on-Trent and one in Newcastle-under-Lyme, both urban areas, which are rated as some of the poorest in England (Experian, 2012). Contact details of all adults aged 50 years and over were taken from each research practice, and were checked by GPs from the practices for exclusions (n=11309). These exclusions were; (i) patient is unable to complete the questionnaire due to illness; (ii) patient is known to have severe learning disabilities or a severe psychological disorder; and (iii) the patient has indicated previously that they would not like to take part in research projects.

The baseline phase of the study was a two-stage mailed survey; firstly the participants were mailed the "Health Survey Questionnaire", which collected

information on socio-demographics, general health, physical function, participation, and bodily pain. Following this, those consenting to further contact and reported pain and/or problems in the hand, hip, knee or foot in the last year were mailed the “Regional Pains Survey Questionnaire”, which focused on these four joint areas. At both stages, questionnaires were mailed with a letter from the GP practice and a study information leaflet, with reminders to be sent to non-responders after two and four weeks (Thomas et al., 2004: b).

Throughout the mailing process the Keele General Practice Research Partnership (KGPRP) staff maintained the checks on the database for deaths and departures. Details such as date of birth and gender given on the returned questionnaires were checked against the GP records to ensure that correct person had completed the questionnaire (Wilkie, 2005; Muller, 2010). If a date of birth on a questionnaire differed to the data on the GP records, the information given by the participant was used as the correct data once it was established that the correct person had completed the questionnaire. The gender of the participants was checked against the name of the respondent in the GP lists if the NorStOP baseline Health Survey questionnaire was inconsistent.

#### **4.2.2 Data administration**

The Arthritis Research Primary Care Centre (ARPCC) at Keele University has data security systems and policies in place to meet legal requirements and obligations set by the Data Protection Act and follow General Medical Council (GMC), Caldecott Guardian and British Computer Society standards and guidelines

(Wilkie, 2005). Personally identifiable data that was held during mailing was stripped from research databases as soon as it was feasible. Individuals in the sample were allocated a unique study number. All questionnaires and survey data were anonymised by separating them from all contact details prior to any analysis (Wilkie, 2005).

Survey questionnaires were designed and processed using *Teleform*<sup>TM</sup> version 7.0 (Cardiff Software Inc, 1998). Once the questionnaires were scanned into computers, Optic Mark Recognition (OMR) technology was used to recognise marked choices for multiple choice questions, and Optic Character Recognition (OCR) and Intelligent Character Recognition (ICR) to recognise machine print and hand-writing. The automatic data entry system was organised using three linked software modules: Designer, Reader, and Verifier. Protocols were devised to facilitate data processing prior to scanning, and checks were conducted to ensure data accuracy. The system reviewed those items that could be interpreted erroneously (e.g. hand-writing) during the data process by the Reader and Verifier modules according to the validation criteria set during the design stage. These procedures were continuously monitored to ensure high levels of data accuracy (Wilkie, 2005).

The “Health Survey Questionnaire” included a body manikin, which the automatic data entry system could not process. Thus, body manikin related data was entered manually, using a transparent overlay to define pain areas (Figure 4.1) into a

Microsoft Access database. The manikin data entry was checked by another data clerk to identify errors, which were later corrected in the database (Muller, 2010).

The processed data was then exported to a SPSS<sup>®</sup> file and checked for anomalies in the full data set. In this process, where date of birth was missing, the signature on the questionnaire was checked against the name on the mailing list to make sure that the correct person filled the questionnaire. Then, the date of birth was copied from the mailing list to the questionnaire database. The same procedure was followed for the missing values of gender. If there were any unreasonable values of height and weight (e.g. heights indicated as greater than 8 feet or weight indicated as less than 3 stones) these were scored as missing. Occupations which could not be coded using the Office for National Statistics (ONS) classification of occupation (ONS, 2000; ONS, 2001) were checked against the original questionnaire to identify data entry errors (Wilkie, 2005).

#### **4.2.3 Ethical approval and informed consent**

Epidemiological research involves collecting an extensive amount of data on participants to describe physical, environmental and behavioural factors. Thus, researchers need to consider the ethical implications of their projects. The World Medical Association (WMA) has developed the Declaration of Helsinki as a statement of ethical principles for medical research involving human participants, including research on human material. They also detail data consent policies and more explicit guidelines for conducting research (Declaration of Helsinki, 1964). Following this, ethical guidelines have been formulated to protect the rights and

privacy of research participants, and to provide a standard for ethical dignity and professional conduct in research and practice. This ethical stance is located within a legal framework that includes the Human Rights Act and Data Protection Act. At the national level, the NHS National Patient Safety Agency (NPSA) includes the National Research Ethics Service (NRES) which developed a process to streamline ethical approval for NHS research projects. These guidelines demand the practice of obtaining informed consent from research participants. Researchers are expected to conduct all research in accordance with these guidelines to advance high standards in quality of research, and to cultivate public confidence, support and participation in these studies. Thus, securing ethical approval helps to maintain ethical standards of research practice, protect participants from harm, protect participants' rights, provide reassurance to the public, and protect researchers from unjustified criticism and harm.

Ethical approval for the NorStOP baseline study was gained from the North Staffordshire Local Research Ethics Committee (REC number: 1351) to conduct the study. Ethical approval for the 3 years follow-up was gained separately (REC number: 05/Q2604/20) and a substantial amendment was made to this application for the 6 years follow-up study.

At baseline, returned health survey questionnaires were taken as consent to be part of the baseline study. The Health Survey questionnaires were sent to participants with a study information sheet to explain the rationale for the study and at baseline, participants were asked to complete a consent form giving / not



giving permission for researchers to access their medical records and contact them again in the future for follow-up studies. At 3 and 6 years follow-up studies consent was again requested for permission for further contact.

#### **4.2.4 Follow-up of the NorStOP cohort**

Participant attrition is a concern in longitudinal studies as loss of participants may result in subsequent bias on findings (Ahern and Le Brocque, 2005). Thus, recruitment and follow-up methods in longitudinal studies should be designed to maximise the retention of these participants (Young et al., 2006).

##### *3 years follow-up*

The baseline responders who consented to further contact were eligible for the follow-up study at 3 years. Prior to the mailing, health informatics staff from the KGPRP checked the current practice registers to identify those who were no longer registered with the current practice. They also scrutinised the National Health Service Strategic Tracing Service (NHSSTS) to determine whether the person died or had changed their address. These checks were undertaken in two-weekly intervals throughout the mailing process. NHSSTS was not always able to identify the new addresses of those who had departed, thus these participants could no longer be followed-up. The same two-stage mailing process (Health Survey and the Regional Pains Survey) was repeated as per baseline sample, reminders for each stage were sent at 2 weeks (postcard only) and 4 weeks (a further questionnaire and a letter from the GP) respectively.

#### *6 years follow-up*

The same procedures for recruitment and mailing at baseline and 3 years were followed at the 6 years follow-up study, with one exception. Participants who were no longer registered with the practices were not traced using NHSSTS. This is because using NHSSTS was found to excessively lengthen the mailing process and not necessarily provide improved response rates (Muller, 2010).

#### **4.2.5 The Health and Regional Pains Survey Questionnaire Contents**

The Health Survey Questionnaire was designed by the NorStOP project team to measure a range of individual, health and contextual variables to allow the exploration of the consequences of health in populations. The survey instrument included previously well-validated questionnaires, a pain manikin and specifically developed single items. The Regional Pains Survey questionnaire included single items and validated questionnaires to explore hip, knee and foot pain, and hand pain or problems only.

One of the intentions of the ICF was to facilitate categorisation of health and contextual variables. The instruments included in the surveys and from which data was used in the analyses in this thesis were categorised in to ICF in the following section (Table 4.1). Additional variables included in the survey to investigate other research questions are not described in this thesis. A critical summary of the outcome measures used in this study is presented in Table 4.3 and 4.4 for ease of reference, and a synopsis of the threats to the validity and reliability of large scale self-report survey data is provided in Table 4.5.

**Table 4.1 Health & Regional Pains Survey Questionnaire contents**

ICF category	Measurement topic	Measurement method	Details	Scoring
Health conditions	■Self-reported health conditions	Single item	“Do you suffer from any of the following?  Chest problems Heart problems Diabetes	Yes/ No Yes/ No Yes/ No
Impairments	■Self-reported impairments	Single items	“Thinking back over the past 3 months have you suffered from any of the following?”  Falls Memory problems Cough with spit Breathless when walking Dizziness Weakness in arms/ legs Deafness Problems with eyesight Raised blood pressure	Yes/ No Yes/ No Yes/ No Yes/ No Yes/ No Yes/ No Yes/ No Yes/ No Yes/ No
	■Extent of bodily pain in the last 4 weeks	Single item	“In the past 4 weeks have you had pain that has lasted for one day or longer in any part of your body?”  “If so, please shade in the body chart, any pain that has lasted for one day or longer in the past 4 weeks	Yes/ No  0-44 areas
	■Selected joint pains in the last year	Single item	Have you had pain in and around your  Hands Hips Knees Feet (adapted from Jinks et al., 2001)	Yes/ No Yes/ No Yes/ No Yes/ No

■ Health Survey Questionnaire items

**Table 4.1b Health & Regional Pains Survey Questionnaire contents**

ICF category	Topic	Method	Details	Scoring
	Joint pain severity and stiffness for hand*	Scale	Australian/Canadian Osteoarthritis Hand Index (AUSCAN) (Bellamy et al., 2002)	None to moderate Severe to extreme
	Joint pain severity and stiffness for hip and knee*	Scale	Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) (Bellamy et al., 1988)	None to moderate Severe to extreme
	Joint pain severity for foot*	Scale	Foot Disability Index (FPDI) (Garrow et al., 2004)	None of the time/ On some days/ On most-every day(s)
	Joint pain chronicity*	Single item	Joint pain chronicity	< 3 months 3 months +
	■Anxiety	Scale	Hospital Anxiety and Depression Scale (HADS) (Zigmond & Snaith, 1983)	(0-7) (8-10) (11+)
	■Depression	Scale	Hospital Anxiety and Depression Scale (HADS) (Zigmond & Snaith, 1983)	(0-7) (8-10) (11+)
	■Cognition and alertness	Scale	Functional Limitations Profile – Cognitive and Alertness behaviour subscale (Bergner et al., 1981)	(0) (22.5-100) (0.1-22.4)

■ Health Survey Questionnaire items

\* Regional Pains Survey Questionnaire items (measured at baseline, 3YFUP and 6YFUP)

**Table 4.1c Health & Regional Pains Survey Questionnaire contents**

ICF category	Topic	Method	Details	Scoring
Body structures	■ Body Mass Index (BMI)	Single item	Height Weight	(<20) (20-24.9) (25-29.9) Obese (≥30)
Activity Limitation	■ Physical functioning	Scale	MOS SF-36 Physical Functioning subscale (Ware & Sherbourne, 1992)	(90.1-100) (70.1-90) (35.1-70) ≥35
Personal factors	■ Socio-demographic characteristics	Single item	Age	50-59; 60-69; 70-79; 80+ yrs
			Gender	Male; Female
		Single Item	Occupational class	Non-manual Manual
			Educational attainment “Have you gained qualifications through study as an adult?”	Yes/ No
	■ Social networks	Scale	Berkman and Syme Social Network Index (Berman & Syme, 1979)	High Med/ high Med/Low
Environmental factors	■ † Help and assistance	Single items	Have a confidant	Yes/ No
			Receipt of help and assistance from family/ friends and/or health/ social care professionals in: <ul style="list-style-type: none"> <li>• Washing and bathing</li> <li>• Looking after your skin, teeth, hair, nails</li> <li>• Putting on and taking off clothes</li> <li>• Using a toilet</li> <li>• Eating and drinking</li> </ul>	No need help Yes, receive help Yes, don't receive help Yes, receive help*
	■ † Aids and adaptations	Single items	Use of aids and adaptations for: <ul style="list-style-type: none"> <li>• Washing and bathing</li> <li>• Looking after your skin, teeth, hair, nails</li> <li>• Putting on and taking off clothes</li> <li>• Using a toilet</li> <li>• Eating and drinking</li> </ul>	Yes/No

■ Health Survey Questionnaire items

† This variables were measured at Health Survey Questionnaire at 6 years follow-up only

\* Receive help from family/ friends and/ or from health professionals/ social services

## Health Conditions

### *Self-reported health conditions*

The baseline Health Questionnaire included 12 single items to measure the experience of health problems and symptoms. Three out of twelve single items were defined as health conditions in accordance with the ICF framework. These were three of the most common chronic health conditions (apart from osteoarthritis) in the older UK population (chest problems, heart problems and diabetes). Each item had a simple yes/no response option. A count variable was created by adding up the occurrence of these conditions; this variable was categorised by combining adjacent numbers of health conditions into one category if their univariable estimates of association were similar (0, 1 and 2-3).

## Impairments

### *Self-reported impairments*

Again with reference to the ICF, the remaining nine items from the twelve single items in the Health Questionnaire were considered to measure impairments (falls, difficulty remembering, cough with spit, breathlessness when walking, dizziness, weakness in limbs, deafness, problems with eyesight, and raised blood pressure). Each item had a simple yes/no response option. A count variable was created by adding up the occurrence of these conditions. This variable was categorised, combining adjacent numbers of impairments into one category if their univariable estimates of association were similar (0, 1-3, 4-5 and 6-9).

### *Peripheral Joint Pain*

Joint pain is the most common symptom of osteoarthritis in older adults (Felson et al., 2000). For population studies of older adults, joint pain is the most appropriate way to define osteoarthritis (Peat et al., 2001). Single items were included in the questionnaire to identify the presence of peripheral joint pain (hand, hip, knee and foot). Participants were asked whether they had suffered from pain for one day or more in the last year for each joint site (hand, hip, knee and foot). Response options were yes/ no (Jinks et al., 2001).

### *Joint-specific characteristics*

Joint specific characteristics were measured in the Regional Pain Survey questionnaire at baseline, 3 and 6 years follow-up. Hand pain severity and stiffness were measured using Australian/Canadian Osteoarthritis Hand Index (AUSCAN) by 6 items (5 pains and 1 stiffness) that capture a combination of common symptoms in hand osteoarthritis (Bellamy et al., 2002). AUSCAN is a valid and reliable measure of hand problems in a community-dwelling population of older adults and was highlighted to have performed well both for its internal reliability and for its relationship with the external constructs (Dziedzic et al., 2007).

Lower limb pain and stiffness were measured using the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) for those with hip and knee pain (Bellamy et al., 1988). Scores for both hand and lower limb pain severity and stiffness ranged from 1 (none) to 5 (severe). Previously knee pain has been assessed in population surveys by a number of measurements such as Short

Form 36 (SF-36) (Ware and Sherbourne, 1992), Health Assessment Questionnaire (HAQ) (Fries et al., 1980), and Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC). WOMAC is a disease specific, self-administered instrument and was developed for use as an outcome measure in people with hip and knee osteoarthritis (Bellamy et al., 1988). Items are reported as subscales of pain, stiffness and physical function. WOMAC was found to be a reliable disease-specific measure for use in large epidemiological surveys, with good test-retest reliability for pain (ANOVA ICC: 0.88  $p>0.74$ ) in the general older adult population (Jinks et al., 2002).

Foot pain was measured using the Foot Disability Index (FPDI) (Garrow et al., 2004). This is a self-complete questionnaire that consists of 19 items to investigate function, pain intensity and appearance, each of which has three possible response categories: none of the time / on some days / on most or every day (s). Only the items that measured pain intensity were used in this analysis (10 and 14 to 17).

Chronicity of pain for hand, hip, knee and foot was measured by asking participants the number of days of hand/hip/knee/foot pain they have had in the last 12 months. The answer options were: less than 7 days / 1 to 4 weeks / >1 months but <3 months / 3 months or more. Joint pain chronicity was calculated by categorising those who reported 3 months or more hand / hip / knee / foot pain as having chronic pain as it is recommended by the criteria established by the




International Association for the Study of Pain (IASP) (Merskey and Bogduk, 1994).

### *Bodily pain*

Bodily pain was assessed to measure musculoskeletal comorbidity using a single item and a body manikin. Participants were asked “In the past 4 weeks have you had pain that has lasted for one day or longer in any part of your body?” and those responding positively were asked to shade their painful areas on a full body manikin. This manikin was separated into 44 mutually exclusive areas and a transparent template was used to mark the borders (Lewis et al., 2002) (Figure 4.1). Those responding negatively to the pain item and not shading the manikin were categorised as the ‘no pain’ group, and those responded positively to the pain item and shaded the manikin were considered to have pain in shaded areas. The bodily pain variable was derived from this variable to capture pain which was not linked to hands, hips, knees or feet pain. The areas included were mutually exclusive. The areas excluded were respectively knee pain (15, 19, 36, 40); foot pain (17, 21, 38, 42); hand pain (6, 10, 27, 31); hip pain (35, 39, 14a if 44 is +ve; 18a if 45 is +ve) (see Fig 4.1). Bodily pain was categorised by combining adjacent numbers of bodily pain into one category if their univariable estimates of association were similar (0; 1-4; 5-9; 10).

**Figure 4.1 The pain manikin**



### *General Health Characteristics*

The Medical Outcomes Study (MOS) Short Form (SF) 36 is a generic health status instrument (Ware et al., 1993). The SF-36 has been recommended for use in primary care research for its succinctness and the quality of its psychometric testing (Ware, 2000; Brazier et al., 1992). The SF-36 Health Survey was developed with an aim to capture physical and mental health, functioning and well-being under one succinct instrument (Ware and Sherbourne, 1992). The eight

domains of health represent the most frequently measured concepts in health surveys and those most affected by disease and treatment (Ware, 1995; Ware et al., 1993). These domains are Physical Functioning (PF); Role-Physical (RP); Bodily Pain (BP); General Health (GH); Vitality (VT); Social Functioning (SF); Role-Emotional (RE); and Mental Health (MH). The selected questionnaire items characterise various pointers of health, including behavioural function and dysfunction, distress and well-being, objective reports and subjective ratings, and both favourable and unfavourable self-evaluations of general health status (Ware et al., 1993). The questions include asking participants to rate their health, problems experienced as a result of their physical and mental health, and the amount of time their physical health or emotional problems interfered with their social activities (such as visiting friends and relatives). The usefulness of the SF-36 in postal surveys of older adults has received criticism due to items being frequently left blank on questionnaires by respondents (Mallinson, 1998). However, later studies which assessed the practicality and validity of using the SF-36 in community-dwelling older people recommended the use of SF-36 to obtain population scores in this population group (Walters et al., 2001; Nante et al., 1999).

The SF-12 Health Survey was developed as a shorter alternative to the SF-36 Health Survey for use in large-scale studies, particularly when overall physical and mental health are the outcomes of interest (Stewart & Ware, 1992). The SF-36 and SF-12 Health Surveys measure the same eight health domains, and each survey provides psychometrically-based physical component summary (PCS) and

mental component summary (MCS) scores. The PCS and MCS scores have a range of 0 to 100 and were designed to have a mean score of 50 and a standard deviation of 10 in a representative sample of the US population. This norm-based score allows comparison among the three surveys and across the studies published in the past 20 years. The SF-12 Health Survey uses just 12 questions to measure functional health and well-being from the patient's point of view. Previous studies recommended the SF-12 Health Survey to be a practical, reliable, and valid measure of physical and mental health, with less respondent burden (Jenkinson and Layte, 1997). In this study the SF-12 score for Physical Health (PCS) and Mental Health (MCS) was used to describe the health characteristics of the responders, and scores were expressed as means with standard deviations.

#### *Anxiety and Depression*

The Hospital Anxiety and Depression Scale (HADS) (Zigmond and Snaith 1983), was used to measure anxiety and depression. This was developed to identify the presence and severity of relatively mild mood disorders (anxiety and depression) in non-psychiatric hospital outpatients. It has been validated for use as a self-completed questionnaire (Herrmann, 1997; Lisspers et al., 1997). The HADS consists of 14 items (7 relating to the anxiety subscale and 7 relating to the depression subscale) with a four point response scale (from 0 representing absence of symptoms, to 3 representing maximum symptomatology). Scores can range from 0-21 on each subscale, higher scores indicating greater depression or anxiety, with cut-off points suggested between 0-7 (non-cases), 8-10 (borderline cases) and >11 (definite cases) (Pallant and Bailey, 2005). The scores were

calculated separately for anxiety and depression for the purpose of this thesis and categorised using the suggested cut-off points (Sackley et al., 2009; Mallen et al. 2006; Mallen et al., 2005).

### *Cognitive Complaint*

Cognitive complaint, which is the subjective reporting of perceived cognitive impairment, is common and more frequently encountered in older age and females (Westoby et al., 2009) was measured using the Sickness Impact Profile (SIP) Cognitive and Alertness subscale (Bergner et al., 1981) at all three time points (baseline, 3 and 6 years follow-ups). This is a ten item scale which was designed to measure sickness related changes in normal life (Bergner et al., 1981) and has been previously administered as a postal questionnaire (Trigg and Wood, 2003). The response options were a simple yes / no, and responses contribute to a summation of a weighted score, which is then converted to a percentage value, with higher scores indicating poorer function. Raw scores were categorised according to the sample distribution, using quartiles. 53% of the responders had a score of 0 (no cognitive impairment category) and the third quartile (75th percentile: 22.4) was divided into two categories as high (22.5-100) and low levels (0.1-22.4) of cognitive impairment. No cognitive impairment category was used as a reference category as higher scores indicates higher levels of cognitive impairment.

## Body Structures

### *Body Mass Index*

Studies suggest that weight problems are associated with osteoarthritis (Felson et al., 1997; Schouten et al., 1992) and obesity could be an important predictor of activity limitation and participation restriction in older people (McDonough and Jette, 2010). Therefore body mass index (BMI), an anthropometric measure to determine if people are at healthy weight, was used to calculate self-reported height and weight scores of the responders. Participants were asked to provide their height and weight using a preferable unit (either metric or imperial) at each time point to calculate BMI. This score was categorised into standard BMI groups which were used in previous studies of disability (WHO, 2004) as <20 (underweight), 20-24.99 (healthy body weight) 25-29.99 (overweight), >30 (obese). Healthy body weight (20-24.99) was used as a reference category.

## Activities

### *Activity limitation*

Activity limitations are difficulties that people may have in executing tasks (WHO, 2001). Activity limitation was assessed using the Physical Functioning subscale (PF-10) of the SF-36 Health Survey Questionnaire (Ware and Sherbourne, 1992), which was measured in NorStOP at all three time points. MOS SF-36 were used in previous population studies to measure activity limitation (Fransen et. al., 2002), and its use in primary care has been recommended (Ware, 2000; Jenkinson and Lyte, 1996). PF-10 consists of ten items to assess the extent of activity limitations

in a variety of physical activities. A criticism of the PF-10 is that the two items in this scale actually measure participation restriction (Wilkie 2005). Participants were asked “Does *your* health *now* limit you in these activities? If so, how much?” The activity items were; i) vigorous activities (such as running, lifting heavy objects, participating in strenuous sports); ii) moderate activities (such as moving a table, pushing a vacuum cleaner, bowling or playing golf); iii) lifting or carrying groceries; iv) climbing several flights of stairs; v) climbing one flight of stairs; vi) bending, kneeling or stooping; vii) walking more than a mile; viii) walking half a mile; ix) walking 100 yards; x) bathing and dressing yourself. The answer options were: yes limited a lot / yes limited a little / no, not limited at all. The scores for the ten items were summed and normalised so that the total score ranges from 0-100, with higher scores indicating better physical functioning. Following initial empirical work, raw scores were categorised using quartiles into lowest limitation (90.1-100), 3<sup>rd</sup> highest limitation (70.1-90.0), 2<sup>nd</sup> highest limitation (35.1-70.0), highest limitation ( $\geq 35.0$ ).

### Contextual Factors

Personal factors:

#### *Age and gender*

Participant’s age was derived from the date of birth provided in the questionnaire and categorised into 10 year age bands (50-59, 60-69, 70-79 and 80 and over) as recommended in previous studies (Thomas et al., 2004:b; Schoenborn, 2000).

### *Ethnicity*

Ethnic origin was measured using a single item in the baseline Health Survey questionnaire as 'White British / European', 'Afro Caribbean', 'Chinese', 'Asian', 'African', or 'Other'. However its distribution was not sufficiently diverse to include this factor in the analysis (i.e. 99.4% of the responders were white).

### *Socio-economic status*

Individual's socio-economic status was measured in three domains as occupational class, educational attainment and perceived adequacy of income:

#### i) Occupational class

Occupational class was assessed by asking participants "*If working what is your job title*" and "*If you are not working, or are retired, what was your last job title?*" and classified according to the Standard Occupational Classification (Office for National Statistics (ONS) (2000) (Table 4.2)). These categories were further revised to *non-manual* (higher managerial, higher professional, lower managerial / professional, intermediate occupations), *manual* (lower supervisory / technical, semi-routine occupations, routine occupations) and *self-employed* (Small employers and own account workers) occupations (Muller, 2010). The *self-employed* category was excluded from the analysis due to ambiguity of this classification as it was not possible to identify the nature of their job (i.e. manual or non-manual).



**Table 4.2      Categories of individual socio-economic classification**

1	Higher managerial and professional occupations a) Large employers and higher managerial roles b) Higher professional occupations
2	Lower managerial and professional occupations
3	Intermediate occupations
4	Small employers and own account workers
5	Lower supervisory, craft and related occupations
6	Semi routine occupations
7	Routine occupations
8	Never worked and long term unemployed

**Source:** Office for National Statistics, 2010

#### *Educational attainment*

Educational attainment was measured by a single item, asking participants *“have you gained qualifications through study as an adult?”* With answer options yes or no. Having adult qualifications (yes) was used as a reference category.

#### *Perceived adequacy of income*

Perceived adequacy of income was measured using a single item which asked *“Thinking about the cost of living as it affects you, which of these descriptions best describes your situation”*. The answer options were: ‘find it strain to get by from week to week / have to be careful with money / able to manage without much difficulty / quite comfortably off’ (Thomas, 1999). This measure was developed by Thomas (1999) and it measures participants’ financial circumstances without revealing the details of the responders’ income. Although measuring income adequacy has been previously criticised for the inherent variability of individuals

subjective evaluation (Ballantyne and Marshall, 2001), it has been validated by a study comprised persons aged 50 years and older in 12 countries from the Survey of Health, Ageing and Retirement in Europe as an objective economic indicator in predicting household financial distress (Litwin and Sapir, 2009).

#### *The Berkman and Syme Social Network Index*

The Berkman and Syme Social Network Index was originally constructed in 1979 to summarise the relationship between increasing social isolation and mortality (Berkman and Syme, 1979) and was used in this analysis to measure participants' social network status. The use of the Berkman and Syme Social Network Index is supported in population studies of adults to explore the role of social integration and support (Handley et al., 2012; Kearney et al., 2011; Shelton et al., 2011; White et al., 2009). The nine item version consists of asking the participants questions such as the number of children they have, number of children they see at least once a month, number of close relatives and friends, the number of religious meetings and/or community groups they attend to measure participants' social network status. Raw index scores were categorised as *“low / medium / medium high / high”* with high scores indicating better social ties.

#### *Having a confidant*

Participants were asked *“Is there any one special person you know that you feel close to; someone you feel you can share confidences and feelings with?”* The answer options were *“Yes or No”* (Michael et al., 2001).

## Participation

### *Self-care*

The Keele Assessment of Participation (KAP) was developed to measure participation for NorStOP. It includes one item that measures self-care restriction and the following section reviews the potential for this item to be used to meet the objectives of this thesis.

**Table 4.3 Psychometric properties of the outcome measurement scales used in this study**

<b>Name of measure / scale</b>	<b>Reliability / Repeatability</b>	<b>Validity</b>	<b>Ability to detect change</b>	<b>Strengths</b>	<b>Cautions</b>
<b>Australian/ Canadian Osteoarthritis Hand Index (AUSCAN)</b> (Bellamy et al., 2002)	Inter-observer and Intra-observer reliability were found to be 'substantial' and 'moderate' respectively. Two sources of data were used; first, a two-staged cross-sectional postal survey of older adults population using self-complete questionnaires; second, through the completion of AUSCAN by participants in a pilot study of clinical assessment of hand problems [median agreement beyond chance (kappa) was 0.75 (range: -0.03, 0.93) and 0.57 (range: -0.02, 1.00)] (Dziedzic et al., 2007).	The index has been subject to validation studies which have supported the clinimetric qualities such as the internal consistency, construct validity, factor structure, responsiveness and clinical relevance (Bellamy et al., 2002).	It is suggested that this scale should successfully detect changes in both pain and function, even when pain and loss of function do not occur concurrently (Allen et al., 2006).	The patient self-completed AUSCAN Indices are reliable, valid and responsive and can be recommended as primary outcome measures for future population-based studies.	Self-completed AUSCAN represent only the perspective of the participant. Additionally, it appears to pool aspects of long-term outcome and disease activity. It may be more appropriate to separate measures of disease activity from the long-term outcome (Kloppenburger et al., 2007).

Name of measure / scale	Reliability / Repeatability	Validity	Ability to detect change	Strengths	Cautions
<b>Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)</b> (Bellamy et al., 1988)	Overall, test-retest reliability of the WOMAC pain subscale has been variable across studies but generally meets the minimum standard; test-retest reliability has been more consistent and stronger for the physical function subscale, but the stiffness subscale has shown low test-retest reliability (McConnell et al., 2001).	Developed with the use of expert opinion, reviews of existing instruments, and surveys of patients with hip and/or knee osteoarthritis. High correlations and overlapping items between the pain and physical function subscales may pose measurement problems (Pua et al, 2009; Stratford and Kennedy, 2004).	It is suggested that activity overlap on the pain and function subscales limits the physical function subscale's ability to detect change over time (Stratford and Kennedy, 2004).	It is accepted as one of the most widely used self-report measures of lower extremity symptoms and function, and had been studied over 30 years in various contexts and populations. It has shown sufficient psychometric properties to be applied in population studies (Jinks et al., 2002).	The reliability of the pain subscale has been variable across studies, albeit meeting the minimum requirements.  The reliability of the physical functioning scale reported to be more consistent and stronger than the pain subscale.
<b>Foot Pain and Disability Index (FPDI)</b> (Garrow et al., 2004)	The construct of FPDI structure (pain intensity, functional limitation and appearance) was verified through confirmatory factor analysis in a general population of adults. Internal consistency for the three constructs was good	The FPDI was found to be a valid measure for use in community-dwelling older adults (Roddy et al., 2009; Roddy and Muller, 2009). A good level of agreement was	It is suggested to have a good ability to detect differences in disability levels reported by community subjects with and without current foot pain (Garrow et al., 2000).	Research into foot pain and function in the general population using postal questionnaires had been hindered by the lack of instruments to measure foot-related disability. FPDI, as a validated measure has potential to fill	The appearance subscale consists of only two items, which makes the scoring problematic (Muller and Roddy, 2009).

Name of measure / scale	Reliability / Repeatability	Validity	Ability to detect change	Strengths	Cautions
	(Cronbach's alpha 0.74, 0.92 and 0.77, respectively). Test-retest repeatability for the individual constructs ranged from fair to substantial (Roddy et al., 2009).	reported when compared with similar items on the ambulation subscale of the Functional Limitation Profile questionnaire (Garrow et al., 2000).		this gap.	
<b>Hospital Anxiety and depression Scale (HADS)</b> (Zigmond & Snaith, 1983)	Satisfactory or good item total correlations observed for anxiety and depression subscales (Herrmann, 1997). HADS was found to perform well in assessing the symptom severity of anxiety disorders and depression in both primary care patients and in the general population (Bjelland et al., 2002).	The concurrent validity was reported to be good to very good (Bjelland et al., 2002).	Re-test reliability shows a high correlation up to two-weeks, which decreases with longer term intervals; showing HADS is stable enough to withstand situational influences (Herrmann, 1997).	The scale is well accepted by general population and hospital patients, with response rates ranging from 95-100% in well-motivated study participants (Herrmann, 1997).	It is important to emphasise that anxiety and depression identified by self-report methods are only valid for screening purposes; definitive diagnosis must rest on the process of clinical examination.
<b>MOS SF-36 Physical</b>	The reliability of the scale and the summary measure	Previously has been validated for use as a generic-	It is suggested that SF-36 is sensitive to changes in health in	The SF-36 has proven useful in surveys of general	Mallinson (2002) proposed in a qualitative

Name of measure / scale	Reliability / Repeatability	Validity	Ability to detect change	Strengths	Cautions
<b>Functioning Subscale</b> (Ware and Sherbourne, 1992)	has been estimated using both internal consistency and test-retest methods. Published reliability statistics have exceeded the minimum standard of 0.70 recommended for measures used in group comparisons in more than 25 studies (Tsai, Bayliss, & Ware, 1997).	health related quality of life measure in general populations (Garratt et al., 1993; Brazier et al., 1992), and in selected populations with knee osteoarthritis (Brazier et al., 1999).	general populations (Busija et al., 2008; Hemingway et al., 1997).	and specific populations, comparing the relative burden of diseases, and in differentiating the health benefits produced by a wide range of different treatments (Ware, 2000).	assessment of the questionnaire that there are problems in which the respondents interpret the questions and suggested further research should explore differences in interpretation between different social groups.  Schroder et al. (2011) reported that the structure of self-rated health in patients with severe Functional Somatic Syndromes (FSS) (e.g. fibromyalgia) differed from people in the general population.
<b>Functional Limitations Profile (FLP) Cognitive and Alertness</b>	Adapted from the Sickness Impact Profile (SIP), this scale is a widely used health status	Not tested.	Not tested.	Unlike the SF-36, the FLP uses weights expressing the severity of individual items, which have	SIP measures behavioural impacts of sickness in terms of dysfunction and does not assess

Name of measure / scale	Reliability / Repeatability	Validity	Ability to detect change	Strengths	Cautions
<b>behaviour subscale</b> (Bergner et al., 1981)	measure. Previous studies support the measurement properties of this measure in population studies (Bruin et al., 1992; Carter et al., 1976), however there is no evidence of psychometric testing to support these assumptions.			been derived from previous research (Fitzpatrick et al., 1998).	levels of positive functioning. FLP-Cognitive and alertness subscale needs to be further tested to confirm its psychometric validity and reliability.
<b>Berkman and Syme Social Network Index</b> (Berkman and Syme, 1979)	There is no evidence of formal psychometric testing to support the use of Berkman and Syme Social Network Index in general population studies.	Not tested.	Not tested.	The index measures the level of social integration, taking into account number as well as relative importance of different ties. Thus, intimate ties are given more weight than group affiliations (Eng et al., 2002).	It is a lengthy scale, thus, it is not always used as an index to obtain a total score. Instead, different items are included/excluded in studies. This causes difficulty when comparing results across studies.
<b>Keele Assessment of Participation (KAP)</b> (Wilkie et al., 2005)	Internal consistency was not examined. Mean observed agreement over a 4-week period for dichotomised responses was 90%.	Cognitive and semi-structured interviews examined face validity, responder burden and content validity and found	Responsiveness and ability to detect minimum clinically importance difference has not been tested.	The instrument measures participation comprehensively and can be applied to the general population as a generic measure of	It has not been tested sufficiently to evaluate its appropriateness for evaluating interventions. Further



Name of measure / scale	Reliability / Repeatability	Validity	Ability to detect change	Strengths	Cautions
	Kappa ranged from 0.20 to 0.71.	that the instrument comprehensively measured participation. Demonstrated high levels of agreement with Reintegration to Normal Living Index and Impact of Participation and Autonomy.		participation. Its brief and concise design minimises the responder burden.	psychometric testing required for its responsiveness.  The brevity of the instrument is at the expense of detail.

**Table 4.4 Strengths and weaknesses of the specifically developed single items in the survey**

<b>Self-reported health conditions and impairments</b>	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>Allows the measurement of the presence or absence of common health conditions and symptoms in older adults and is classified in-line with the International Classification of Functioning (ICF).</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>The response option (yes / no) for each item does not measure the severity of these symptoms.</li> <li>Participants are asked to think back over the past 3 months, this might introduce recall bias.</li> <li>Low agreement was found when self-reported morbidity was compared with computerised general practice consultation records 2-years prior to the survey (Barber et al., 2009)</li> </ul>
<b>Selected peripheral joint pain</b>	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>It is adapted from a previously validated measure of knee pain (Jinks et al., 2001).</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>Participants are asked to think back over the past year to remember if they had suffer from pain for each joint site; as participants may not recall the pain for specific joint site, this might introduce recall bias.</li> <li>Self-reported joint pain may be arising from causes other than Osteoarthritis (e.g. bursitis, tendonitis).</li> </ul>
<b>Joint Pain Chronicity</b>	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>The criteria for chronic pain (<math>\geq 3</math> months) is based on the recommendations of the International Association for the Study of Pain (IASP) (Merskey and Bogduk, 1994).</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>Participants are asked to remember the number of days of hand/hip/knee/foot pain they have had in the last 12 months. This might be difficult to recall, which may impact on the precision of these estimates (recall bias).</li> </ul>
<b>Self-reported height and weight</b>	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>This score is categorised into standard Body Mass Index (BMI) groups which were used in previous studies of disability (WHO, 2004), allowing comparisons with the wider literature.</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>It relies on the self-reported health and weight measurements of the responders, which may have been subject to recall / information bias.</li> <li>There is an on-going discussion in the literature whether BMI is a reliable measure of body fatness for people.</li> </ul>

Single item which asked patients to indicate their occupation or if retired their previous occupation; this was coded to provide data on Occupational Class	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>• Responses to this measure were classified according to the Standard Occupational Classification (Office for National Statistics (ONS) (2000).</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>• Participants were classified into categories based on their answer to the question "If working what is your job title" and "If you are not working, or are retired, what was your last job title?" As the study population consist of older people who may have been involved in variety of roles throughout their working life, there is a potential for misclassification of their occupational class.</li> </ul>
Perceived adequacy of income	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>• This measure was originally developed by Thomas in 1999 and it measures the responder's financial circumstances without revealing the details of their income.</li> <li>• It has been validated as an objective economic indicator in predicting household income by a multinational study (Litwin and Sapir, 2009).</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>• It is "subjective" and can be influenced by negative appraisal.</li> </ul>
Educational Attainment	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>• Identifies whether participants gained qualifications as an adult; providing another level of socio-economic measure to the survey data.</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>• This item does not specify the type of qualifications gained through study as an adult, thus it might be misinterpreted depending on what the participant considers these are, which may result in misclassification.</li> </ul>
Extent of bodily pain	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>• Use of pain drawings which were than scored for the presence or absence of pain in each of 45 body areas has been tested in a previous study by margolis et al (1986) and scorers achieved a high rate of inter-rater agreement with relatively little training.</li> <li>• This method has been shown to be repeatable (Lacey et al 2005).</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>• Those, who responded "Yes" to the pain item but did not shade any areas on the manikin, were excluded from analyses. Those who responded "No" to the pain item but shaded at least one area on the manikin were also excluded from analyses. This may have influenced the results.</li> </ul>

Receipt of help and assistance from family/ friends and/or health/ social care professionals+	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>• This item breaks down daily activities into individual tasks such as 'washing and bathing' and 'using toilet' to aid to recall the difficulties participants may experience, thus require help when carrying out specific tasks.</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>• Validity has not been tested.</li> <li>• Participants were asked to recall the details of how they conducted daily daily activities in the past 4-weeks and whether they have had any help from family and friends or a health / social care professional. This may have resulted in recall / information bias.</li> <li>• Participants may have had a tendency to response this item positively to appear in a better light, (i.e. social desirability bias).</li> </ul>
Use of aids and adaptations+	
<p><u>Strengths:</u></p> <ul style="list-style-type: none"> <li>• This item lists commonly used aids and appliances when conducting the essential activities of daily living with an easy 'yes / no' option to aid the participants to recall the information.</li> </ul>	<p><u>Weaknesses:</u></p> <ul style="list-style-type: none"> <li>• Validity has not been tested.</li> <li>• Although a selection of aids and adaptations are provided in the questionnaire, this list is not exhaustive. Thus, participants might use alternative tools/ methods to help with their daily activities that are not represented in this list. This could result in misclassification.</li> <li>• Some items such as 'aids to help you wash and dress' may be open to different interpretations as what these aids might be. Again, this may also cause misclassification.</li> <li>• Participants were asked to recall the details of how they conducted daily daily activities in the past 4-weeks and whether they have used any aids and appliances to help with these activities. This may have been difficult to remember, thus, resulted in recall / information bias.</li> </ul>

+This measures are detailed in Chapter 8

**Table 4.5      A summary of the potential threats to the validity and reliability of self-reported survey data**

<b>Acquiescence bias</b> (Watson, 1992)	<ul style="list-style-type: none"> <li>❖ Refers to the tendency to respond positively.</li> <li>❖ Negatively and positively worded items should be evenly distributed to reduce this bias.</li> <li>❖ The process and interpretation of the data should be conducted with caution.</li> </ul>
<b>Extreme and moderate response bias</b> (Couch and Keniston, 1960)	<ul style="list-style-type: none"> <li>❖ Reflects the tendency for subjects to respond consistently using particular sections of the data.</li> <li>❖ Demographic factors may influence extreme responding.</li> <li>❖ Can be controlled by altering the response format.</li> </ul>
<b>Negative affectivity bias</b> (Watson et al., 1987)	<ul style="list-style-type: none"> <li>❖ Negative affectivity markedly inflates the associations between variables.</li> <li>❖ If identified, can be controlled by statistical methods.</li> </ul>
<b>Social desirability bias</b> (Paulhus, 1991)	<ul style="list-style-type: none"> <li>❖ Refers to the tendency to answer self-report items in such a way as to deliberately or unconsciously represent oneself in a favourable light.</li> </ul>
<b>Sampling bias</b> (Lijmer et al., 1999)	<ul style="list-style-type: none"> <li>❖ It is a consistent error that arises due to the sample selection.</li> <li>❖ It means the data collected may not be accurate, or represent the target sample.</li> </ul>
<b>Response bias</b> (Porta, 2008)	<ul style="list-style-type: none"> <li>❖ Systematic error due to differences in characteristics between those who choose or volunteer to take part in a study and those who do not.</li> </ul>
<b>Information bias</b> (Sanderson et al., 2007)	<ul style="list-style-type: none"> <li>❖ Bias in an estimate arising from measurement errors.</li> <li>❖ Also referred to as observational bias and misclassification.</li> <li>❖ The occurrence of information bias may not be independent of the occurrence of selection bias.</li> </ul>
<b>Recall bias</b> (Rothman et al., 2008)	<ul style="list-style-type: none"> <li>❖ It occurs when the way a respondent answers a question is affected not just by the correct answer, but also by the respondent's memory.</li> </ul>
<b>Measurement bias</b> (Porta, 2008)	<ul style="list-style-type: none"> <li>❖ It occurs when surveys do not measure what they intended to measure.</li> <li>❖ This results from flaws in the instrument, question wording, question order, interviewer error, timing and question response options.</li> </ul>

#### **4.2.6 Keele Assessment of Participation Restriction (KAP)**

The Keele Assessment of Participation (KAP) was developed as a self-complete instrument to provide estimates of participation restriction in population surveys (Wilkie, 2005). This was the first instrument developed specifically to measure participation restriction in population surveys (Magasi and Post, 2010). The KAP was developed to measure participation in line with the conceptual model proposed in the ICF. In the ICF, participation is seen as the outcome of a complex relationship between an individual's health condition, personal factors, and the external factors of the environment in which the person lives (WHO, 2001). The KAP consists of 11 items to measure participation restriction in different aspects of life (Table 4.6), reflecting the domains proposed to capture participation (Wilkie et al., 2005). Each of the 11 items are an aggregated measure and so responders must consider a number of relevant tasks when responding to one item. Each item is qualified by an individual's judgement of their performance because the occurrence of participation restriction is best perceived by the individual who experiences it, rather than describing deviation from an established normative standard (Wilkie et al., 2005). The cognitive testing of the KAP demonstrated that when individuals answered these questions they considered personal and environmental factors relevant to them and how these contribute to either preventing or facilitating participation. This was in line with the intended conceptual model and provided support for face validity of participation restriction (Wilkie, 2005).

One of the KAP items provides an aggregated measure of self-care (i.e. puts the self-care tasks such as washing, toileting, dressing, feeding, maintaining health under one umbrella). It is phrased to capture the individual's perception of their self-care needs being met. 'During the past 4 weeks, my self-care needs (examples are washing, toileting, dressing, feeding, maintaining health) have been met, **as and when I have wanted**' (response options: *all the time / most of the time / some of the time / a little of the time / none of the time*) (Table 4.3). The conceptual model means that individuals will consider the interaction between the individual and their environment when responding.

**Table 4.6 Keele Assessment of Participation**

1	During the past 4 weeks, I have moved around in my home, <b>as and when I have wanted*</b>
2	During the past 4 weeks, I have moved around outside my home, <b>as and when I have wanted*</b>
3	During the past 4 weeks, my <b>self-care</b> needs (examples are washing bathing, toileting, dressing, feeding, maintaining health) have been met, <b>as and when I have wanted*</b>
4	During the past 4 weeks, my home has been looked after, <b>as and when I have wanted*</b>
5	During the past 4 weeks, my things (belongings) has been looked after, <b>as and when I have wanted*</b>
6	Do you have relatives or any other people who depend on you? (Yes/ No) † If yes, during the past 4 weeks, were those people looked after <b>as and when you have wanted?*</b>
7	During the past 4 weeks, I have met and spoken to other people, <b>as and when I have wanted*</b>
8	During the past 4 weeks, I or someone else on my behalf, have managed my money, <b>as and when I have wanted*</b>
9	Do you choose to take part in paid or voluntary work? (Yes/No) † If yes, during the past 4 weeks, have you taken part in paid or voluntary work <b>as and when you wanted*</b>
10	Do you choose to take part in education or training courses? (Yes/No) † If yes, during the past 4 weeks, have you taken part in education or training courses <b>as and when you have wanted?*</b>
11	Do you choose to take part in social activities? (Yes/No) † If yes, during the past 4 weeks, have you taken part in social activities <b>as and when you have wanted?*</b>

\*Response options: All the time/ most of the time/ some of the time/ a little of the time/ none of the time. † Filter questions were added following pre-testing stage (Wilkie et al., 2005)

## Psychometric testing and development of the instrument

### *Pre-pilot studies: face validity, responder burden and content validity*

The review criteria suggested by the Scientific Advisory Committee of the Medical Outcomes Trust (Lohr et al., 1996) were used to inform and guide the development and testing of the KAP (Wilkie et al., 2005). The psychometric properties of the instrument were tested in two interview studies (cognitive interviews (n=11); qualitative interviews (n=4)) and a pilot study. Cognitive interviews examined face validity, responder burden and content validity (Wilkie, 2005). For this, purposive a convenience sample of adults aged 50 years and over was selected as a representative of older people with participation restrictions across a range of life areas (n=11). Participants completed the draft version of the KAP measure and were timed and observed for any difficulties they encountered during completion of the instrument. During this process, semi-structured interviews and open-ended probing were used to explore deeper meaning and interpretation of questions (Wilkie, 2005). Analysis of face validity reflected on the way items were completed with reference to the conceptual model of participation restriction. Responder burden was assessed by observing the difficulties and problems faced by participants when completing the KAP. Content validity was tested using the participant's opinions on the comprehensiveness of the KAP, considering whether it captured their own involvement in life situations (Wilkie, 2005). Qualitative interviews further tested the face and content validity of KAP in a different sample of participants (n=4) purposively sampled from a rheumatology ward to represent those with range of health conditions and restrictions. Semi-structured in-depth individual interviews were undertaken by an experienced



qualitative researcher to explore participants' accounts of the impact of joint pain, and the comprehensiveness of the KAP was examined through exploring the impact of joint pain on individuals in relation to the domains captured by the KAP instrument. For this, the interviewer was provided with the WHO definition of the domains covered by the questionnaire, from which the questions for the interview were generated (Wilkie, 2005).

### *Results of the pre-pilot studies*

The cognitive interview participants were divided into two categories. A healthy group (3 participants who had no complaints of joint pain or any other health problems) and an unhealthy group (8 patients receiving treatment on rheumatology ward in a local district general hospital). With regards to face validity, all items were understood and answered as intended, and all questions were regarded as appropriate and relevant by participants. Considering responder burden, there were no observed difficulties in completing the questionnaires, and for the content validity no additional items were highlighted by participants (Wilkie, 2005; Wilkie et al., 2005).

The qualitative interview participants also found the items relevant, felt able to communicate the problems they were having, and reported that the instrument was presented in an acceptable format. All restrictions mentioned in the interviews could be mapped to a domain of the KAP (Wilkie, 2005; Wilkie et al., 2005).

### Pilot questionnaire

The KAP was included in the survey instrument mailed to a random sample of 1461 adults aged 50 years and over, drawn from one GP register within the North Staffordshire General Practice Consortium. The inclusion of the KAP in this population based cross-sectional study allowed the meaning of responses to the filter questions, completion rates, and missing data to be examined as well as the distribution of KAP items in terms of restrictions to be observed (Wilkie, 2005).

Of 1461 questionnaires mailed, 1117 responded with completed questionnaires (adjusted response rate 71.7%). The completion rates for all KAP items were reasonably high; the completion of the KAP self-care item was 98.9% (n=1106). Of those, 53% of responders had no participation restriction and 5% reported restriction in self-care (Wilkie, 2005; Wilkie et al., 2005).

### *Face validity and conceptual discrimination*

The measurement model for filter questions was based on the assumption that responders who answered 'no' to the filter questions did not or could not have participation restriction in that domain (e.g. if they answered 'no' to the filter question 'have any dependents' they would not have participation restriction in 'looking after others'). Thus, to examine the correctness of this assumption, those 'no' responses indicating no participation were matched with corresponding frequency items. To support the assumption, participants who answered 'no' to the filter questions would be expected to indicate that they do not take part at all in the matched task (Wilkie, 2005). To establish conceptual discrimination, KAP items

were matched to other survey items from the same domain (e.g. self-care with self-care) that were intended to capture participation restriction measured from the perspective of the society (as opposed to person-perceived approach taken in KAP).

The proportion of responders who indicated 'no' in a filter question and also indicated that they did not participate in the corresponding task ranged from 64% for those who answered item 6 to 98% for those who answered item 10; this proportion was 98.9% for the self-care item (Table 4.6). There was no distinct association between person-perceived participation restriction and the frequency with which people engaged in activities in these domains. The observed agreement between the KAP self-care item and the paired question about the frequency (take a bath – most days or less) was 40%, indicating conceptual distinctions between person-perceived participation restriction and society-perceived participation restriction (Wilkie, 2005).

#### *Repeatability of the KAP*

Responders who gave permission for further contact were randomly designated to two groups and sent further questionnaires to address (i) the repeatability of the KAP, and (ii) validity of the KAP compared to items measuring participation derived from other instruments. Responses to the first and second wave of the KAP were then compared for both the main and filter questions. The repeatability of the KAP was examined by calculating the proportion of participants with agreement and agreement beyond chance (kappa K) for dichotomous variables.

Repeatability was assessed for the original five response options for each item, and the categories of 'none' and 'any' participation restriction (Wilkie, 2005).

Of possible 314 who responded at baseline, 196 returned a repeat questionnaire four weeks later (adjusted response rate 62.4%). For the four filter questions the mean observed agreement was 87.5%, and the chance corrected agreement ranged from moderate to substantial. The mean observed agreement for the five response options to each participation item was 75.1% (range: 68%-83%), and the chance corrected agreement was highest for the item which refers to mobility within the home ( $K_w = 0.64$ ; with 95% CI 0.54-0.74) and the lowest for the item referring to managing money ( $K_w = 0.34$ ; CI 0.09-0.56). Better repeatability was seen for the dichotomised response (mean agreement: 90.4%; range: 85.3-94.4%), with chance corrected agreement ranging from slight to substantial. The mean observed agreement for the five response options to KAP self-care item was 82.1%, and the chance corrected agreement ranged from slight to moderate ( $K_w = 0.45$ ; CI 0.33-0.55). Better repeatability was seen for the dichotomised response (mean agreement: 93.9%; range: 85.3-94.4%), with chance corrected agreement ranging from slight to moderate ( $K_w = 0.37$ ; CI 0.23-0.51) (Wilkie et al., 2005).

### Construct validity sub-studies

#### *Convergent validity and discriminant validity*

To examine construct validity of the KAP, a similar construct to that of the KAP was needed. During the mapping process two instruments were identified as containing a reasonable proportion of items that measured participation restriction

which could be deemed to capture similar construct to the KAP (Impact on Participation and Autonomy (IPA) (Cardol et al., 2001) and the Reintegration to Normal Living (RNL) (Wood-Dauphinee et al., 1988)). These instruments were used to examine the convergent and discriminative validity. The original response options were replaced with a uniform scale to allow responders to indicate how often they performed the participation task, and response scale for each item was dichotomised to indicate 'participation' and 'participation restriction' (Wilkie, 2005).

There were 102 responders who completed both KAP and RNL, and 104 responders completed both KAP and IPA. The mean percentage agreement for the 10 pairs of corresponding items (convergent validity) between KAP and RNL was 79.3% (range: 72-84%), and the mean percentage agreement for the KAP and IPA (23 items) was 76% (range: 74-97%). The highest observed agreement between items of the KAP and the RNL was for self-care, and lowest agreement was for looking after dependants.

Mean agreement between the 60 pairs of non-corresponding items (discriminant validity) between the KAP and RNL was 76% (range: 57-89%) and the mean agreement for the 160 pairs of non-corresponding items between the KAP and IPA was 82.8% (range: 66-97%), suggesting that individual's perception does not simply reflect the frequency of participation restriction. The high levels of agreement within some of the matched pairs also suggest that participation frequency may be more influential on the perception of participation restriction for some tasks than others.

### *Interpretation and summary*

The findings of the pilot studies support the validity and practicality of applying the KAP in a population survey to estimate participation restriction. These studies have shown that KAP items are comprehensible, can be completed in line with the conceptual model, and relevant and understandable by potential responders. The face validity test suggests that KAP captures person-perceived participation and the construct says that this differs from frequency. KAP items also have presented good convergent validity when compared with other constructs of participation restriction (RNL and IPA), with participation restriction correlating with items selected both for their correspondence and their difference to the items measured by KAP.

However, it was difficult to interpret the levels of chance-corrected agreement for the KAP self-care item and the categories of amount of participation restriction due to the effects of low prevalence. This is because Kappa statistics rely on a comparison between the observed and the expected agreement. When the prevalence of participation restriction is low, the possible agreement above chance can only be small due to high expected agreement; which then makes it difficult to achieve even moderate kappa values (Sim and Wright, 2005). Also the changes that occurred in the prevalence of participation restriction in the repeat mailing were not investigated. Thus, it is unknown whether these changes were due to a random bias or a systematic error as the levels of repeatability for other questionnaires included in the survey did not suggest any change in status of the concepts they were measuring. The change in prevalence could be due to several

factors such as the fluctuating nature of participation restriction, or the variance in the responder's appraisal of an item as person-perceived measures may be influenced by contextual factors (Wilkie, 2005). Although the issues with the repeatability of KAP can introduce a level of uncertainty to the longitudinal analysis in this thesis, it would be reasonable to accept that any participation restriction measured from a person-perspective rather than the societal norm will have variable estimations in between measurements due to change in personal and environmental factors over time.

The KAP's performance in validity and reliability tests indicated that it can be used as an aggregated measure to identify and provide estimates of participation restriction in postal surveys of population of adults aged 50 years and over. Thus, as a validated measure of participation, the KAP offers a good opportunity to measure self-care as participation from an individual's perspective, along with other components of NorStOP which offer the potential to look at health, joint pain and contextual factors and their links with self-care in line with the ICF framework.

## **4.3 Response to NorStOP**


### **4.3.1 Baseline survey response**

11309 people aged 50 years and older from the three general practices were identified as eligible for the NorStOP study by the KGPRP staff, and 11230 people were mailed after 79 people were excluded prior to mailing (22 had died, 33 had left the practices, 6 were unwilling to take part in the study and 18 had a severe psychiatric disorder, illness or learning difficulties) the Health Survey Questionnaire.

Following the mailing of the first wave of the questionnaires, 175 further exclusions were made due to: subsequent deaths and departures from the practice (n=45), returned questionnaires with addressee unknown (n=105), and those with comprehension / memory problems (n=25). During the mailing 255 people refused to take part in the survey, 109 had ill health and 2813 did not respond. Thus, 7878 responded to the Health Survey Questionnaire from the possible 11055 people, giving the adjusted response of 71.3% (Thomas et al., 2004:b) (Figure 4.2).



**Figure 4.2 Flowchart of response to NorStOP study at baseline**



\* **GP screen** - 6 were identified by their GP as unwilling to take part and 18 were stated to have a severe psychiatric disorder, illness or severe learning disabilities, hence unable to complete the questionnaire.

\*\* **Adjusted baseline response** - Adjusted response refers to the response rate calculated with those who were excluded from the mailing process, for example through death, stroke, dementia or incorrect address information, removed from the denominator.

### **4.3.2 Response at 3 years and 6 years follow-ups**


#### *3 years follow-up*

In the baseline study, 5366 people gave permission for further contact; of these 308 were excluded prior to mailing due to deaths and GP screenings and 5058 were mailed 3 years Health Survey Questionnaire. Of these, 57 were excluded during mailing due to death or departure, 31 dropped out due to being unable to complete the questionnaire due to ill health, 78 refused to return a questionnaire, 658 did not respond and 4234 returned Health Survey questionnaires. The adjusted response rate for 3 years follow-up was 84.7% (Thomas et al., 2007) (Figure 4.3).

#### *6 years follow-up*

Of the 4234 responders at three years, 3596 responders gave permission for further contact. Of this group 186 people were excluded prior to mailing at 6 years (181 died, 5 were screened as non-eligible by GPs) and 3410 people were mailed the Health Survey questionnaire. Following mailing, 37 people were excluded (change of address n=6; ill health n=5; wrong person n=4; deaths and departures n= 22), leaving 3373 eligible participants. Of those 542 did not respond (refusals n=73; ill health n= 40; non-response n=429). Completed questionnaires were received from 2831 people (adjusted response rate 84%) (Figure 4.3).

**Figure 4.3 Study consort from baseline to 6 years follow-up**



### 4.3.3 Completeness of the data

Completeness is an important attribute of data quality (Nagurney et al., 2005). The ultimate objective of data quality assessment is to fully understand the characteristics of the data set and determine strategies for the data analyses. Table 4.7 shows the levels of missing data associated with the key variables from the NorStOP baseline, 3 and 6 years follow-up studies used in this thesis. The levels of missing data on the KAP self-care item was very low, less than 2% at each time point. There were higher levels of missing data in certain constructs such as the SF-12 Physical and Mental Health Component, SF-36:PF10 Physical Functioning scale and the SIP Cognitive Alertness Score, where multiple items were required in order to construct a score.

**Table 4.7 Levels of missing data among items and scales within the NorStOP at baseline, 3 years and 6 years follow-up studies**

<b>Item/ Scale</b>	<b>Baseline (n=7878)</b>	<b>3 years (n=4234)</b>	<b>6 years (n=2831)</b>
	<b>%</b>	<b>%</b>	<b>%</b>
KAP Self-care item	1.9	1.9	1.2
SF36-PF10 Physical functioning scale	6.8	4.5	1.6
Hospital Depression Score (HAD)	2.4	1.8	1.4
SIP Cognitive Alertness Score	6.6	5.0	8.5
Perceived Adequacy of Income	2.8	1.6	1.8
Educational attainment	2.7	1.9	1.6
SF-12 Physical Health Component (PCS)	14.7	11.0	10.2
SF-12 Mental health Component (MCS)	14.7	11.0	10.2

## 4.4 The impact of non-response at baseline and attrition across the six years

### 4.4.1 Responders compared to non-responders at baseline

At baseline, a further 175 (1.5%) people were excluded following mailing and 3177 (28%) did not respond. There was a significant difference in the age and gender of responders, non-responders and those excluded (both  $p < 0.001$ ). Response was higher in the 60-69 and 70-79 age groups, and in women compared with men (71.8 % cf 68.2%). Exclusions were common amongst the oldest age group (80+ years) and non-response were higher in the youngest age groups (50-59 years). Proportionately there were more males in the non-responder (30.0% cf 26.9%) and exclusions (1.8% cf 1.3%) groups compared to women (Table 4.8).

**Table 4.8** Age and gender distribution of respondents, exclusions, and non-responders of NorStOP Health Survey at baseline

Overall n=11230	Exclusions n=175* (%)	Responders n=7878 (%)	Non-responders n=3177(%)	†P-value
<b>Age groups</b>				
50-59	53 (1.4)	2521 (65.6)	1273 (33.1)	p<0.001
60-69	28 (0.5)	2352 (74.4)	783 (24.8)	
70-79	40 (1.4)	2030 (75.6)	602 (22.4)	
80+	52 (3.4)	975 (63.1)	519 (33.6)	
<b>Gender</b>				
Females	81 (1.3)	4416 (71.8)	1651 (26.9)	p<0.001
Males	94 (1.8)	3462 (68.2)	1526 (30.0)	

\* Numbers do not add up due to missing data in exclusions

† Chi-squared test ( $\chi^2$ )

#### **4.4.2 The NorStOP compared to the local and national population**

Table 4.6 shows the comparison of the NorStOP responders to the population of England and Staffordshire by age, gender and ethnicity. This table shows that the age structure of the Staffordshire population is very similar to the population of England. Although the Staffordshire population had less than 10% of adults who were 80 years and older, compared to the English national population where adults 80+ years comprise 11.3% of the total population. However the NorStOP responders comprised a higher proportion of adults in the 80+ years category (12.2%) than the Staffordshire population, making the distribution of this age group closer to the English national sample. The proportion of those in the 60-79 years old groups was higher in the NorStOP responders compared to England and Staffordshire populations. Thus, indicating a possible response bias by age. However this difference is more likely to be related to the sampling frame than to the bias in the response to the survey.

Again, the gender structure of England and Staffordshire was very similar within each age group. There were slightly higher percentages of females in the NorStOP responders compared to the Staffordshire and England populations, but this difference was not markedly higher to suggest a response bias by gender.

The proportion of those who were white was markedly higher in the Staffordshire population and NorStOP responders compared to the English national population in each age group. This underlines that the NorStOP cohort is not representative of England as a whole in terms of ethnicity, however the effect of such bias is likely

to be insignificant due to the small proportion of non-white people overall aged 50 years and over in England (Table 4.9).

**Table 4.9 Comparison of the NorStOP responders to those of England and Staffordshire population by age, gender and ethnicity proportions**

	<b>50-59</b>	<b>60-69</b>	<b>70-79</b>	<b>80+</b>
<b>Age:</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
England <sup>a</sup>	38.4	28.3	22.0	11.3
Staffordshire <sup>a</sup>	40.6	28.7	21.0	9.7
NorStOP responders <sup>b</sup>	32.2	30.8	24.9	12.2
<b>Female:</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
England <sup>a</sup>	50.5	51.9	56.5	68.0
Staffordshire <sup>a</sup>	49.7	51.0	56.0	68.4
NorStOP responders <sup>c</sup>	53.9	52.7	57.1	69.5
<b>Ethnicity (white):</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
England <sup>a</sup>	95.3	95.1	97.3	98.5
Staffordshire <sup>a</sup>	98.7	98.7	99.5	99.4
NorStOP responders <sup>d</sup>	99.3	99.5	99.5	99.6

<sup>a</sup>Calculated from 2001 Census: standard tables downloaded from Nomis <https://www.nomisweb.co.uk/home/Census2001.asp>, accessed on 19 February 2009;

<sup>b</sup>Age group at baseline responders to NorStOP Health Survey Questionnaire; <sup>c</sup>Gender at baseline responders to NorStOP Health Survey Questionnaire stratified by age group;

<sup>d</sup>Ethnicity (white) at baseline responders to NorStOP Health Survey Questionnaire stratified by age group

## 4.5 Sample analysis

### 4.5.1 Descriptive characteristics of responders at baseline

#### Demographic and socio-economic characteristics

The survey responders at baseline had a mean age of 66.3 years (SD: 10.3); and the mean age was slightly higher for female responders compared to male responders (Mean age: 66.9 years (SD:10.6) cf 65.5 years (SD:9.8)). Just over half of the baseline sample were women (56%), and majority were from white ethnic backgrounds (99.4%). The slight increase in the proportion of women in this sample could be related to the evidence suggesting that women are more likely to respond to population surveys than men (Moore and Tarnai, 2002; Gannon et al., 1971). The ethnic homogeneity of this population could be explained by the fact that although black and minority ethnic groups make up over 16% of the population of England, this reduces to 8% in those aged 60 and over (ONS, 2009).

One in four responders at baseline lived alone, and 67% were married. Recent research also suggests a significant increase in the numbers of older people living alone in the recent years (Cambridge CHPR (Centre for Housing & Planning Research) (2010)), with women being more likely to fall into this category (Blood, 2010).

In terms of socio-economic status, the majority of responders had manual occupations (68.5%) and only 14% would describe their perceived adequacy of income as being 'comfortable'. 81.9% consider their income 'difficult to get by' with



or that they 'had to be careful', and 3.9% found it to be 'a strain'. The local economy of the North Staffordshire area has traditionally been dominated by coal mining, heavy manufacturing industries and the ceramics industry. The historically recent decline of these industries has had a significant impact upon the socio-economic dynamics of its population. Thus, the North Staffordshire area is known to be highly deprived in relation to health and employment compared to the rest of the country (Health and Social Profile of Staffordshire (Adults) 2010). However, the Department of Work and Pensions (Households Below Average Income) (2011) reports that 1.7 million pensioners (14%) live below the poverty line in the UK and 8% of pensioners are in persistent poverty (DWP, 2012). Previously the National Statistics for England reported that participants in lower income quintiles were more likely to report high prevalence of chronic diseases (National Statistics (The information centre) 2005).

In terms of level of education amongst responders, over 70% of participants report not having adult qualifications. Education is associated with material resources in Britain; people who are only educated up to secondary level are associated with a low level of material resources, whilst those who progress to further education (adult qualifications) come to be associated with attaining higher levels of resources (Burholt and Windle, 2006) (Table 4.10).

**Table 4.10 Descriptive characteristics of NorStOP Health Survey baseline responders (n=7878): demographic and socio-economic characteristics**

<b>Age Mean (SD)</b>		
	Overall	66.3 (10.3)
	Female	66.9 (10.6)
	Male	65.5 (9.8)
<b>Female n (%)</b>		4416 (56.1)
<b>Ethnicity (white) n (%)</b>		7633 (99.4)
<b>Marital Status † n (%)</b>		
	Married	5197 (66.8)
	Separated	55 (1.0)
	Divorced	440 (5.7)
	Widowed	1559 (20.0)
	Cohabiting	116 (1.5)
	Single	411 (5.3)
<b>Live alone n (%)</b>		1915 (24.3)
<b>Occupational class* n (%)</b>		
	Manual	4619 (68.5)
	Non-manual	2119 (31.4)
<b>Perceived adequacy of income † n (%)</b>		
	Comfortable	1083 (14.1)
	Little difficulty	3094 (40.4)
	Have to be careful	3179 (41.5)
	Strain	302 (3.9)
<b>Higher educational attainment † n (%)</b>		
	YES	2204 (28.8)
	NO	5460 (71.2)

† Totals do not add up to the 7878 due to the missing values

\* Totals do not add up to 7878 due to the removal of the self-employed category

### Health characteristics

The mean SF-12 PCS and MCS scores of responders were 40.7 (SD: 12.5) and 48.8 (SD: 11.2) respectively. Physical health and mental health are linked; individuals with mental health problems are at increased risk of a range of physical problems and conditions and poor physical health is also associated with the increased risk of developing mental health problems (Coalition Government's

Public Health White Paper Healthy Lives, Healthy People (2010). In terms of anxiety and depression, the majority of responders were not anxious or depressed (59.6% and 77.3% respectively). Of those with anxiety, 21.9% were borderline and 18.5% were deemed to be “definite” cases. Of those with depression, 14% were borderline and 9.7% were deemed as definite. This is similar to the findings of the Health Survey for England which studies chronic diseases and the health of older people in England. It reported that approximately 10-15% of the population aged 65 and over have depression, which is severe in 3-5% of older people (National Statistics (The information centre) 2005). Depression can also co-occur with cognitive impairment and pain (Sherbourne et al, 2009; Fishbain et al., 1997; Magni et al., 1993). Of those with cognitive impairment, 21.6% reported low levels and 24.9% had reported high levels. However, over half of responders had no cognitive impairment (53.5%). A previous population survey of community-dwelling older adults in the UK also reported high levels of cognitive impairment among community-dwelling adults. However the sample population was aged 75 and older in this study (Rait and Jones, 2005). In terms of joint pain, 43.8% reported hand pain, 51.5% reported knee pain, 33.1% reported hip pain, and 38.3% reported foot pain in the last 12 months. These estimates are comparable with previous studies of joint pain in community-dwelling older people. As mentioned in earlier chapters, the definition and measurement of joint pain is varied vastly across the studies. For example the prevalence of hand pain has been reported to be in the regions of 12% to 30% (Dziedzic et al., 2007; Urwin et al., 1998) and the prevalence of knee pain was reported to be in the regions of 19% (Urwin et al., 1998) and 47% (Jinks et al., 2004).

Over 40% of responders were overweight, 17% were obese, and less than 5% were underweight. It is reported that a quarter of the UK adult population is obese and nearly two thirds are beyond the threshold for being overweight in the UK. Obesity rates increase with age (up to the age of 70 years) and decline thereafter (Peeters et al., 2003). The most prevalent health problems experienced were breathlessness when walking (35.5%), followed by weakness in arm and leg (34%) and raised blood pressure (31%). A Health Survey conducted in New Zealand reported that increasing BMI and waist circumference were related to increased prevalence of cardiovascular disease, diabetes, high blood pressure and osteoarthritis (Turley et al., 2006). The Health Survey of England (2005) also reported that hypertension, arthritis, and having had a joint replaced were more common in overweight or obese adults, supporting the view that health problems and multiple conditions are interrelated (Table 4.11).

**Table 4.11 Health characteristics of baseline responders to NorStOP Health Questionnaire (n=7878)**

<b>SF-12 Physical Component Score</b>	
(0-100) Mean (SD)	40.7 (12.5)
<b>SF-12 Mental Component Score</b>	
(0-100) Mean (SD)	48.8 (11.2)
<b>HAD* Anxiety † n (%)</b>	
Non cases	4564 (59.6)
Borderline cases	1679 (21.9)
Definite cases	1414 (18.5)
<b>HAD* Depression † n (%)</b>	
Non cases	5932 (77.3)
Borderline cases	1077 (14.0)
Definite cases	669 (8.7)
<b>Cognitive Alertness †† n (%)</b>	
None	3936 (53.5)
Low level	1588 (21.6)
High level	1832 (24.9)
<b>Body Mass Index (BMI) n (%)</b>	
20 – 24.99	2744 (36.5)
25 – 29.99	3070 (40.9)
>30	1339 (17.8)
<20	1355 (4.7)
<b>Peripheral Joint Pain n (%)</b>	
Hand pain	3449 (43.8)
Knee pain	4060 (51.5)
Hip pain	2608 (33.1)
Foot pain	3020 (38.3)
<b>Selected health problems n (%)</b>	
Chest problems	1698 (21.6)
Heart problems	1410 (17.9)
Deafness	1467 (18.1)
Eyesight*	1688 (21.4)
Raised blood pressure	2445 (31.0)
Diabetes	643 (8.2)
Falls	1007 (12.8)
Difficulty remembering things	2148 (27.3)
Cough with spit	1460 (18.5)
Breathlessness when walking	2797 (35.5)
Dizziness/ unsteadiness	1931 (24.5)
Weakness in arm/leg	2677 (34.0)

\* Totals do not add up to the 7878 due to the missing values

† Hospital Anxiety and Depression Scale (Non cases: 0-7, Borderline cases: 8-10, Definite cases: 11+)

†† Functional Limitations Profile, Cognitive and Alertness behaviour subscale (None: 0, high levels: 22.5-100, low levels: 0.1-22.4)

\*Excluding need for spectacles

#### **4.5.2 Responders compared to lost at follow-up at 3 years**

Of 7878 responders at baseline, 3644 (46%) did not return a questionnaire at 3 year follow-up. Of those lost at follow-up, 2512 (69%) did not consent to further participation, 365 (10%) were excluded from the study and 767 (21%) did not respond to the 3 years follow-up questionnaire (Figure 4.3).

Compared to those lost to follow-up, responders at 3 years were younger ( $M = 64.3$   $SD = 9.3$ ), had better physical and mental health (Mean PCS: 42.0 (12.4); Mean MCS: 49.6 (11.1)), better physical functioning (Mean: 64.5 (30.4)), less depression and lower levels of cognitive impairment ( $p \leq 0.001$ ). There were no significant differences between responders and those who were lost to follow-up in terms of gender ( $p = 0.149$ ). However, there were significant differences in the socio-economic status of the responders and those lost to follow-up. Proportionately, fewer responders found their income to be of strain (3.5%) and, responders tended to possess adult qualifications (35.6%) ( $p \leq 0.001$ ) (Table 4.12). These differences indicate that the responders at 3 years were characteristically different from those lost to follow-up, which may affect the estimates of frequency of self-care restriction and the strength of association with other factors in the longitudinal estimates.

**Table 4.12 Responders versus loss to follow-up at 3 years**

<b>n= 7878</b>		<b>Total loss to follow-up n= 3644</b>			<b>Significance test</b>
	<b>Responders n=4234</b>	<b>Non-response n= 767</b>	<b>Non-consent n= 2512</b>	<b>Excluded n= 365</b>	
<b>Age</b>					
Mean (SD)	64.30 (9.3)	63.4 (10.2)	69.4 (10.6)	74.5 (9.9)	p≤ 0.001
<b>Gender n (%)</b>					
Female	2350 (55.5)	390 (50.8)	1518 (60.4)	158 (43.3)	p= 0.149
Male	1884 (44.5)	377 (49.2)	994 (39.6)	207 (56.7)	
<b>SF-12 PH</b>					
0-100: Mean (SD)	42.0 (12.4)	40.9 (12.5)	39.4 (12.4)	32.8 (11.4)	p≤ 0.001
<b>SF-12 MH</b>					
0-100: Mean (SD)	49.6 (11.1)	48.5 (11.0)	47.9 (11.2)	44.4 (11.7)	p≤ 0.001
<b>HAD depression score* † n (%)</b>					
Non cases	3411 (82.0)	573 (76.9)	1731 (71.6)	217 (61.0)	p≤ 0.001
Borderline cases	484 (11.6)	105 (14.1)	413 (17.2)	72 (20.2)	
Definite cases	263 (6.3)	67 (9.0)	272 (11.2)	67 (18.8)	
<b>Cognitive Alertness†† n (%)</b>					
None	2308 (57.4)	371 (52.1)	1133 (49.4)	124 (37.2)	p≤ 0.001
Low -level	859 (21.4)	163 (22.9)	495 (21.6)	71 (21.3)	
High-level	852 (21.2)	178 (25.0)	664 (29.0)	138 (41.4)	
<b>Activity limitation (PF-10)</b>					
0-100: Mean (SD)	64.5 (30.4)	63.8. (31.6)	55.5 (33.3)	38.0 (32.8)	p≤ 0.001
<b>Perceived adequacy of income n (%)</b>					
Comfortable	660 (15.8)	122 (16.3)	252 (10.5)	49 (13.9)	p≤ 0.001
Little difficulty	1729 (41.5)	283 (37.9)	939 (39.3)	143 (40.5)	
Be careful	1631 (39.1)	297 (39.8)	1107 (46.3)	144 (40.8)	
Strain	147 (3.5)	45 (6.0)	93 (3.9)	17 (4.8)	
<b>Higher educational attainment n (%)</b>					
YES	1481 (35.6)	238 (32.0)	395 (16.4)	90 (25.0)	p≤ 0.001
NO	2674 (64.4)	505 (68.0)	2011 (83.6)	270 (75.0)	

\* Numbers do not add up due to missing data

† Hospital Depression Score (Non cases: 0-7, Borderline cases: 8-10, Definite cases: 11+)

†† Functional Limitations Profile, Cognitive and Alertness behaviour subscale (None: 0, high levels: 22.5-100, low levels: 0.1-22.4)

#### **4.5.3 Responders compared to loss to follow-up at 6 years**

Of 4234 responders at 3 years, 2831 had responded (67%) to the six year follow-up questionnaire. 1403 (33%) were lost at follow-up. Of those who were lost at follow-up, 638 (45%) did not consent to take further part in the study at 3 years. A further 223 (16%) were excluded from the study and 542 (39%) did not respond to the six year follow-up questionnaire (Figure 4.3).

Again, the difference between the responders and those lost at follow-up at 6 years were statistically significant for demographic, health and socio-economic characteristics (Table 4.10). Responders at 6 years were younger (Mean age 65.9 (8.6)), predominantly female (65.4%) ( $p=0.043$ ), with better physical and mental health (Mean PCS: 43.2 (12.1); Mean MCS: 50.1 (10.9)), higher physical functioning (Mean: 61.9 (29.5)) ( $p\leq 0.001$ ), less depression ( $p\leq 0.001$ ) and lower levels of cognitive impairment ( $p\leq 0.001$ ). Responders had the higher proportion of people with higher educational achievement (40.4%) ( $p\leq 0.001$ ) and those whose income was comfortable (16.7%) compared to those who were lost to follow-up at 6 years ( $p=0.024$ ) (Table 4.13). Thus, the cohort at 6 years was a healthier cohort compared to those who were lost to follow-up. This cohort was used to study the empirical differences between two things in community-dwelling older people with joint pain. First, task-specific activity limitation and self-care restriction. Second, the link between environmental facilitators and self-care restriction (Chapter 8). A healthier cohort may affect the estimates of self-care restriction and task-specific activity limitation, and the strength of associations with other health, socio-



demographic and environmental factors at the cross-sectional study of older adults with joint pain at 6 years.

**Table 4.13 Responders versus loss to follow-up at 6 years**

<b>n= 4234</b>		<b>Total loss to follow-up n= 1403</b>			<b>Significance test</b>
	<b>Responders n=2831</b>	<b>Non-response n= 542</b>	<b>Non-consent n= 638</b>	<b>Excluded n= 223</b>	
<b>Age</b>					
Mean (SD)	65.9 (8.6)	67.1 (9.9)	70.9 (9.5)	74.5 (9.4)	p≤ 0.001
<b>Gender n (%)</b>					
Female	1598 (56.4)	291 (53.7)	374 (58.6)	87 (39.0)	p= 0.043
Male	1233 (43.6)	251 (46.3)	264 (41.4)	136 (61.0)	
<b>SF-12 PH</b>					
0-100: Mean (SD)	43.2 (12.1)	39.1 (12.2)	38.8 (11.9)	33.8 (10.9)	p≤ 0.001
<b>SF-12 MH</b>					
0-100: Mean (SD)	50.1 (10.9)	47.4 (11.4)	47.1 (11.2)	46.7 (10.8)	p≤ 0.001
<b>HAD depression score*†</b>					
Non cases	1737 (82.0)	312 (75.4)	320 (71.1)	108 (65.1)	p≤ 0.001
Borderline cases	235 (11.1)	55 (13.3)	81 (18.0)	31 (18.7)	
Definite cases	146 (6.9)	47 (11.4)	49 (10.9)	27 (16.3)	
<b>Cognitive alertness†† n (%)</b>					
None	1053 (53.8)	165 (44.0)	170 (44.9)	49 (35.0)	p≤ 0.001
Low -level	482 (24.6)	103 (27.5)	102 (26.4)	35 (25.0)	
High-level	424 (21.6)	107 (28.5)	115 (29.7)	56 (40.0)	
<b>Physical Functioning</b>					
0-100: Mean (SD)	61.9 (29.5)	51.2 (32.3)	48.6 (31.6)	36.6 (28.2)	p≤ 0.001
<b>Perceived adequacy of income n (%)</b>					
Comfortable	463 (16.7)	67 (12.7)	81 (13.3)	29 (14.0)	p= 0.024
Little difficulty	1168 (42.0)	212 (40.2)	264 (43.5)	90 (43.5)	
Be careful	1085 (39.0)	230 (43.6)	249 (41.0)	85 (41.1)	
Strain	63 (2.3)	19 (3.6)	13 (2.1)	3 (1.4)	
<b>Higher educational attainment n (%)</b>					
YES	1126 (40.4)	168 (31.5)	115 (18.5)	72 (33.6)	p≤ 0.001
NO	1659 (59.6)	365 (68.5)	508 (81.5)	142 (66.4)	

\* Numbers do not add up due to missing data

† Hospital Depression Score (Non cases: 0-7, Borderline cases: 8-10, Definite cases: 11+)

†† Functional Limitations Profile, Cognitive and Alertness behaviour subscale (None: 0, high levels: 22.5-100, low levels: 0.1-22.4)

## **4.6 Discussion**

### **4.6.1 Representativeness of the general population**

The use of general practice registers provided a representative sample of older adults in the general population who were not involved in other similar research at the time of the baseline survey. The sampling frame (n=11309) was similar in demography (age and gender) to the population of England, albeit there were differences in ethnic background.

A repeat mailing strategy was used in NorStOP to increase the response rate, as this method was previously successful in population studies (Nakash et al., 2006; Jinks et al., 2001). Acceptable response rates depend on many factors and an acceptable response rate for self-completed surveys is reported to be between 65-75% (Bowling, 2002; Sitzia and Wood 1998). The response to the NorStOP Baseline Health Questionnaire was 71%, which is within the suggested range. The sample analysis identified that non-response was higher in the youngest (50-59 years) age group, whereas exclusions were higher in the oldest age group (80+) compared to the mailed sample. However, there was no evidence of responders at the baseline Health Survey questionnaire been different than those in the England population in terms of age groups and gender. The proportion of white adults was slightly higher in responders at baseline compared to the English population, especially in the 50-79 age groups. However, considering that both the responders at baseline and the population of England's ethnicity was  $\geq 95\%$  white, this is unlikely to cause a biased sample in terms of the study sample's representativeness of the general population. Thus, the findings from the NorStOP

cross-sectional sample at baseline can be generalised to the national population with careful consideration.

#### **4.6.2 Study attrition**

Despite the two-stage mailing process being repeated at each follow-up and support of NHSSTS to boost response rates, there was a significant reduction in sample size from baseline to 3 and 6 years follow-up. In particular a reduction in the proportion of males aged 70 years and over from lower socio-economic backgrounds and with poorer health. This anticipated bias may affect results of the longitudinal analysis at 3 years by underestimating the onset and persistence of self-care restriction and weakening the associations with known risk factors. However there is potential to use statistical methods, for example imputation, to assess the effect of attrition over time.

#### **4.6.3 Data quality**

NorStOP was a large population survey, administered as a self-completed postal survey. Postal surveys are a simple, cost efficient way of reaching a widely dispersed population, many of whom would not normally attend a healthcare setting (Bates and Rogstad, 2000). Self-completion postal surveys eliminate interviewer bias (i.e. the researcher can't influence the respondents' answers by using verbal or facial expression to affirm or reject a particular answer). However, this also means that the researcher has no control over whether the intended person fills in the survey; anyone at the mailing address may have picked the form up and filled it in. However, problems completing the questionnaire due to specific

health problems such as reduced sight and physical functioning may have required help from a significant other or a friend to complete the questionnaire which may have introduced bias to the response by encouraging / discouraging certain answers.

To minimise human error in data entry, the questionnaire data were scanned and processed using the *Teleform*<sup>TM</sup> software package (Cardiff Software Inc., 1998) with the exception of the pain manikins, where the data were entered manually into a Microsoft Access database, using a transparent overlay to define pain areas. The data entry accuracy was audited through manually checking one in ten questionnaires for errors. Jinks et al (2003) reported that these processes enable accurate survey data transfer to be attained with *Teleform*<sup>TM</sup> in a study which evaluated a computer-assisted data entry procedure (including *Teleform*<sup>TM</sup>) for large-scale mailed surveys.

The missing data amongst the important variables such as the KAP self-care item, HADS scores, perceived adequacy of income and educational attainment were less than 3.0% across the 6 years follow-up. However, there were some instances of higher levels of missing data ( $\geq 5\%$ ). In particular where multiple items were required to calculate an overall score, such as SF-12 Physical and Mental Health scores. Thus, reducing the sample size available for the analysis. Issues with missing covariate data is explored in a later chapter.

#### **4.6.4 Measurement of self-care restriction using KAP**

Traditionally, self-care was measured as an individual's capacity to perform specific tasks of daily living (e.g. difficulty and dependence in washing and bathing, dressing and toileting (see Chapter 1 and 2)). In contrast, the KAP measures self-care as a biopsychosocial construct in line with the ICF model, where self-care is an outcome involving the interaction between the individual and their environment. It measures self-care from the individual's perspective and the individual judges whether their self-care needs are *"met as and when I wanted"*. There are no other published instruments to measure self-care as participation restriction, despite it being argued as a principal component of health-related quality of life (Wilkie et al., 2004; Wilkie et al., 2005). This is important as most health measurement instruments do not separate participation restriction from activity limitation (Wilkie et al., 2004; WHO, 2001). The philosophical concept and theoretical rationale behind the KAP, along with its simple format for application in epidemiological studies provides a good foundation to describe and investigate the extent of self-care restrictions in the community-dwelling older adults.

The test of construct validity demonstrated that there was some distinction between person-perceived participation restriction and the frequency with which people engaged in activities in these domains. The observed level of agreement between the self-care item and the paired question about frequency suggested that participants reported some specific functional limitations without indicating participation restriction in the corresponding item. This was because limitation was not considered important or severe enough to influence participation (Wilkie et al.,

2005). The response to NorStOP across 6 year follow-up also shows that less than 2% of data in KAP self-care item were missing, suggesting a high rate of data completeness.

Although KAP appears to be a good measure of self-care restriction in population surveys, it is not without limitations. The aggregated nature of the KAP self-care item means that the specific nature of the restriction (e.g. washing and bathing, dressing) cannot be identified. In developing the epidemiology of self-care restriction and associated factors in a population of older adults, this may not be a problem, but to develop clinical interventions for rehabilitation purposes, more information into the specific restrictions would be needed, such as to establish needs for specific dependencies in order to provide environmental facilitators (e.g. help and assistance in washing, aids and appliances for cooking).

Levels of kappa statistics need to be considered with percentage agreement and the low prevalence of certain items when judging the repeatability of the KAP. The assessment of repeatability suggests that person-perceived restriction may not be stable over time, but the other psychometric tests indicate that the KAP measures the construct at each time point. The prevalence of limitation in self-care tasks such as bathing is reported to be in the range of 29-40% in older adults (Gill et al., 2009; Gill et al., 2006:a; Naik et al., 2004). In contrast the prevalence of self-care restriction measured by the KAP in the pilot study was lower, which may indicate differences in the constructs (limitation in capacity at the level of the individual versus person-perceived participation including the use of environmental factors)

(Wilkie et al., 2005). Attrition bias also may have resulted in reduced estimates of participation restriction in self-care in the study sample compared to the “true” prevalence of these restrictions in the general population of older adults. However statistical methods can further assess the extent of this on observed estimates.

## **4.7 Conclusions**

Secondary analysis of the NorStOP dataset provides a practical, cost-effective, and suitable method of accessing a large population survey containing data that can be used to investigate self-care restrictions and joint pain in community-dwelling older people. The strength of using NorStOP is that it contains the KAP which measures participation in line with the ICF construct of participation and includes a number of variables to examine the associations with health conditions, activities, personal and environmental factors and joint pain. The limitation is the study attrition; this is an inevitable occurrence in studies where people are followed up over a long period of time. Thus, the issues with the selective loss to follow-up at three and six years will be considered in the following chapters for their potential to affect the analysis.

**Table 4.14      Summary of key findings**

- This chapter examined the suitability of the NorStOP dataset to meet the objectives of this thesis by reviewing the study design, the psychometric properties of the data collected, and sample derivation.
- The Health Survey instrument included previously validated questionnaires; a pain manikin and specifically developed single items. The Regional Pains Survey included single items and explored hip, knee and foot pain and hand pain and problems only.
- The KAP's performance in validity and reliability tests indicated that it can be used as an aggregated measure to identify and provide estimates of participation restriction in postal surveys of population of adults aged 50 years and over. Thus, as a validated measure of participation, the KAP offers a good opportunity to measure self-care as participation from an individual's perspective.
- Response to the NorStOP Health Survey Questionnaire was n=7878 at baseline (adj. response 71.3%); n=4234 at three years follow-up (adj. 87.7%); and n=2831 at six years follow-up (adj. 83.9%) studies.
- There was no evidence of significant difference between the age and gender distribution of the responders at baseline and the England population. Thus, the findings from this cross-sectional sample can be generalised to the national population with careful consideration.
- Despite the two-staged mailing process being repeated at each follow-up, and support of the NHSSTS to boost response rates, there was a significant study attrition from baseline to three and six years follow-up, particularly with a reduction in the proportion of males aged 70 years and over, from lower socio-economic backgrounds and with poorer health. Employing statistical methods such as imputation may help to minimise the effects of attrition in longitudinal analysis of this data.
- The missing data amongst the important variables such as the KAP item, HADS scores, perceived adequacy of income and educational attainment were less than 3% across the six years follow-up.
- The strength of using NorStOP is that it contains the KAP, which measures participation in line with the ICF, and includes a number of variables to examine the associations with health conditions, activities, personal and environmental factors and joint pain.



## Chapter 5

### **The prevalence and distribution of restricted self-care and associated factors in the general population at baseline**

#### **5.1 Introduction**

Cross-sectional studies are used to estimate prevalence and examine the relationship between the outcome of interest, and other explanatory factors in a defined population, at a specific time point (Porta, 2008). First of all it is important to quantify the size and nature of a problem within specific populations. Prevalence is used to quantify the size of a health problem in populations, and is defined as *the number of events of a given condition, in a given population at a chosen time* (Porta, 2008). When this term is used to refer to the total number of people with a specific condition at a specific point time, it is called the 'point prevalence'. When it is used to refer to observing the total number of people with a condition *at any time during a specified period*, it is called the 'period prevalence' (Porta, 2008). Prevalence estimates of conditions in the general population play an important role in public health policy making and service provision (Valanis, 1999). Understanding how prevalence may differ with demographic factors, such as age and gender, can help to generate hypotheses on causation. Identifying factors that are associated with self-care restriction can contribute further to the generation of hypotheses on causal mechanisms and provide some insight into potential prevention and intervention strategies. This is a key component in building the epidemiology of self-care, however, it is important to emphasise that the temporal

sequence of cause and effect cannot be determined in cross-sectional studies (Porta, 2008).

Self-care, as a form of participation in which the role of environmental factors is acknowledged, has not been examined before in population studies; hence there are no estimates of prevalence or empirical studies of the associations with self-care restriction. This study sets out to test this conceptual model of self-care (i.e. that self-care restriction will be associated with health and contextual factors). Age, female gender, physical inactivity / limitation, lower income, excessive weight and mental health problems are factors commonly associated with poor functional outcomes (Leroux et al., 2005; Woolf and Pfleger, 2003; Badley and Ibanez, 1994). Previous research into specific limitations in self-care activities (e.g. bathing) reported that the capacity to carry out this activity independently depends on more than one factor and can be multiply determined by both physiological and psychological factors (Mullick 2005; Naik et al., 2004). Emotional distress and physical limitation are key factors independently linked to limitation in self-care activities such as bathing and dressing (Verbrugge and Jette, 1994). The role of joint pain is explored specifically in this study. With an ageing population, and the expected rise in the number of people with musculoskeletal conditions, understanding the link with self-care is important.

This chapter describes the cross-sectional analysis of the North Staffordshire Osteoarthritis Project (NorStOP) baseline data. This includes the methods and results for studies that: (i) estimate the prevalence of person-perceived restricted

self-care in the general population of adults aged 50 years and over; (ii) describe the distribution of self-care restriction with age and gender; (iii) determine the associations between self-care restriction and selected health conditions, impairments, activity limitation and other contextual factors, with a specific focus on joint pain.

## **5.2 Aims and objectives**

Using cross-sectional analyses of the baseline data from the NorStOP in community-dwelling adults aged 50 years and over the aims of this work are to:

- i. Estimate the four-week period prevalence of person-perceived self-care restriction and its distribution by age and gender
- ii. Investigate the strength of cross-sectional associations between restricted self-care and selected health conditions, impairments, activity limitation and contextual factors with a specific focus on joint pain
- iii. Derive a parsimonious model of factors which are independently associated with restricted self-care
- iv. Examine factors that may explain the relationship between joint pain and self-care restriction

## 5.3 Methods

### 5.3.1 Dataset

The analyses described in this chapter uses data collected at baseline in the NorStOP study. Participants were community-dwelling adults aged 50 years and over in the general population who responded and completed the Keele Assessment of Participation (KAP) self-care item at baseline (n=7725).

### 5.3.2 Outcome


As outlined in chapter 4, the self-care item from the KAP was used to define participation and participation restriction. The item asks *“During the past 4 weeks, my self-care needs (examples are washing, toileting, dressing, feeding, maintaining health) have been met, as and when I have wanted”*. The answer options were *“All the time / Most of the time / Some of the time / A little of the time / None of the time”*. Those who answered *“All the time”* and *“Most of the time”* were classed as *restricted*, those who answered *“Some of the time”*, *“A little of the time”*, and *“None of the time”* were classed as *not restricted* in self-care; this provides a dichotomous outcome variable.

### 5.3.3 Covariates

The ICF framework was used to guide the cross-sectional analysis of the baseline data. Covariates included in the study were classified according to the ICF model of human function and disability (see Chapter 4). As the diagram shows, in the ICF disability and functioning are viewed as outcomes of interactions between health conditions (diseases, disorders and injuries) and contextual factors (Figure 5.1).

Contextual factors considered in this study were the *internal* personal factors such as age, gender and socio-economic characteristics. *External* environmental factors were not measured in NorStOP at baseline.

**Figure 5.1** ICF Model of Human Function and Disability (WHO, 2001)



**Source:** WHO, 2001

## **5.4 Statistical analysis**

The statistical analysis conducted in this study was designed and planned by the author of this thesis with the advice of the supervisory team. The analysis was performed by the author using SPSS Statistics Version 17.0 and Stata Version 11.0.

### **5.4.1 Prevalence of person-perceived self-care restriction and its distribution by age and gender**

Prevalence of restricted self-care was calculated by dividing the number of responders who reported restriction in self-care by the number of responders who completed the self-care (KAP) item and reported as a percentage.

Prevalence of restricted self-care was calculated overall and within age and gender groups to assess for potential confounding. Differences in prevalence of restricted self-care within age groups were determined using Pearson *Chi-squared* test for trend and by percentage difference with 95% confidence intervals for gender.

To explore the effect of gender on the distribution of restricted self-care with age, the prevalence of restricted self-care for each age group was stratified by gender.

### *Age standardisation using indirect method*

Age is one of the most common and important confounding factors in health studies as it is more often than not related to both the outcome of interest and the independent variable (Hennekens and Buring, 1987). Standardisation methods are used to enable valid comparisons between groups studied at different places and / or times (Wells, 2002). Age standardisation is a method that removes the confounding effect of age in order to make unbiased comparisons (Field, 2005). The two most common standardisation techniques used in the literature are the *direct* and *indirect* methods. Both methods are similar in principal, as they are based on taking a weighted average of category specific rates. However, the *direct method* is generally used for comparisons between two populations, and the *indirect method* is used when the age-specific rates for the study population are not known but the total number of events is known, or when calculating rates for small populations where fluctuations in age-specific rates can affect the reliability of rates calculated using the direct method (Armitage and Berry, 1994) (Table 5.1).

**Table 5.1      Age standardisation methods**

<b>Direct method</b>	<b>Indirect method</b>
$SR = (\text{SUM}(r_i * P_i)) / \text{SUM } P_i$	$SR = (C / \text{SUM}(R_i * p_i)) * R$

**SR** is the age-standardised rate for the population being studied

**ri** is the age-group specific rate for age group i in the population being studied

**Pi** is the population of age group i in the standard population

**C** is the observed number of events\* in the population being studied

**SUM(Ripi)** is the expected number of events in the population being studied

**Ri** is the age-group specific rate for age group i in the standard population

**pi** is the population for age group i in the population being studied

**R** is the crude rate in the standard population

\* 'Events' can include deaths, incident or prevalent cases of disease or other conditions, or health care utilisation occurrences.

**Source:** Australian Institute of Health and Welfare (AIHW) Knowledgebase, 2005

The age and gender structure of England differs to that of NorStOP (Chapter 4). Indirect standardisation was used to compare the actual number of self-care restriction with the number expected in the population of England and Wales, as the census data were not available only for England. The observed prevalence of restricted self-care in NorStOP was adjusted using the most recent available estimates of the national population for England and Wales at the time this thesis was undertaken (Census 2001). Calculation was applied to age groups stratified by gender respectively. Figures in each group were added together and divided by the total number to acquire the total (final) standardised prevalence estimate of restricted self-care in people aged 50 years and over in the general population.

The NorStOP population who responded to the KAP self-care item at each age band was shown in column (a). The number of people who reported restricted self-care in each age band was shown in column (b). The age-specific prevalence was calculated and shown in column (c).

Rate/ 1000 person-years

= (Observed number of cases/ number in NorStOP population) x 1000

= ((b)/ (a)) x 1000

To compare the prevalence of restricted self-care in the NorStOP sample to the population of England and Wales, whose age distributions may differ, the observed rate was adjusted to the reference population using UK Census (2001) data. Age group specific rates, which occurred in the study population, were applied to the appropriate age groups in the England and Wales population, and



the number of cases would have been expected if the study population rates were identical to national rates were calculated as below:

Expected number of cases

= Prevalence (per 1000 individuals) in the study population x (number of people in the national population/ 1000)

= (c) x (d)

This was calculated for each age-group stratified by gender and the results were shown in column (e).

The standardised rate

= (Total expected number of cases/ Total national population) x 1000

= (Total (e)/ Total (d)) x 1000

The combined standardised prevalence rate

= (total expected male cases + total expected female cases) x 1000

(total standard male population + total standard female population)

#### **5.4.2 Factors associated with restricted self-care in the general population**

Following prevalence estimates, factors associated with restricted self-care in the general population were investigated through a set of univariable and multivariable logistic regression analyses. These were conducted by the author using SPSS Version 17.0.

When carrying out a univariable analysis it is important to understand that the relationship between two variables could be confounded by another factor (Henneken and Buring, 1987). A confounder variable is associated with both the probable cause and the outcome (Jordan et al, 1998). In order to control the potential effects of confounding, multivariable techniques were used to assess independent associations with restricted self-care. Therefore each variable that was associated with restricted self-care in the univariable analysis was adjusted for age and gender, and then all other associated factors in the multivariable analysis, to identify their independent association with restricted self-care using a backward-stepwise logistic regression.

Logistic regression method was used to investigate the associations between self-care restriction and selected covariates, and adjust for potential confounders, because the dependent variable was binary / dichotomous and all independent variables were binary or categorical (Field, 2005; Jordan et al., 1998). Logistic regression is used for the prediction of the probability of occurrence of an event by fitting data to a logistic function (Agresti, 2007). Logistic regression is carried out through a logistic or logit transformation that is used to link the dependent variable to the set of other predictive or explanatory variables (Tranmer and Elliot, 2008). As both the dependent variable and explanatory variables were categorical, the logit modelling was used. The logit modelling is represented as:

$$y = \text{logit}(p) = \text{Log} [ p / (1 - p) ]$$

Predictive variables were also coded 1 for exposed and 0 for non-exposed. The model can be represented as below:

$$p_i = p(y=1 | X = x_i)$$

$$\text{Log}(p/1-p) = \text{logit}(p) = \beta_0 + \beta_1 x_i$$

In the above formula,  $p_i$  represents the probability of having self-care restriction and  $x_i$  is the predictive variable (e.g. gender). Therefore the parameter  $\beta_0$  gives the log odds of a study participant having self-care restriction in non-exposed (when  $x_i = 0$ ) and  $\beta_1$  shows how this odds differ in exposed (when  $x_i = 1$ ) (Tranmer and Elliot, 2008).

This logistic regression model was also used to assess the association of self-care restriction with a number of variables ( $x_1, x_2, x_3, \dots, x_i$ ), by calculating corresponding odd ratios from the model coefficients ( $\beta_1, \beta_2, \dots, \beta_i$ ) to estimate independent associations with restricted self-care in the multivariate analysis.

Odds ratios with 95% confidence intervals (CI) were used to determine the direction and strength of associations between self-care restriction and selected covariates. Odds ratios can be calculated using 2 x 2 tables as below:

		Exposure	
		Yes	No
Self-care restriction	Yes	a	b
	No	c	d
		$n^1$	$n^2$

The odds ratios were calculated as the ratio of self-care restriction within the exposed group, to the odds of self-care restriction within the non-exposed group (Field, 2005).

$$\begin{aligned}\text{Odds of self-care restriction in the exposed} &= (a / n^1) / (c / n^1) \\ &= a / c\end{aligned}$$

$$\begin{aligned}\text{Odds of self-care restriction in the non-exposed} &= (b / n^2) / (d / n^2) \\ &= b / d\end{aligned}$$

$$\begin{aligned}\text{Odds of ratio of the self-care restriction} \\ \text{among the exposed to the non-exposed} &= (a/c) / (b/d) \\ &= ad / bc\end{aligned}$$

A ratio of 1.0 represents that the odds of self-care restriction in both exposed and non-exposed groups are the same; hence there is no difference between the two groups. However when this ratio is over 1.0 it is indicative of a positive association with the independent variable, meaning that the odds of having self-care restriction is higher in the exposed than the non-exposed. The opposite is true when a ratio is less than 1.0, therefore the odds of having self-care restriction would be lower in the exposed than the non-exposed (i.e. the exposed are less likely to have self-care restriction) (Silman and Macfarlane, 2002). In order to assess the precision of these estimates, confidence intervals (CI) were calculated to provide information on the boundaries, within which we believe the true value of the population mean will fall.

Cross-tabulations of restricted self-care were performed with each variable. Associations were presented as odds ratios with 95% confidence intervals, first unadjusted and then adjusted for age and gender and all other variables that were associated with restricted self-care from self-reported health conditions and impairments, activity limitation and contextual factors.

**Table 5.2 Variables included in the multivariable analysis**

ICF component	Variable	Categories
Personal Factors	Age*	0-59, 60-69, 70-79, 80+
	Gender*	Male, Female
	BMI	Normal, underweight, overweight, obese
Health Condition	Number of health conditions	0,1,2-3
Impairment	Number of Impairments	0, 1-3, 4-5, 6-9
	Number of non-OA body pain areas	0, 1-4, 5-9, 10+
	Number of joint pains	0,1,2,3,4
	Anxiety	Non-case, borderline case, definite case
	Depression	Non-case, borderline case, definite case
	Cognitive impairment	None, low level, high level
Activity limitation	SF-36 physical functioning scale	Lowest level, 3 <sup>rd</sup> highest, 2 <sup>nd</sup> highest, highest
Contextual factors	Occupational Class	Non-manual Manual
	Higher educational attainment	YES NO
	Perceived adequacy of income	Strain Be careful Little difficulty Comfortable
	Have a confidante	YES NO
	Berkman and Syme Social Network Index	High Medium high Medium Low

\* Age and gender included as potential confounders

#### **5.4.3 Parsimonious model of key factors associated with restricted self-care**

Variables included in the multivariable analysis were factors that were significantly associated with restricted self-care in the univariable analysis after adjusting for age and gender ( $p < 0.05$ ). These factors were also checked for collinearity with other variables in the model, using 2 x 2 tables. It is suggested, that highly correlated variables could result in increased variance, hence a loss of significance in multivariable analysis (Field, 2005). Backwards-stepwise binary logistic regression was used to select / remove variables from the multivariable model. This approach was chosen as it is suggested that in statistical models each parameter introduced to the model adds some uncertainty to it (Harrell, 2001). Excessively complex models suffer from over fitting and have poor predictive power (Hosmer and Lemeshow, 1995).

The fixed statistical level rule was predetermined for the removal from the model as  $p=0.05$ , and for re-entry to the model as  $p=0.01$ . Recommended values in the literature for a decision rule range from 0.15 to 0.20 (Hosmer and Lemeshow, 1989). These somewhat narrow values were chosen for providing precision in these estimates and acquiring the most parsimonious model of factors. In order to achieve the maximum available sample size, the variables that remained associated with restricted self-care following adjustment for all other factors were included in the parsimonious model. This was achieved by running the backward-stepwise model and then, when it had found the final set of predictors, refitting the model as a parsimonious model of key factors associated with restricted self-care. Adjusted odds ratios with

95% confidence intervals were calculated to determine the strength of association for each variable.

#### *Validity of the parsimonious model*

Model assumptions play an important part in determining the quality of associations, therefore when a model satisfies its underlying assumptions, estimates would be more accurate (Field, 2009). It is also important to consider that the performance of the model could be overestimated on the sample of participants that were used to construct the analysis, resulting in an optimistic prediction of the outcome (Harrell, et al., 2001 and 1996; Houwelingen and Le Cessie, 1990).

Calibration and discrimination are often used in studies with binary outcomes to measure performance (Steyerberg et al., 2010). Calibration refers to the ability of a predictive model to match predicted and observed rates across the entire spread of the data. A predictive model, in which the number of observed cases supports the number of cases predicted by the model, indicates good calibration. Good calibration is crucial for attaining a realistic risk adjustment (Houwelingen and Le Cessie, 1990). A common measure of calibration used is the Pearson Goodness-of-fit test. The Pearson Goodness-of-fit test provides the sum of differences between the observed and the expected frequencies of outcome (counts of observations); each squared and divided by the expectation. The resulting value can be compared to the chi-squared distribution to determine the goodness of fit (Laub and Kuhl, 2005).

Models which can differentiate between people with and without the expected outcome are considered to have good discrimination (Cook, 2008). For binary outcomes, discrimination can be quantified by the c-index which is the area under the receiver operating characteristic (ROC) curve (Steyerberg et al., 2000). The c-index is based on the ranks in individuals with and without the outcome and ranges from 0.5 (no predictive ability) to 1 (perfect predictive ability), with higher values indicating better discrimination (Pepe, 2003). The c-index is obtained by plotting sensitivity against 1-specificity for different cut-off points for the predicted risk.

Internal validity of the parsimonious model was conducted using the original baseline sample employed in this study (NorStOP<sup>1</sup>). The external validity of the parsimonious model was tested in the NorStOP<sup>2</sup> for comparison of the results with an alternative sample population. The NorStOP<sup>2</sup> sample was another sub-cohort of NorStOP. Participants in NorStOP<sup>1</sup> were recruited during March and April 2002 and used as the main sample in this thesis. Participants in the NorStOP<sup>2</sup> study sample were recruited between July 2002 and June 2003 with an aim to provide a sampling-base for a clinical epidemiology study of knee pain and problems (CAS-K; Peat et al., 2004) (see Chapter 4).



Validation was carried out in following stages:

i) Apparent performance:

Apparent performance is defined as the estimate of a model performance from the full, original sample (Steyerberg, 2009). The apparent performance is deemed to be optimistic because both the development and the testing of the model is carried out on the same data. Multivariable logistic regression was used to perform estimates. The dependent variable was self-care restriction measured by the KAP self-care item. Coefficients were the variables from parsimonious model of key factors associated with restricted self-care. ORs with their respective 95% confidence intervals were used to summarise the contribution of the each predictor in the model. The fitting of the model to the data was assessed using Pearson Chi Squared statistic ( $X^2$ ). Large values of  $X^2$  and small p-values indicate a lack of fit of the model (Donati et al., 2004). The c-index (ROC curve) was used as a summary measure to evaluate how accurately the model discriminated between respondents with or without restricted self-care.

ii) Bootstrap performance:

The bootstrap method was introduced by Efron in 1979 as a resampling procedure for estimating the distribution of statistics based on independent observations. Bootstrapping is a computer intensive method for a resampling technique. It repeatedly generates samples from an underlying population by extracting samples with substitution from the original data, of the same size as the original dataset (Efron and Tibshirani, 1993). Each of these resamples will

randomly differ from the original sample. As the cases in these resamples vary, the statistic calculated from one of these resamples will also have different values (Cugnet, 1997).

The bootstrapping approach was conducted using Stata 11 as below;

**Step 1.** An empirical probability distribution,  $F_n$ , was constructed from the original sample by placing a probability of  $1/n$  at each point,  $x^1, x^2, \dots, x_n$  of the sample. This is the empirical distribution function of the sample, which is the nonparametric maximum likelihood estimate of the population distribution,  $F$ .

**Step 2.** Resample was drawn from the empirical distribution function,  $F_n$ , as a random sample of size  $n$  with replacement.

**Step 3.** The  $X^2$  and c-index statistics ( $T_n^*$ ) were calculated for the resample

**Step 4.** Steps 2 and 3 was repeated  $B$  (500) times

**Step 5.** The relative frequency histogram was constructed from the  $B$  number of  $T_n^*$ s by placing a probability of  $1/B$  at each point,  $T_n^{*1}, T_n^{*2}, \dots, T_n^{*B}$ . The distribution obtained was the bootstrapped estimate of the sampling distribution of  $T_n$ . This distribution was then used to make inferences about the model calibration and discriminative powers, which was estimated by  $T_n^*$ .

(Efron and Tibshirani, 1993)

iii) Split-sample performance:

In the split-sample (reduced sample, e.g. two-thirds of the sample) approach the data is split into two sets and the first part is used to develop the model and the second part is used to measure its performance. Thus, model performance is tested on similar yet independent datasets (Steyerberg et al., 2001). Split-sample performance was carried out in Stata 11 by randomly selecting 65% of the data. The performance of the parsimonious model was then tested on this sample using multivariable logistic regression to obtain odds ratios with 95% confidence intervals. HL  $X^2$  statistic and ROC curve (c-index) were also obtained.

iv) Expected optimism:

Expected optimism of the parsimonious model was calculated as the difference between the estimated bootstrap and the test performance (iv= (ii – iii)).

v) Optimism corrected performance:

The optimism corrected performance of the parsimonious model was calculated as the difference between the estimated apparent performance (i) and the optimism (iv) (v= i - (ii – iii)). It is understood that the bigger the absolute difference, the bigger the optimism (Steyerberg et al., 2003).

#### **5.4.4 Factors explaining the relationship between joint pain and self-care restriction**

The relationship between joint pain and self-care restriction may be explained in addition to confounders, by mediators or moderators (Wilkie et al., 2007: a-b; Machado et al., 2008). *“A variable that occurs in a causal pathway from an independent variable to an outcome variable and causing variation in the outcome variable and itself is caused to vary by the original independent variable is called an intermediate variable”*. This can also be known as a mediator (Porta, 2008, p.131). A factor that modifies the measure of effect of an assumed causal factor is referred as the moderator or modifying factor (Porta, 2008). The ICF model suggests that contextual factors can influence the relationship between participation restriction, health conditions, impairments and activity limitation (WHO, 2001). Thus, it may be plausible to assume that these factors may be mediating / moderating the relationship between joint pain and self-care restriction.

Multivariable logistic regression was used to test the hypothesis that each of the variables that were included in the parsimonious model influence the association between joint pain and restricted self-care following the methods proposed by Baron and Kenny (1986). This method suggests that if the relationship between variable x (e.g. pain) and outcome y (e.g. self-care restriction) is significantly reduced when adjusted for variable z (e.g. depression), variable z can be considered as a potential mediator (i.e. it explains some of the mechanism between pain and participation restriction). The “potential mediators” were the variables from the parsimonious model.

The relationship between the number of pain sites and self-care restriction were adjusted for each separately.

## 5.5 Results

### 5.5.1 Prevalence and age and gender distribution of restricted self-care

At baseline, 7725 (98%) responders completed the KAP self-care item in returned questionnaires (n=7878). 886 (11.5%; 95% confidence interval (CI): 10.8, 12.2) participants reported restricted self-care. There was a linear trend with increasing age ( $p \leq 0.001$ ) and restriction was higher in women (% difference 2.3 %; CI: 0.8, 2.7) (Table 5.3).

**Table 5.3 The 4-week period prevalence of restricted self-care; overall and stratified by age and gender**

	Restricted in self-care
<b>Overall</b>	886 (11.5%)
<b>Age group</b>	
50-59	174 (7.0%)
60-69	216 (9.3%)
70-79	280 (14.2%)
80+	216 (22.9%)
<b>Gender</b>	
Female	540 (12.5%)
Male	346 (10.2%)

When stratified by gender in each age group it was noted that there were a high proportion of older females. However, the difference in the prevalence of self-care restriction between men and women aged 80+ was not significant (% difference 5.0 % (95% confidence interval -0.6, 10.3)).

#### *Age standardisation*

There were differences between the age and gender structures of NorStOP and the population of England and Wales. To remove the effect of differences in age, standardisation calculations were carried out using indirect standardisation. Calculations were done both for age groups and gender.

The observed rate of self-care restriction (overall) in adults aged 50 years and over in the general population (study population) was 11.5% and the standardised (England and Wales population) overall rate of restricted self-care was 11.2%. In females the observed prevalence of self-care was 12.4% and the standardised rate was 12.3%. The observed rate of prevalence in males was 10.2% and 9.8% in the standardised population (Table 5.4 & 5.5).

These results indicate that the observed prevalence rate of restricted self-care in the NorStOP population is generalisable to the population of adults aged 50 years and over in England and Wales based on the age and gender structure.

**Table 5.4** Standardisation of the prevalence of restricted self-care by age groups

	a	b	c	d	e
Age Group	NorStOP Population	Observed Restricted self-care	Observed prevalence b/a x 1000	England and Wales Population /1000	Expected Restricted self-care
<b>Females</b>					
50-59	1332	95	71	3306	234,726
60-69	1249	122	97	2487	241,239
70-79	1133	170	150	2152	322,800
80+	622	153	246	1475	362,850
Total	4336	540	564	9420	1,161,615
<b>Males</b>					
50-59	1158	79	68	3248	221,479
60-69	1066	94	88	2351	207,314
70-79	843	110	131	1677	218,861
80+	322	63	196	716	140,049
Total	3389	346	483	7992	787,703

**Table 5.5** The standardised prevalence of restricted self-care

Observed estimates in NorStOP		Standardised to England and Wales	
<b>Overall</b>		<b>Overall</b>	
11.5%		11.2%	
<b>Female</b>	<b>Male</b>	<b>Female</b>	<b>Male</b>
12.4%	10.2%	12.3%	9.8%

### 5.5.2 Factors associated with self-care restriction

#### *Associations between health conditions / impairments and restricted self-care*

##### i) Health conditions

All health conditions were significantly associated with restricted self-care in the univariable analysis. Adjusting for age and gender did not reduce the

strength of the relationship, suggesting age and gender did not confound the association between restricted self-care and number of health conditions (Table 5.6). There was a dose and response relationship between restricted self-care and an increase in numbers of conditions. (2-3 health conditions adj. OR: 3.1; 95% CI 2.5, 3.8). In the multivariable analysis, there was no significant relationship between health conditions and self-care restriction when adjusted for other associated factors.

#### ii) Impairments

All self-reported impairments were significantly associated with restricted self-care in the univariable analysis before and after adjusting for age and gender. There was a dose-response relationship between restricted self-care and increasing numbers of impairments (for 6-9 impairments adj. OR: 14.8; 95% CI 9.7, 22.4). When adjusted for all other factors associated with self-care in the multivariable analysis, these associations were no longer statistically significant (Table 5.6).

#### iii) Body mass index

Only obesity was significantly associated with restricted self-care in the univariable analysis after adjusting for age and gender (adj. OR: 2.1; 95% CI 1.7, 2.5). However, BMI was not associated with restricted self-care once adjusted for other variables in the multivariable analysis (Table 5.6).



#### iv) Anxiety

Anxiety was significantly associated with restricted self-care in the univariable analysis. There was a trend with self-care restriction and increasing anxiety; the odds of reporting restricted self-care was four and a half times higher in those with definite cases compared to non-cases (adj. OR for definite cases: 4.5; 95% CI 3.8, 5.4). When adjusted for other factors associated with restricted self-care, associations were no longer statistically significant (Table 5.6).

#### v) Depression

Depression was significantly associated with restricted self-care in the univariable analysis before and after adjusting for age and gender. There was a trend with self-care restriction and increasing depression. The odds of reporting self-care restriction were almost ten times higher for definite cases of depression comparing to non-cases (adj. OR for definite cases: 9.8; 95% CI 8.0, 11.9). Associations remained statistically significant in the multivariable analysis (adj. OR for definite cases: 3.3; CI 2.4, 4.6) (Table 5.6).

#### vi) Cognitive complaint

Cognitive complaint was significantly associated with restricted self-care in the univariable analysis. The odds of reporting self-care restriction in those individuals with high levels of cognitive impairment were five times higher than those who had scored zero (none) (adj. OR for high levels: 5.1; 95% CI 4.3, 6.2). Adjusting for other variables associated with restricted self-care reduced the odds of reporting restricted self-care in those with both high and low levels

of cognitive impairment, but associations remained statistically significant (adj. OR for high levels: 1.7; CI 1.3, 2.3) (Table 5.6).

vi) Peripheral joint pain

Restricted self-care was significantly associated with the site, number, and pattern of peripheral joint pain in the univariable analysis. A dose-response relationship was observed between comorbid joint pain and self-care restriction. The odds of reporting self-care restriction increased with the increasing number of peripheral joint pain, and when multiple sites of joint pain were reported (adj. OR: 5.0; 95% CI 3.9, 6.5 for four joint pains). (adj. OR for Multiple LL: 2.7; CI 1.6, 2.9 and adj. OR for Hand+LL: 3.0; CI 2.4, 3.8). Therefore only the number of peripheral joint pain was included in the multivariable analysis. However, associations with restricted self-care were not significant when adjusted for other factors (Table 5.7).

vii) Bodily pain (pain elsewhere other than hand, hip, knee or foot pain)

Restricted self-care was significantly associated with bodily pain in the univariable analysis. There was a dose-response relationship between bodily pain and restricted self-care. The odds of reporting self-care restriction in those individuals who reported ten or more sites of pain elsewhere were five times higher than those who reported no bodily pain (adj. OR: 5.1; 95% CI 4.0, 6.5). In the multivariable analysis there was no significant relationship between bodily pain and restricted self-care (Table 5.7).

### *Associations between activity limitation and restricted self-care*

The strongest association with self-care restriction in the univariable analysis was with activity limitation. There was a positive trend between the association with restricted self-care and activity limitation. The odds of reporting self-care restriction in people with highest activity limitation scores were 13 times higher than those with lowest levels of activity limitation (adj. OR: 13.3 95% CI: 9.2, 19.1). Associations remained statistically significant in the 2<sup>nd</sup> highest and highest categories after adjusting for other variables in the multivariable analysis (adj. OR for highest: 7.6 95%CI: 4.2, 13.6) (Table 5.8).

### *Associations between contextual factors and restricted self-care*

#### i) Socio-demographic factors

##### *Age*

Age was significantly associated with restricted self-care in the univariable analysis before and after adjusting for gender. There was a positive trend with increase in age and restricted self-care. Strongest associations occurred in those who were 80 years of age and older (adj. OR 3.9; 95% CI 3.1, 4.8). Associations with age and restricted self-care remained significant in the multivariable analysis for those 70 years and older (adj. OR for 80+: 1.7; CI 1.2, 2.5) (Table 5.9).

##### *Gender*

Female gender was significantly associated with restricted self-care before and after adjusting for age in the univariable analysis, albeit the association

was not strong (adj. OR: 1.2 95% CI 1.0, 1.3). Gender was not significantly associated with restricted self-care in the multivariable analysis once adjusted for other associated factors (Table 5.9).

## ii) Socio-economic factors

### *Occupational class*

Occupational class was significantly associated with restricted self-care in the univariable analysis after adjusting for age and gender. The odds of reporting restricted self-care in those who were in the manual jobs category were twice as high compared to those in the non-manual jobs category (adj. OR 2.3; 95% CI 1.9, 2.8). Associations were not statistically significant in the multivariable analysis once adjusted for other factors in the model (adj. OR 1.2; CI 0.7, 2.0) (Table 5.9).

### *Educational attainment*

Educational attainment was significantly associated with restricted self-care in the univariable analysis. The odds of reporting restricted self-care in those with no adult qualifications was twice as high comparing to those who gained adult qualifications (adj. OR: 2.3 95% CI 1.9, 2.8). Associations reduced but remained statistically significant in the multivariable analysis (adj. OR: 1.5 CI 1.1, 2.0) (Table 5.9).

### *Perceived adequacy of income*

Perceived adequacy of income was significantly associated with restricted self-care in the univariable analysis before and after adjusting for age and

gender (adj. ORs for strain: 9.5; 95% CI 6.2, 14.5). In the multivariable analysis, associations remained statistically significant for those who had to 'be careful' and find it 'strain' to get by (adj. ORs for strain: 2.6; CI 1.3, 5.0) (Table 5.9).

### iii) Social networks

#### *Social networks*

Social network scores were significantly associated with restricted self-care in the univariable analysis before and after adjusting for age and gender. The odds of reporting self-care restriction was three times higher in those with 'low' social network scores compared to those with 'high' scores (adj. OR: 2.7; 95% CI 1.9, 2.8). However, associations were not statistically significant when adjusted for other factors in the multivariable analysis (Table 5.9).

#### *Having a confidant*

Having a confidant was significantly associated with restricted self-care before and after adjusting for age and gender, albeit the association was not strong (adj. OR: 1.4; 95% CI 1.1, 1.8). Associations were not statistically significant when adjusted for other factors in the multivariable analysis (Table 5.9).

**Table 5.6 Associations between restricted self-care and selected health conditions and impairments; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care**

Health conditions & Impairments	Total n	Restricted n (%)	Crude (95%CI)	ORs	Adj. OR (95% CI) (Age and Gender)	Adj. OR (95% CI) for all other variables
<b>Health conditions</b>						
0	4872	392 (8.0)	1		1	◇ Removed due to non-significant association
1	2160	325 (15.0)	2.0 (1.7, 2.4)		1.9 (1.6, 2.2)	
2-3	693	169 (24.4)	3.7 (3.0, 4.5)		3.1 (2.5, 3.8)	
<b>Impairments</b>						
0	1432	42 (2.9)	1		1	◇ Removed due to non-significant association
1-3	4509	392 (8.7)	3.2 (2.3, 4.4)		2.8 (2.0, 3.9)	
4-5	1562	368 (23.6)	10.2 (7.3, 14.2)		8.1 (5.8, 11.2)	
6-9	222	84 (37.8)	20.1 (13.4, 30.3)		14.8 (9.7, 22.4)	
<b>Body Mass Index (BMI)+</b>						
20 – 24.99	2695	273 (10.1)	1		1	◇ Removed due to non-significant association
25 – 29.99	3016	288 (9.5)	0.9 (0.8, 1.1)		1.1 (0.9, 1.3)	
<20	345	57 (16.5)	1.8 (1.2, 2.4)		1.3 (0.9, 1.8)	
>30	1316	205 (15.6)	1.6 (1.3, 2.0)		2.1 (1.7, 2.5)	
<b>HAD + anxiety score</b>						
None case						◇ Removed due to non-significant association
Borderline	4486	302 (6.7)	1		1	
Definite	1645	228 (13.9)	2.2 (1.9, 2.7)		2.3 (1.9, 2.7)	
	1393	317 (22.8)	4.1 (3.4, 4.8)		4.5 (3.8, 5.4)	
<b>HAD + depression score</b>						
Non cases	5833	335 (5.7)	1		1	1
Borderline	1046	255 (24.4)	5.2 (4.4, 6.3)		4.9 (4.1, 5.9)	2.2 (1.6, 2.9)
Definite	662	260 (39.3)	10.6 (8.8, 12.8)		9.8 (8.0, 11.9)	3.3 (2.4, 4.6)
<b>Cognitive complaint ++</b>						
None	3863	193 (5.0)	1		1	1
Low levels	1559	170 (10.9)	2.3 (1.9, 2.8)		2.2 (1.8, 2.7)	1.6 (1.1, 2.2)
High levels	1803	426 (23.6)	5.9 (4.9, 7.1)		5.1 (4.3, 6.2)	1.7 (1.3, 2.3)

\*Numbers do not add up due to missing data

+BMI (20–24.99:Normal, 25–29.99:Overweight <20:Underweight >30:Obese

+ Hospital Anxiety and Depression Scale (Non cases: 0-7, Borderline cases: 8-10, Definite cases: 11+)

++ Functional Limitations Profile, Cognitive and Alertness behaviour subscale (None: 0, high levels: 22.5-100, low levels: 0.1-22.4)

◇ Removed in the backward-stepwise logistic regression due to non-significance

**Table 5.7 Associations between restricted self-care and selected health conditions and impairments; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care in the univariable analysis**

Health conditions & Impairments	Total n■	Restricted n (%)	Crude ORs (95%CI)	Univariable Associations Adj. ORs (95% CI)	▲ Multivariable Associations Adj. ORs (95% CI)
<b>Site of peripheral joint pain</b>					
Hand pain					
NO	4128	329 (8.0)	1	1	~ Not included in the analysis
YES	3379	519 (6.9)	2.0 (1.8, 2.4)	2.0 (1.7, 2.4)	
Hip pain					
NO	4943	411 (8.3)	1	1	-
YES	2558	427 (16.7)	2.2 (1.9, 2.5)	2.1 (1.8, 2.5)	
Knee pain					
NO	3536	267 (7.6)	1	1	-
YES	3995	583 (14.6)	2.0 (1.8, 2.4)	2.0 (1.7, 2.3)	
Foot pain					
NO	4553	347 (7.6)	1	1	-
YES	2959	496 (16.8)	2.4 (2.1, 2.8)	2.3 (2.0, 2.7)	
<b>Number of peripheral joint pain</b>					
0	1730	102 (5.9)	1	1	◇ Removed due to non-significant association
1	1767	120 (6.8)	1.2 (0.9, 1.5)	1.2 (0.9, 1.5)	
2	1622	171 (10.5)	1.9 (1.5, 2.4)	1.9 (1.4, 2.4)	
3	1275	197 (15.5)	2.9 (2.3, 3.7)	2.7 (2.1, 3.5)	
4	880	218 (24.8)	5.3 (4.1, 6.8)	5.0 (3.9, 6.5)	
<b>Pattern of peripheral joint pain</b>					
No joint pain	1730	102 (5.9)	1	1	~ Not included in the analysis
Hand only	403	25 (6.2)	1.0 (0.7, 1.6)	1.0 (0.7, 1.6)	
Single LL*	1364	95 (7.0)	1.2 (0.9, 1.6)	1.2 (0.9, 1.6)	
Multiple LL	947	121 (12.8)	2.3 (1.8, 3.1)	2.7 (1.6, 2.9)	
Hand+ LL	2380	465 (46.4)	3.1 (2.5, 3.9)	3.0 (2.4, 3.8)	-
<b>Bodily Pain</b>					
0	2675	193 (7.2)	1	1	◇ Removed due to non-significant association
1-4	2588	247 (9.5)	1.4 (1.1, 1.6)	1.4 (1.1, 1.7)	
5-9	1232	192 (15.6)	2.4 (1.9, 2.9)	2.5 (2.0, 3.1)	
10+	595	155 (26.1)	4.5 (3.6, 5.7)	5.1 (4.0, 6.5)	

\*LL (lower limb)

~ Not included due to high correlation with another variable in the multivariable analysis

■ n= for univariable associations

▲ denominator (overall n=) for multivariable analysis was n=4357

◇ Removed in the backward-stepwise logistic regression due to non-significance

**Table 5.8**      **Associations between restricted self-care and activity limitation;**  
**Univariable associations adjusted for age and gender and**  
**Multivariable associations adjusted for all other variables**  
**associated with restricted self-care in the univariable analysis**

<b>Activity Limitation</b>	<b>Total n■</b>	<b>Restricted n (%)</b>	<b>Crude ORs (95%CI)</b>	<b>Univariable Associations Adj. ORs (95% CI)</b>	<b>▲ Multivariable Associations Adj. ORs (95% CI)</b>
<b>Activity limitation score*</b>					
lowest	1318	34 (2.5)	1	1	1
3rd high	1953	60 (3.0)	1.2 (0.8, 1.8)	1.4 (0.7, 1.7)	1.4 ( 0.8, 2.7)
2nd high	1732	126 (6.8)	2.8 (1.9, 4.1)	2.6 (1.7, 3.8)	1.9 (1.0, 3.5)
highest ≥35	1437	565 (28.2)	15.2 (10.7,21.7)	13.3 (9.2, 19.1)	7.6 (4.2, 13.6)

\* PF-10 Physical functioning score: lowest (90.1-100), 3<sup>rd</sup> highest (70.1-90.0), 2<sup>nd</sup> highest (35.1-70.0), Highest (≥35.0)

■ n= for univariable associations

▲ denominator (overall n=) for multivariable analysis was n=4357



**Table 5.9 Associations between restricted self-care and contextual factors; Univariable associations adjusted for age and gender and Multivariable associations adjusted for all other variables associated with restricted self-care in the univariable analysis**

Contextual factors	Total n	Restricted n (%)	Crude ORs (95%CI)	Univariable Associations Adj. ORs (95% CI)	▲ Multivariable Associations Adj. ORs (95% CI)
Age					
50-59	2316	174 (7.5)	1	1	1
60-69	2099	216 (10.2)	1.4 (1.1, 1.7)	1.4 (1.1, 1.7)	1.1 (0.8, 1.5)
70-79	1696	280 (16.5)	2.2.(1.8, 2.7)	2.2 (1.8, 2.7)	1.4 (1.0, 1.9)
80+	728	216 (29.6)	4.0 (3.2, 5.0)	3.9 (3.1, 4.8)	1.7 (1.2, 2.5)
Gender					
Male	3043	346 (11.3)	1	1	◇ Removed due to non-significant association
Female	3796	540 (14.2)	1.3 (1.1, 1.4)	1.2 (1.0, 1.3)	
Occupational Class					
Non-manual	2094	120 (5.7)	1	1	◇ Removed due to non-significant association
Manual	4530	590 (13.0)	2.5 (2.0, 3.0)	2.3 (1.9, 2.8)	
Higher Educational attainment *					
Yes	2181	114 (5.2)	1	1	1
No	5345	731 (13.7)	2.9 (2.3, 3.6)	2.3 (1.8, 2.8)	1.5 (1.1, 2.0)
Perceived Adequacy of Income					
Comfortable	1065	41 (3.8)	1	1	1
Little difficulty	3044	256 (8.4)	2.3 (1.6, 3.2)	2.2 (1.6, 3.0)	1.5 (0.9, 2.5)
Be careful	3115	459 (14.7)	4.3 (3.1, 6.0)	4.2 (3.0, 5.8)	2.1 (1.3, 3.4)
Strain	295	67 (22.7)	7.3 (4.9, 11.1)	9.5 (6.2, 14.5)	2.6 (1.3, 5.0)
Social Networks					
High	1249	72 (5.8)	1	1	◇ Removed due to non-significant association
Medium high	682	56 (8.2)	1.5 (1.0, 2.1)	1.4 (1.0, 2.0)	
Medium	2663	274 (10.3)	1.9 (1.4, 2.5)	1.7 (1.3, 2.2)	
Low	1529	264 (17.3)	3.4 (2.6, 4.5)	2.7 (2.0, 3.6)	
Have a confidant					
YES	6839	751 (11.0)	1	1	◇ Removed due to non-significant association
NO	724	107 (14.8)	1.4 (1.1, 1.8)	1.4 (1.1, 1.8)	

\* Educational attainment yes = Have adult qualifications, no= Do not have adult qualifications

■ n= for univariable associations

▲ denominator (overall n=) for multivariable analysis was n=4357

◇ Removed in the backward-stepwise logistic regression due to non-significance

### **5.5.3 Derivation of the parsimonious model of key factors associated with restricted self-care**

A parsimonious model of key factors associated with restricted self-care was derived from the multivariable analysis using backward-stepwise logistic regression. The final fitted model (n=6506) included key variables independently associated with restricted self-care, which were age, activity limitation, depression, cognitive alertness, perceived adequacy of income and educational attainment.

Strongest associations occurred between restricted self-care and the highest activity limitation score (adj. OR: 7.6; 95% CI 4.2, 13.6). This was followed by definite cases of depression (adj. OR: 3.8; CI 3.0, 4.9) and inadequacy of income (adj. OR for strain: 2.5; CI 1.5, 4.1). The risk of reporting self-care restriction for those with high levels of cognitive impairment was nearly twice as high as those with no cognitive impairment (adj. OR high levels: 1.8; CI 1.4, 2.3). Age was also independently associated with restricted self-care. Those who were 80 years and older had twice the odds of reporting self-care restriction than those who were up to 59 years of age (adj. OR for 80+: 1.9; CI 1.4, 2.5). Those who did not have adult qualifications were also more likely to have self-care restrictions even after adjusting for all other factors. However the association was not as strong as in other key factors (adj. OR: 1.1; CI 1.0, 1.5) (Table 5.10).

**Table 5.10 Final (re-fitted) parsimonious model of key factors associated with restricted self-care in older adults in the general population**

Overall n=6506	Total n	Restricted n (%)	Adjusted ORs for all covariates (95% CI)
<b>Age</b>			
50-59	2229	174 (7.8)	1
60-69	1991	216 (10.8)	1.3 (1.0, 1.7)
70-79	1599	280 (17.5)	1.6 (1.2, 2.0)
80+	687	216 (31.4)	1.9 (1.4, 2.5)
<b>Activity limitation</b>			
lowest (90.1-100)	1266	34 (2.6)	1
3rd highest (70.1- 90)	1864	60 (3.2)	1.1 (0.7, 1.8)
2nd highest (35.1-70)	1645	126 (7.6)	1.9 (1.1, 3.0)
highest ≥35	1731	565 (32.6)	5.6 (3.6, 8.7)
<b>Depression†</b>			
Non cases (0-7)	5105	335 (6.6)	1
Borderline cases (8-10)	857	255 (29.7)	2.1 (1.7, 2.6)
Definite cases (≥11)	544	260 (47.8)	3.8 (3.0, 4.9)
<b>Cognitive complaint ††</b>			
None (0)	3531	193 (5.4)	1
low levels (0.1-22.4)	1410	170 (12.0)	1.5 (1.2, 1.9)
high levels (22.5-100)	1565	426 (27.2)	1.8 (1.4, 2.3)
<b>Perceived adequacy of income</b>			
Comfortable	943	41 (4.3)	1
Little difficulty	2659	256 (9.6)	1.6 (1.1, 2.4)
Be careful	2658	459 (17.2)	2.1 (1.4, 3.0)
Strain	246	67 (27.2)	2.5 (1.5, 4.1)
<b>Higher educational attainment</b>			
Yes	1968	114 (5.8)	1
No	4538	731 (16.1)	1.1 (1.0, 1.5)

† Hospital Depression Score (HAD)

†† Functional Limitations Profile, Cognitive and Alertness behaviour subscale

### *Validation of the parsimonious model*

#### i) Internal validity

The apparent performance of the parsimonious model was determined using the original baseline sample of NorStOP1 (n=6506). The Hosmer and Lemeshow goodness of fit test (HL) suggested that the parsimonious model was not a good fit for the data (Pearson's  $X^2 = 684.20$ ,  $p = 0.0007$ ). However, the area under the ROC curve was 0.83 (95% CI 0.81, 0.85), which indicated an 83% probability of those with self-care restriction having higher predicted probability than those without self-care restriction, for a random pair of cases with or without self-care restriction. Bootstrapping improved the goodness of fit significantly (Pearson's  $X^2 = 284.36$ ,  $p = 1.00$ : c-index= 0.83 (CI 0.81, 0.85), and the split-sample performance yielded similar, but improved results to the apparent performance of the model (Pearson's  $X^2 = 546.22$ ,  $p = 0.07$ : c-index= 0.82 (CI 0.80, 0.84)). The expected optimism of the model was 0.01, suggesting that the parsimonious model had very low levels of optimism. The optimism corrected performance of the model was 0.82, proposing that this model has a reasonable power in predicting self-care restriction in those aged 50 and over in the general population (Table 5.11).

**Table 5.11 Internal validity (NorStOP<sup>1</sup>) of the parsimonious model**

Overall n= 6506		Apparent ORs 95% CI	Bootstrap ORs 95% CI	Split-sample ORs 95% CI
<b>Age groups</b>				
	50-59	1	1	1
	60-69	1.3 (1.0, 1.7)	1.3 (1.0, 1.7)	1.6 (1.9, 2.3)
	70-79	1.6 (1.2, 2.0)	1.6 (1.2, 2.0)	1.7 (1.3, 2.4)
	80+	1.9 (1.4, 2.5)	1.9 (1.4, 2.5)	2.0 (1.4, 2.9)
<b>Activity limitation</b>				
	lowest	1	1	1
	3 <sup>rd</sup> highest	1.1 (0.7, 1.8)	1.1 (0.7, 1.8)	1.0 (0.5, 1.6)
	2 <sup>nd</sup> highest	1.9 (1.2, 3.0)	1.9 (1.2, 3.0)	1.5 (1.0, 2.5)
	highest	5.6 (3.6, 8.7)	5.9 (3.7, 9.3)	4.8 (2.9, 7.9)
<b>Depression +</b>				
	Non cases (0-7)	1	1	1
	Borderline (8-10)	2.1 (1.7, 2.6)	2.1 (1.7, 2.6)	2.0 (1.5, 2.6)
	Definite (≥11)	3.8 (3.0, 4.9)	3.8 (3.0, 5.0)	3.2 (2.3, 4.4)
<b>Cognitive complaint ++</b>				
	None (0)	1	1	1
	low levels (0.1-22.4)	1.5 (1.1, 1.9)	1.5 (1.1, 1.9)	1.6 (1.2, 2.1)
	high levels (22.5-100)	1.8 (1.4, 2.3)	1.8 (1.4, 2.3)	1.9 (1.4, 2.6)
<b>Perceived adequacy of Income</b>				
	Comfortable	1	1	1
	Little difficulty	1.6 (1.1, 2.4)	1.6 (1.1, 2.4)	2.1 (1.2, 3.5)
	Be careful	2.1 (1.4, 3.0)	2.1 (1.4, 3.1)	2.7 (1.6, 4.5)
	Strain	2.5 (1.5, 4.2)	2.5 (1.5, 4.3)	3.8 (2.0, 7.3)
<b>Higher educational attainment</b>				
	Yes	1	1	1
	No	1.1 (1.0, 1.5)	1.1 (0.8, 1.6)	1.1 (0.7, 1.7)
<b>Goodness of fit test</b>				
	Pearson X <sup>2</sup>	684.20	284.36	546.22
	P value	≤0.001	1.00	0.07
	c-index	0.83 (0.81, 0.85)	0.83 (0.81, 0.85)	0.82 (0.80, 0.84)

+ Hospital Depression Score (HAD)

++ Functional Limitations Profile, Cognitive and Alertness behaviour subscale

## ii) External validity

The external validity of the parsimonious model was tested on the NorStOP<sup>2</sup> dataset with sample size of n=5022. In the apparent performance HL test showed that the model was a reasonable fit for the data (Pearson's  $X^2 = 531.66$ ,  $p = 0.60$ ). The area under the ROC curve was 0.83 (95% CI 0.79, 0.83), again suggesting the model had a good discriminative power in identifying those with high and low risk of reporting self-care restriction. Bootstrapping improved how the model fitted the data (Pearson's  $X^2 = 309.60$ ,  $p = 1.00$ : c-index= 0.82 (CI 0.79, 0.83)) and the split-data sample method had similar results to the apparent model, albeit the model calibration was even lower (Pearson's  $X^2 = 471.89$ ,  $p = 0.34$ : c-index= 0.80 (CI 0.77, 0.82)).

The expected optimism of the model was 0.02, suggesting that the model was not optimistic. The optimism corrected performance of the model was 0.81, indicating that the external validity of the parsimonious model was similar to the model tested on the original data (internal validation). Thus, this model had a reasonable predictive power on different population sample, and therefore the model could be used in other populations to estimate restricted self-care in adults aged 50 years and over in the general population (Table 5.12).

**Table 5.12 External validity (NorStOP<sup>2</sup>) scores of the parsimonious model**

Overall n= 5022		Apparent ORs 95% CI	Bootstrap ORs 95% CI	Split-sample ORs 95% CI
<b>Age groups</b>				
	50-59	1	1	1
	60-69	1.2 (1.0, 1.6)	1.2 (1.0, 1.7)	1.4 (1.0, 2.0)
	70-79	1.3 (1.0, 1.8)	1.3 (1.0, 1.8)	1.4 (1.0, 2.1)
	80+	1.8 (1.3, 2.5)	1.8 (1.3, 2.6)	2.0 (1.4, 3.0)
<b>Activity Limitation</b>				
	lowest	1	1	1
	3 <sup>rd</sup> highest	1.0 (1.0, 1.6)	1.0 (1.0, 1.6)	1.0 (0.6, 1.8)
	2 <sup>nd</sup> highest	1.7 (1.1, 2.7)	1.7 (1.1, 2.7)	1.6 (1.0, 2.7)
	highest	4.8 (3.1, 7.5)	4.8 (3.1, 7.7)	4.1 (2.4, 6.8)
<b>Depression †</b>				
	Non cases (0-7)	1	1	1
	Borderline (8-10)	2.1 (1.6, 2.7)	2.1 (1.6, 2.7)	2.1 (1.5, 3.0)
	Definite (≥11)	4.6 (3.5, 6.2)	4.6 (3.5, 6.2)	4.5 (3.1, 6.3)
<b>Cognitive complaint ††</b>				
	None (0)	1	1	1
	low levels (0.1-22.4)	1.5 (1.1, 2.0)	1.5 (1.1, 2.0)	1.4 (1.0, 2.0)
	high levels (22.5-100)	1.5 (1.2, 2.0)	1.5 (1.2, 2.0)	1.3 (1.0, 1.8)
<b>Perceived adequacy of Income</b>				
	Comfortable	1	1	1
	Little difficulty	1.0 (0.7, 1.4)	1.0 (0.7, 1.4)	1.1 (0.7, 1.8)
	Be careful	1.5 (1.0, 2.9)	1.5 (1.0, 2.1)	1.5 (1.0, 2.3)
	Strain	1.8 (1.1, 3.0)	1.8 (1.1, 3.0)	2.4 (1.3, 4.3)
<b>Higher educational attainment</b>				
	Yes	1	1	1
	No	1.0 (0.7, 1.5)	1.0 (0.7, 1.5)	1.0 (0.6, 1.4)
<b>Goodness of fit test</b>				
	Pearson X <sup>2</sup>	531.66	309.60	471.89
	P value	0.60	1.00	0.34
	c-index	0.83 (0.79, 0.83)	0.82 (0.79, 0.83)	0.80 (0.77, 0.82)

† Hospital Depression Score (HAD)

†† Functional Limitations Profile, Cognitive and Alertness behaviour subscale

#### **5.5.4 Factors explaining the relationship between joint pain and self-care restriction**

Univariable analysis showed strong associations between joint pain and restricted self-care. However joint pain was not independently associated with restricted self-care and was not included in the parsimonious model.

A multivariable logistic regression was fitted to examine the associations between joint pain and self-care restriction adjusted for each factor from the parsimonious model to assess for potential mediation. The association between the number of joint pains and restricted self-care remained after adjustment for each factor. The association between the number of joint pains attenuated most with adjustment for activity limitation and the depression. There was no attenuation when adjusted for age and educational attainment (Table 5.13).



**Table 5.13 Multivariable analysis of the associations of joint pain and restricted self-care adjusted for parsimonious factors**

		Association with restricted self-care ORs (95% CI)
<b>Peripheral Joint Pain by number</b> Crude n=7274	0	1
	1	1.2 (0.9, 1.5)
	2	1.9 (1.5, 2.4)
	3	2.9 (2.3, 3.7)
	4	5.3 (4.1, 6.8)
<b>Peripheral Joint Pain by number</b> Adjusted for age n=7274	0	1
	1	1.2 (0.9, 1.5)
	2	1.8 (1.4, 2.4)
	3	2.7 (2.1, 3.5)
	4	5.0 (3.9, 6.5)
<b>Peripheral Joint Pain by number</b> Adjusted for activity limitation n=6890	0	1
	1	0.8 (0.6, 1.1)
	2	0.9 (0.7, 1.2)
	3	1.0 (0.8, 1.4)
	4	1.3 (1.0, 1.8)
<b>Peripheral Joint Pain by number</b> Adjusted for depression n=7116	0	1
	1	1.1 (0.8, 1.5)
	2	1.6 (1.2, 2.0)
	3	2.0 (1.5, 2.6)
	4	2.9 (2.2, 3.8)
<b>Peripheral Joint Pain by number</b> Adjusted for cognitive alertness n=6891	0	1
	1	1.1 (0.8, 1.4)
	2	1.4 (1.1, 1.9)
	3	1.9 (1.5, 2.5)
	4	3.2 (2.5, 4.3)
<b>Peripheral Joint Pain by number</b> Adjusted for perceived adequacy of income n=7096	0	1
	1	1.1 (0.9, 1.5)
	2	1.8 (1.4, 2.3)
	3	2.7 (2.1, 3.5)
	4	4.3 (3.3, 5.6)
<b>Peripheral Joint Pain by number</b> Adjusted for educational attainment n= 7115	0	1
	1	1.2 (0.9, 1.6)
	2	1.9 (1.5, 2.5)
	3	2.9 (2.3, 3.8)
	4	5.3 (4.1, 6.9)

### **5.5.5 Summary of results**

The overall prevalence of restricted self-care was 11.5% in community-dwelling older adults in the general population. There was a linear trend with increasing age, and self-care restrictions were higher in women.

Self-reported health conditions, impairments, activity limitation and contextual factors were significantly associated with restricted self-care in the univariable analysis, after adjusting for age and gender. The strongest associations occurred with self-reported impairments, activity limitation and depression. Restricted self-care was associated with each peripheral joint pain; the association with joint pain increased with the increasing number of joint pains. However joint pain was not independently associated with self-care restriction once adjusted for other factors. This is because it co-occurs with other health and socio-demographic factors that are strongly associated with self-care, such as activity limitation and depression. Age, depression, cognitive alertness, activity limitation, educational attainment and perceived adequacy of income were included in a parsimonious set of statistically significant and independently associated factors with restricted self-care. This model was validated using bootstrapping and split-data methods both for internal and external validation. The optimism corrected performance of the model proposed that this model had a reasonable power in predicting self-care restriction in those aged 50 years and over in the general population.

## 5.6 Discussion

### 5.6.1 Principal findings

#### *Prevalence and distribution of person-perceived self-care restriction*

One in ten community-dwelling older adults in the general population was restricted in self-care. Prevalence increased with age, and was higher in women. Individuals who are older and with higher levels of self-care restriction may not have been able to complete this self-complete postal questionnaire due to their severe activity limitation and / or psychological impairments. Age standardisation of the observed prevalence rates of the study to the England and Wales population confirmed that the study was generalisable to the wider national population (observed: 11.5% cf standardised: 11.2%). However, the generalisability of the results to other populations may be limited by differences in the socio-economic characteristics and contextual factors, which may vary across different populations. The study population resided in urban areas of the North Staffordshire area in England, where the local population is highly disadvantaged, both in terms of health deprivation and employment, when compared to the rest of the country. This is partially due to the historic decline of traditional industries (Comprehensive Area Assessment, 2009). More than half of the Stoke-On-Trent population were reported to be in the most deprived national quintile (Association of Public Health Observatories, 2010).

7725 people completed the self-care item out of 7878 participants at baseline. Those who did not complete the self-care item (1.9%) did not differ from those who completed the self-care item in terms of age and gender. As this was a small proportion of the sample, the potential effects of the missing data in the statistical analysis could be considered to be minor.

The study population was highly heterogeneous (99.4% white). Consequently the results of this study could not be generalised to other ethnic groups. However, there is no empirical evidence to suggest that the prevalence of restricted self-care might be different in other ethnic groups or in rural populations.

#### *Associated factors with restricted self-care in older people*

The study findings provide empirical support for the ICF model of participation restriction. Restrictions in self-care were associated with bodily, personal and contextual factors. Activity limitation measured by PF-10 scale had the strongest association with restricted self-care both in the univariable and multivariable analysis. However, there were previous concerns about linking the measurement components of PF-10 (physical functioning scale) to the ICF's domain of activity limitation in population surveys. Previous studies questioned the measurement validity of PF-10, suggesting that this instrument does not necessarily differentiate between the participation restriction and activity limitation domains of ICF (Ayis et al., 2010; Pollard et al., 2006; Wilkie, 2005). Also, the PF-10 had items such as measuring limitation in bathing or dressing, which would expectedly have high correlations with self-care restriction. Nevertheless, in this secondary analysis of a

large dataset, the PF-10 was the most suitable available instrument to measure activity limitation. Although there may be some correlations between the measurement properties of PF-10 and the KAP self-care item, it is appreciated that the ICF framework implies that activity limitation is one component that may lead to participation restriction. Although this study does not completely confirm this it does suggest that for those who are restricted in self-care, most have activity limitation and indicates that this may be a key mechanism for restriction.

The strong associations with depression, cognitive alertness and perceived adequacy of income could be the result of negative reporting bias, which is a major problem in the assessment of health care outcomes (McGauran et al., 2010). Responders with poorer backgrounds and depression may generally be less satisfied with their quality of life (Moore et al., 2005) and have a negative perception of their needs being met 'as and when they have wanted' (Wilkie, 2005). Previous research also suggested that cognitive impairment and depression co-occurs with pain (Sherbourne et al, 2009; Fishbain et al., 1997; Magni et al., 1993).

Those who perceived their incomes as inadequate were nearly three times more likely to report self-care restriction compared to those who were comfortable with their income even after adjusting for all other health and contextual factors in the multivariable analysis. Social disadvantages have been previously associated with disability in those with chronic conditions (Grundy, 2000). It was reported that failure to afford a reasonable standard of living could have an impact on an

individual's health status (i.e. diet, housing, access to health and social services) and may result in an increase of reporting psychological disorders such as anxiety and depression (Emerson et al., 2006). In a recently published report, the equality profile of older people with high support needs was examined, and this highlighted that a higher proportion of people who used support services were from disadvantaged backgrounds (Blood and Bamford, 2010). It is important to acknowledge that given the two key factors independently associated with restricted self-care are socio-economic factors in this study, it would be reasonable to conclude these factors play an important role in the participation restriction process.

Age was another factor that remained significantly associated with restricted self-care in the parsimonious model after adjusting for other key factors. Restriction was higher in the oldest age group (80+). This supports the literature review conducted in Chapter 1 which suggests that ageing population is associated with increase in prevalence of chronic health conditions and disability (Denton and Spencer, 2010; Martin et al., 2010; Nusselder et al., 2006; Wilkie et al, 2006; Gill et al, 2006:a).

A simple method proposed by Baron and Kenny (1986) was used to examine the potential mediation of the link between the number of joint pains and self-care restriction. Using this method, mediation was identified if the strength of association was reduced with the addition of a third variable. The limitations to this approach were that a formal test for mediation (e.g. Sobel's test) was not used and

newer techniques are now being used to examine how extensively mediation is being quantified (for example using structural equation modelling). However following the approach outlined by Baron and Kenny, the results suggest that the link between joint pain and restricted self-care in older people is potentially mediated by activity limitation and depression. Previous studies have highlighted that activity limitation (broadly defined in studies to include limitations in walking, bathing, transferring, dressing and personal care) is on the causal path both for joint pain and participation restriction (Arola et al., 2010; Corti and Rigon 2003) and this provides empirical support for the proposed ICF model of participation.

The strength of the association between joint pain and self-care restriction also attenuated with the addition of cognitive complaint and to a lesser extent with perceived adequacy of income. It could be hypothesised that, once older adults have pain, cognitive impairment or an inadequate income may prevent them from managing the joint problem or from adapting to continue to manage their self-care as they would like to. Further testing is required to explore these relationships further.

The elimination of health conditions and impairments from the parsimonious model is explained by the high correlation between these variables and remaining predictors in the model. Depression was highly correlated with anxiety, other health conditions and impairments in the 2X2 tables, and the same was true for the peripheral joint pain (by site, count and/or pattern) and activity limitation. It is important to recognise that the result of this analysis does not indicate that

variables taken out of the model are not important factors associated with restricted self-care in older people. Similarly, correlations between other health conditions and impairments and activity limitation could have contributed to the strength of relationship between activity limitation and restricted self-care as strong links were retained between these factors in the multivariable analysis.

### **5.6.2 Comparison with other studies**

Previous studies of self-care estimated the prevalence of limitations in older people to be in the region of 29% to 40% (Gill et al., 2009; Gill et al., 2006; a; Naik et al., 2004). Self-care was measured in this study as one form of participation and from the perspective of participants deciding if their needs were met “as and when they wanted”. Therefore, it is understood that an individual may experience task-specific activity limitation in self-care (e.g. difficulties in getting in and out of bath) but access to environmental facilitators may enable them to self-care (e.g. fitting a bath seat, handrails, or non-slip mats or receiving personal help and assistance). This may explain the gap between the observed prevalence estimates of self-care restriction when compared to other population studies of ADL limitations / dependency.

Age characteristics of the study sample had demonstrated agreement to previous studies' findings on the associations of increasing age with increased prevalence of chronic conditions and comorbidity (Palacios-Cena et al., 2012; Tsang et al., 2008; Cigolle et al., 2007; Schafer et al., 2002). This study observed strong links between co-morbid health, number of peripheral joint pains and restricted self-



care. The risk of self-care restriction increased with increasing numbers of peripheral joint pain. Similarly, the odds of reporting self-care restriction was increased in those with peripheral joint pain by pattern, when multiple sites of joint pain were taken into account. Previous research supports this finding by stating that dose-response deterioration in general health status is associated with each extra joint pain sites that is symptomatic (Dawson et al., 2004). Multiple joint pains might also result in an increased severity of pain, which was previously related to functional limitations (Van Dijk et al., 2008). Van Dijk and colleagues (2008) studied comorbidity, limitations in activities and pain in patients with osteoarthritis of the hip and knee, and found morbidity count and severity is associated with increased pain and limitations in activities. This cross-sectional study was conducted in the Netherlands, and participants (n=288) were recruited from three rehabilitation centres and two hospitals. However this study excluded those over the age of 85 years. Limitations in activities were measured using WOMAC and SF-36 physical functioning domains. Relationship between comorbidity and limitation in activities were determined using multivariate linear regression analysis. As opposed to this study, dependent variables were WOMAC and SF-36 physical functioning scores, and the Visual Analogue Scale (VAS) pain score. This study was characteristically different than the NorStOP study as it was conducted on a sample of participants that originated from Orthopaedic department listings as opposed to the general population. Further, participants mean age was 66; somewhat younger than the mean NorStOP sample age. However, findings from this thesis and Van Dijk et al's study were congruent. Both pointed out the

relationship between comorbidity and activity limitation in older adults with osteoarthritis.

A recent study into musculoskeletal comorbidities in cardiovascular disease, diabetes and respiratory disease aimed to quantify the contribution of comorbidity to activity limitations in populations (Slater et al., 2011). This study stated that comorbid arthritis significantly increased the risk of activity limitations. The study population recruited through the Canadian Community Health Survey was different to the NorStOP population in terms of demographic and socio-economic status; they were younger (20+ cf 50+), with higher income and education levels. Nevertheless, the results pointed to the same finding; that chronic musculoskeletal conditions play an important role in activity limitations in those with other conditions such as cardiovascular disease, diabetes and respiratory disease. An earlier study by Adamson and colleagues (2004) examined the association of chronic degenerative diseases with locomotor activity limitation and social participation among older British women. The study was a cross-sectional survey of 4286 women aged 60-79 years from 23 towns in England, Scotland and Wales. The prevalence of locomotor activity limitation and difficulties with social participation increased with increasing age and with increasing number of chronic diseases.

Depression, measured by the Hospital Anxiety and Depression Scale, was also strongly associated with restricted self-care in older people in this thesis. Previously it was reported that people with depressive thoughts and feelings are

more limited in activities and more restricted in participation (Muller, 2009; Bresling et al., 2006). Most previous research into depression has been conducted through longitudinal studies. A recent study by Machado et al. (2008) examined the role symptoms, activity limitation and depression play in social participation of older adults with osteoarthritis. Participants were 184 community-dwelling adults aged  $\geq 55$  years with physician diagnosis of osteoarthritis and they were interviewed at two time points 18 months apart. Measures of activity limitation and participation restrictions were derived by factor analysis of questions about difficulties experienced in daily life. This study found that the relationship between physical symptoms and difficulties in participation was partially mediated by activity limitation and depressive symptoms. Findings suggested that these factors act as a pathway to subsequent participation restrictions.

Another study investigated the relationship between major depression, subclinical depressive symptomology and activity limitation over a four year period in adults aged 18 to 60 years. A brief diagnostic interview was used to gather information on major depression, activity limitations, health conditions and other socio-demographic characteristics. This study found that major depression had a significant adverse impact and leads to limitation in daily activities (Breslin et al., 2006).

An earlier study carried out by Gureje and colleagues (2001) examined the persistent pain syndromes among primary care patients across 15 sites in 14 countries. In this study, 3197 randomly selected primary care patients completed

baseline and 12-month follow-up assessments of pain and depressive symptoms. This study found a strong and symmetrical relationship between persistent pain and depression and stated that the impairment of daily activities was the central component of this relationship (Gureje et al., 2001).

Similarly, the odds of reporting self-care restriction in those individuals with low levels of cognitive alertness score were five times higher than those who were classified as non-cases. The suggestion of strong associations between cognitive impairment and participation restriction was also advocated by previous research, although the direction of the causal association remains to be investigated further (Wilkie et al., 2007:a; Jagger et al., 2005). Buchner and Larson (1987) found that people with cognitive impairment were more susceptible to falls and at higher risk of developing mobility problems. It could be assumed that people with a high risk of falls may restrict activities such as bathing to avoid such risk. It was also suggested that cognitive impairment may also impede the individual's ability to make appropriate self-care decisions (Cameron et al., 2010).

### **5.6.3 Limitations**

This study was based on cross-sectional data. In cross-sectional studies, a purposeful sample of a population is used to estimate the relationship between an outcome of interest and population parameters as they are presented at a particular period of time (Porta, 2008). Since both the outcome and the variables are measured at a single time point, these studies cannot determine cause and effect relationships.

This study was based on self-reported data with varying time periods. It is possible, that some variables (e.g. pain in the last year) might be affected by information or recall bias (Porta, 2008). Similarly, mild problems such as pain, symptoms and activity limitation could also go unreported due to the responders' differing perceptions.

This study was a secondary analysis of the data collected for a general health survey that included questionnaires to investigate self-reported health conditions, activity limitations, participation restriction and contextual factors based around the individual's life. Whilst there were disadvantages to conducting a secondary analysis using this large dataset, (such as lack of familiarity with data, absence of key variables (e.g. environmental variables), and dealing with the complexity of data) there were many more advantages. First of all it was cost effective; it saved time as data was already collected. This allowed sufficient time for a detailed, intensive statistical analysis to answer complex questions. Secondly, it is based upon a high-quality dataset featuring a large and representative sample of general population. The survey response (71.3%) provided results which were representative of the target population (Mangione, 1995). Nevertheless the sample analysis highlighted differences in the non-responders and the responders in terms of age and gender, as there was a greater response from females and in the middle age groups (60-69 and 70-79) (Chapter 4). This could suggest that the oldest age group, which may be the one with most self-care restrictions according to the results of this study, may have been under represented in this survey. Therefore the observed prevalence of restricted self-care may actually be an

underestimation of the correct prevalence in the general population. It is possible that the non-response in the oldest age groups could be due to the study method of collecting data through a postal survey. It can be hypothesised that the prevalence of restricted self-care in individuals who were unable to complete the survey due to activity limitation and / or physical or psychological impairments could be significantly higher.

The parsimonious model was derived from 6506 participants, due to missing data (17%) in six covariates measured. The age and gender structure of these 6506 people were significantly different than the age and gender structure of those who were missing. Proportionately, those included in the parsimonious model were younger compared to those who responded at baseline but had missing covariate data ( $p \geq 0.001$ ), and had fewer women ( $p \geq 0.001$ ). Considering that this study established that self-care restriction increased with age and was higher in women based on the general population of older adults at baseline, associations between the key covariates identified with parsimonious model and self-care restriction may have been an underestimation due to missing data. The model was also solely based on a statistical decision, where inclusion and exclusion of predictors to the model was dependent on the statistical significance level. Rather strict significance values (0.05 for removal and 0.01 for re-entry cf 0.15 and 0.20 recommended by Hosmer and Lemeshow, 1989) were used in the multivariable model due to high levels of significance with all variables at the univariable analysis. This could have resulted in an elimination of other important factors from the model. However, the

internal and external validation of the parsimonious model suggested that the model had a reasonable predictive power.

## **5.7 Conclusions**

Person perceived self-care restriction is reported by one in ten persons over the age of 50 years in the general population and increases with age. It is associated with joint pain, although much of this relationship may be explained by other factors, such as activity limitation and contextual factors. To understand how self-care restriction may occur in older adults with joint pain, a focused analysis within this group is required. Prevalence increases with increasing age and is higher in women. Self-care restriction is associated with a number of factors including health conditions, impairments, activity limitation and contextual factors, providing support for the ICF model of participation; the proposed model of self-care in this thesis.

Whilst the findings reported in this chapter are useful for identifying how common self-care restrictions are in community-dwelling older people and associated factors, the cross-sectional analysis cannot ascertain patterns of occurrence and / or clearly identify potential predictors. The following chapters describe longitudinal analyses of self-care restriction.

**Table 5.14     Summary of key findings**

- This chapter reported the baseline cross-sectional analyses of the NorStOP, with an aim to estimate the prevalence of self-care restriction in older people in the general population, and determine the associations between self-care restriction and health conditions, impairments, activity limitation and other contextual factors.
- The prevalence of self-care restriction was 11.5% (95% CI: 10.8, 12.2). There was a linear trend with increasing age ( $p \leq 0.001$ ) and restriction was higher in women (% difference 2.3%; CI: 0.8, 2.7).
- The results of the age standardisation indicated that the observed prevalence of restricted self-care in the NorStOP population is similar to the estimated prevalence in England and Wales population.
- Self-reported health conditions, impairments, activity limitation and contextual factors were significantly associated with restricted self-care in the univariable analysis, after adjusting for age and gender.
- A parsimonious model of key factors associated with restricted self-care was derived from the multivariable analysis using backward-stepwise logistic regression. These key factors were age, activity limitation, depression, cognitive alertness, perceived adequacy of income and educational attainment.
- Joint pain was not independently associated with self-care restriction once adjusted for other factors, therefore was not identified as a key factor in the parsimonious model. One explanation for this is that joint pain co-occurs with other health and socio-demographic factors that are strongly linked to self-care.
- The internal and external validity of the parsimonious model was tested using the apparent, bootstrap and split-sample techniques and the model had a reasonable power in predicting self-care restriction in those aged 50 years and over in the general population.
- As cross-sectional analysis cannot ascertain patterns of occurrence and/or clearly identify potential predictors, following chapters describe longitudinal analysis of self-care restriction.



## **Chapter 6**

### **Person-perceived self-care restriction in the middle and old age: risk of onset and persistence over a 3 years period**

#### **6.1 Introduction**

The previous chapter described the prevalence of person-perceived self-care restriction and its relationship with joint pain in the general population through a set of cross-sectional analyses. The next stage in building a picture of the epidemiology of restricted self-care is to describe the ‘natural history’, in terms of onset and persistence of self-care restrictions in older people. Studying ‘natural history’ aims to yield information about the course of a disease or condition and generally requires longitudinal methods. Although ‘natural history’ is broadly defined as the course of a disease from pathological onset to resolution in epidemiology, these stages vary vastly across human conditions. Thus, more often than not the term ‘natural history’ refers to a model or framework that incorporates social and cultural interactions as well as the biological and healthcare processes involved in the disease process (Porta, 2008). In this thesis, self-care restriction is the health-related outcome of interest, not a disease or health condition. It is assumed in this chapter that the concept of ‘natural history’ can be applied also to such health-related states.

This is the first study to examine self-care as defined in the International Classification of Functioning (ICF) and there is no estimate of its natural history (i.e. in terms of the onset and persistence of self-care restriction). For comparison,

there are studies which have reported the rate of onset and persistence of dependency and / or loss of independence in self-care tasks; definitions that would be classed under the umbrella term of activities of daily living (ADLs) (e.g. dressing, eating, toileting) and instrumental activities of daily living (IADL) (e.g. meal preparation, grocery shopping, using the telephone, taking medications and managing money) in older adults. Despite conceptual differences between ADL and person-perceived self-care restriction these may nevertheless provide useful information on the natural history of self-care disability in general.

The Index of Independence in Activities of Daily Living, (Index of ADL), was developed by Katz and colleagues (1963) from observations of a large number of activities performed by a group of patients with fracture of the hip, to use as a measure of function which could be used to evaluate chronically ill and aging populations. The index ranks individuals according to adequacy of their performance. This adequacy is expressed as a grade (A, B, C, D, E, F, G, or Other) which summarises overall performance in bathing, dressing, toileting, transferring, continence and feeding (Table 6.1). This scale has been extensively used in studies of the natural history of disease and results of treatments.

**Table 6.1 Index of Independence in Activities of Daily Living**

These specific definitions for ranking appear below the index	
<b>A</b>	Independent in feeding, continence, transferring, going to toilet, dressing, and bathing
<b>B</b>	Independent in all but one of these functions
<b>C</b>	Independent in all but bathing and one additional function
<b>D</b>	Independent in all but bathing, dressing, and one additional function
<b>E</b>	Independent in all but bathing, dressing, going to toilet, and one additional function
<b>F</b>	Independent in all but bathing, dressing, going to toilet, transferring and one additional function
<b>G</b>	Dependent in all six functions
<b>Other</b>	Dependent in at least two functions, but not classifiable as C,D,E, or F

**Source:** Katz et al. 1963 page: 95

Over the years, a number of instruments measured ADL, although a few of them considered those with musculoskeletal conditions. Of those, the most commonly used instrument in arthritis is the Stanford Health Assessment Questionnaire (HAQ) (Bruce and Fries 2003; Fries et al., 1980). The HAQ was developed as a comprehensive measure of outcome in patients with a wide variety of rheumatic diseases, including osteoarthritis. The domain of disability is assessed in the HAQ by eight categories of dressing, arising, eating, walking, hygiene, reach, grip, and other common activities. There are four possible response options to rate difficulty in functioning, which range from 0-3 (without any difficulty (0), with some difficulty (1), with much difficulty (2), and unable to do (3)). The scoring system used in the HAQ reflects the disease activity, which is expressed as the degree of difficulty in performing ADLs. Each of the disability items has a corresponding aids / devices variable that is used to identify what type(s) of assistance, if any, is used to help with these activities. Again, the scoring system used for this range from 0 (No assistance is needed) to 3 (The patient usually needs BOTH a special device and

help from another person). However, the inclusion of environmental facilitators in HAQ increases the disability scores, as it assumes the more help is needed/ used by individuals the more dependent they are.

Latterly, Kempen and Suurmeijer developed the Groningen Activity Restriction Scale (GARS) to measure both ADL and IADL disability in community based studies with respect to the aid and services provided by professional home help and district nursing agencies (Kempen and Suurmeijer 1990). The GARS included 18 items (e.g. for ADL: dressing, getting in and out the bed, washing, toileting, feeding, using stairs, walking outdoors, take care of feet and toenails; for IADL: prepare breakfast / lunch / diner, do light / heavy household activities, wash and iron, make the beds and do the shopping) and five response options for each item to gauge what responders are able to do and not do independently. The response options ranged from: (1) yes, I can do it fully independently without any difficulty; (2) with some difficulty; (3) with great difficulty; (4) no, I cannot do it fully independently, I can only do it with someone's help; (5) No, I cannot do it at all, and I need complete help (Suurmeijer et al., 1994). One disadvantage of the GARS is that it is not taking the use of aids and adaptations into account which may promote independent function in older people, and similar to the HAQ, the disability score of the individuals increase with the increased use of environmental facilitators to function.

Another ADL measure developed originally in Sweden specifically for the use in Occupational Therapy (OT) is the Evaluation of the Daily Activity Questionnaire

(EDAQ) (Nordenskiöld et al., 1996). This measure encompasses 14 domains: where the level of difficulty is measured in personal and IADL (6-14 activities in each: 138 activities in total). Each of these domains has two sections: section (A) is scored without aids, alternate methods or help, and section (B) is scored with aids or alternate methods. As opposed to HAQ and GARS, the scoring system in EDAQ reflects the use of aids and assistance as an improvement in function. However, although this instrument's strength lies in the large number of domains it covers to provide a thorough OT assessment, it has been criticised for the potential responder burden for being too comprehensive (Archenholtz and Dellhag, 2008).

Previous studies of the onset of ADL disability have suggested that the disability process follows a complex and sometimes transient course (Stuck et al, 1999; Norburn et al., 1995). Reynolds and Silverstein (2003) investigated the role of various factors in the onset of ADL and IADL to examine the onset of disability both in the aggregate, and as an individual phenomenon. They did so in order to develop a long term strategy for better targeted home and community based services. In this study, participants were considered as 'dependent' or 'disabled' if they could not perform such activities without help. This study confirmed that the onset of ADL and IADL disability is a complex process, predicted by a variety of different factors. Amongst factors associated with the onset of ADL disability, the strongest predictors were identified as increasing age, being female and having arthritis.

In a series of studies involving a cohort of 754 community-dwelling adults aged 70 years and over who provided monthly reports of disability by telephone interview, Gill and colleagues revealed the high incidence of episodes of disability, their recurrent nature, and the effects of persistent episodes of disability. Over a median follow-up of 51 months, they found that over half the cohort experienced an episode of disability (dependence or difficulty in bathing, dressing, walking, or transferring) (Hardy and Gill, 2004). Of those, 81% recovered (i.e. regained independence in all 4 ADLs measured) within a year of their initial disability episode and a majority of these maintained independence for at least 6 months. This study concluded that disability is a recurrent rather than enduring condition and suggested interventions to maintain independence after recovery are needed (Hardy and Gill, 2004).

In a further analysis within the same cohort, but focused on bathing disability (defined as the 'inability to wash and dry one's whole body without personal assistance'), the authors reported that over the course of 6 years, 58.4% had at least one episode of bathing disability, and 34.0% had multiple episodes (Gill et al., 2006: a). The incidence of bathing disability was higher in women, in those who were physically frail, and in those aged 80 years and over. The fact that many of these incident episodes of bathing disability occurred in the absence of disability in other ADL domains (e.g. dressing, transferring, walking) suggested that bathing disability may be a 'sentinel event'. This study concluded that given the recurrent nature of bathing disability, the focus for intervention must not be solely on the prevention but also restoration and maintenance of independent bathing in older

people who become disabled (Gill et al., 2006:a). In a subsequent study of bathing disability and the risk of long-term admission to a nursing home, participants were followed up with monthly telephone interviews for a median of 75 months. The study determined the occurrence of persistent disability in bathing (defined as present for at least 2 consecutive months) and the time to the first long-term nursing home admission (defined as longer than 3 months). The authors found that the occurrence of persistent disability in bathing was independently associated with the risk of a long-term nursing home admission (Gill et al., 2006: b).

Balzi and colleagues (2010) measured factors potentially associated with high risk of ADL disability in a population of older adults ( $\geq 65$ ) in a prospective cohort study at baseline (n=1155), and at 3 years (n=897) follow-up studies. The 'incidence' of disability was defined as a new ADL disability at 3 years follow-up in those participants without ADL disability at baseline. Worsening disability was defined as increased ADL disability in those who already had ADL disability at baseline. Of those participants who were free of ADL disability at baseline, 8.4% developed ADL disability, and of those who were already disabled, 51.0% had a worsening ADL disability over a 3 years follow-up.

From this small and limited selection of studies, it can be assumed that, like estimates of prevalence, estimates of the rate of onset and persistence of self-care disability are likely to vary depending on the definition used and the population studied. Direct comparisons between the current study and the existing literature

will therefore be difficult. In addition, the studies by Gill and colleagues illustrate that the period of follow-up and the frequency of repeated measurement can yield very different estimates of the episode incidence of self-care disability and that this will differ from estimates of cumulative incidence such as those provided by Balzi and colleagues. The 3 years period of follow-up from Balzi and colleagues is the same as that used in the current thesis and, although the measure of self-care disability in this thesis is different, we might nevertheless hypothesise from these previous studies that: (a) the state of self-care restriction is reversible and therefore a proportion of those reporting self-care restriction at baseline will report no such restriction at 3 years follow-up ('back-transitions'), (b) the rate of onset of self-care restriction will be lower than the rate of persistence, (c) the rate of onset will be higher in women and in older age groups. It must also be assumed that the cumulative incidence of self-care restriction at 3 years will under-estimate the episode incidence of self-care restriction.

## **6.2 Aims and objectives**

The overall aim of this study was to describe the longitudinal course of self-care restriction in older people in the general population over a three year period. Using baseline and 3 years data from the NorStOP in community-dwelling adults aged 50 years and over the objectives of these analyses were:

- i. To estimate the risk of onset and persistence of self-care restriction at 3 years



- ii. To describe the relationship between the risk of onset and risk of persistence of self-care restriction at 3 years and baseline demographic, socio-economic and health characteristics
- iii. To explore the direction and strength of association between the risk of onset and risk of persistence of self-care restriction at 3 years and the site, number, and pattern of peripheral joint pains at baseline

## **6.3 Methods**

### **6.3.1 Dataset**

The analyses described in this chapter use data collected at baseline and 3 years in the NorStOP study. Participants were adults aged 50 years and over in the general population who responded and completed the Keele Assessment of Participation (KAP) self-care item at baseline and 3 years follow-up. Of those 7725 responders who completed KAP self-care item at baseline, 5425 consented to further contact, 4089 (75%) of whom were successfully followed up at 3 years.

### **6.3.2 Outcome**

Longitudinal studies involve following participants over time, measuring an outcome variable and risk factors, at least at two different points in time (Jewell and Hubbard, 2006). Incidence is defined as number of new cases in a specified population in a given period and may be measured as a frequency count, a rate, or a proportion (Porta, 2008). The incidence rate is defined as the rate at which new events occur in a population, and the cumulative incidence is defined as the

proportion of people presenting with the onset of a health-related event during a specified time interval (Porta, 2008). In this study participants were followed at two time points with a 3 years interval. Therefore, it was not possible to measure the incidence of episodes of self-care restriction, as it was not possible to observe the timing of the event or change in state. Therefore, instead of incidence, the term 'onset' was used in this study to refer to *the risk of self-care restriction at 3 years in people free of self-care restriction at baseline*. 'Persistence' was defined as *the risk of self-care restriction at 3 years in people reporting self-care restriction at baseline*. These two variables – onset and persistence of self-care restriction at 3 years – were used as outcome variables in multivariable analyses investigating baseline determinants.

### **6.3.3 Covariates**

#### *Demographic, socio-economic and health characteristics*

Longitudinal studies comprise of studying the same group of individuals over an extended period of time and are an efficient way to discover associations between factors that are not visibly related. Examining the demographic, socio-economic and health characteristics of the study cohort can highlight these associations with the outcome and provide further information on risk factors.

Demographic and socio-economic variables used in this analysis were selected for their associations with restricted self-care at baseline. Age and gender were used to estimate the distribution of the onset and persistence of restricted self-care in the general population of adults aged 50 years and over. In order to describe the

relationship between the risk of onset and risk of persistence of self-care restriction, and the baseline socio-economic and health characteristics of those responders at 3 years, participants were described using perceived adequacy of income (find it strain to get by from week to week / have to be careful with money / able to manage without much difficulty / quite comfortably off), educational attainment (whether they had gained qualifications through study as an adult) along with the health characteristics using 12-item Health Survey (SF-12) physical and mental health component scores. A linear trend in self-care restriction with increasing age was observed at baseline and self-care restriction was higher in women. Individuals' perceived adequacy of income and educational attainment were found to be independently associated with self-care restriction in the multivariable analysis where other socio-economic factors such as occupational class were eliminated from the parsimonious model at baseline. These variables were described in detail in Chapters 4 and 5.

### *Joint pain*

Peripheral joint pain was included in this study to explore the direction and strength of associations between the risk of onset and risk of persistence of self-care restriction at 3 years. Peripheral joint pain was defined as pain in the past year in hand, hip, knee, and / or foot. This was analysed in three ways: by site (each site – yes/no), as a simple count of the number of peripheral joint areas reported as painful (0-4) and by pattern (no pain, hand only, single lower limb, multiple lower limb, hand+ lower limb).

#### 6.3.4 Statistical Analysis

The primary analyses were conducted on cases with complete data at baseline and 3 years. Estimates of the onset and persistence of self-care restriction at 3 years were expressed as percentages, overall, and stratified by age and gender. Differences in onset and persistence between males and females were tested using *chi-squared* test. Differences between age groups (50-59, 60-69, 70-79, 80+ years) were tested using *chi-squared* test for trend.

##### *Relationship with baseline socio-economic and health characteristics*

Descriptive characteristics such as socio-economic (adequacy of income and educational attainment) and health characteristics (SF-12 physical and mental health scores) of the study participants were cross-tabulated using frequencies and percentages to compare those with the onset of self-care restriction against those who remained free of restriction. Differences between the groups were tested using *chi-squared* for non-parametric variables (e.g. educational attainment and perceived adequacy of income) and the *independent samples t-test* for parametric variables (e.g. PCS and MCS scores). These were expressed as p-values. The analyses were repeated comparing those with persistent self-care restriction to those who recovered from self-care restriction at 3 years.

##### *Relationship with selected peripheral joint pain at baseline*

Those who reported onset and persistence of restricted self-care at 3 years were cross-tabulated with joint pain by site, pattern and number at baseline. Links between the onset and persistence of restricted self-care at 3 years and joint pain

by site, pattern and number at baseline were investigated using univariable and multivariable logistic regression analysis (adjusting the associations for all joint specific characteristics) as this method was highlighted as the most appropriate method in the previous chapter (Chapter 5). Associations were presented as odds ratios with 95% confidence intervals.

### *Sensitivity analysis*

Missing data is common in longitudinal analyses due to various factors such as missing items on the questionnaire, non-response and study attrition. There is increasing recognition that studies which restrict analysis to participants with complete data can be biased and improvident (Spratt et al., 2010). Multiple imputation (MI) is increasingly endorsed in epidemiology to adjust for the bias and data loss that may occur in analyses limited to study participants with complete data (Spratt et al., 2010). MI is a technique that allows for uncertainty about the missing data by creating several copies of the dataset in which missing values are substituted by imputed values sampled from an earlier predictive distribution that is estimated from the observed data (Rubin, 1996; Schafer, 1997).

As described in chapter 4, the extent and the characteristics of the study attrition at NorStOP 3 years sample could introduce bias to the longitudinal analyses results. Therefore MI was utilised in this study in an attempt to understand and control this possible bias by examining whether the estimates based on partial and complete datasets had differed.

Missing data mechanisms are categorised into three categories: (i) missing completely at random (MCAR); here the assumption is that the attrition is independent of both observed and unobserved data, (ii) missing at random (MAR); attrition depends on the observed data, but not on unobserved data, and (iii) missing not at random (MNAR); attrition depends on unobserved data (Twisk and Vente, 2002). It not possible to distinguish between MAR and MNAR from the observed data alone, but the MAR assumption can be made more plausible by including more explanatory variables in the analysis (White et al., 2010). The missing data mechanism is important, as different types of missing data require different types of analysis. Under MAR assumption, a set of covariates  $X$  is observed and the missing values,  $Y_{mis}$ , depend on the observed variables  $X$ . There is no statistical test to prove this assumption, however a common approach to see if MAR assumption is plausible, is to conduct univariable and multivariable logistic regression to determine if covariates  $X$  were associated with  $Y_{mis}$  (outcome variable) (Vargas-Chanes et al., 2003). In the preliminary analysis conducted, baseline values of the key factors associated with restricted self-care were predictive of the missing outcome data at 3 years, suggesting covariate dependent missingness. The assumption regarding the lack of association with unobserved quantities in this dataset meant the data was MAR, since the missingness can actually be predicted (but is random after controlling for missingness due to observed quantities) (Horton and Kleinman, 2007). Under the MAR assumption, identifying the variables for the imputation process is an important step and the number of variables included in the imputation process affects the quality of the sensitivity analysis. This is because MI's ability to reduce bias in a given analysis

depends on the measured variables that are associated with both the manner in which data are missing and the outcome variable (Engels and Diehr, 2003). Therefore, only six variables from the parsimonious model developed at baseline were selected to be included in the imputation process (age, activity limitation (PF10), depression (HADs), cognitive complaint (SIP), perceived adequacy of income and educational attainment).

Two major approaches to MI were identified by the author to create multiply imputed datasets. The first approach identified was the approach taken by Stata's *mi imputes* command. This assumes the joint distribution of all variables in the imputation model to be multivariate normal. The second approach identified was based on each conditional density of a variable given other variables, a user-written *ICE* (imputation by chained equations approach) programme (Royston, 2005; 2007; 2009). As the pattern of missingness was deemed to be arbitrary in this dataset, an *ICE* programme was used for imputation. An advantage of the *ICE* approach is that variables are not assumed to have a multivariate normal distribution (Von Hippel, 2007). *ICE* is used to perform multivariable imputation via chained equations, and the *mim* command analyses multiply imputed data by performing data analysis and combining these individual analyses using Rubin's combination rules (Rubin, 1987). Recommendations for the number of imputations vary in the literature, from 5 to 50 depending on the amount of missing data (Graham et al., 2007). For the sensitivity analysis different m sets of imputations were carried out (5; 20; 30; 40) to see if different results were acquired. As results

were similar and the power of the analysis decreased with the number of imputations, 20 m sets was decided as the optimum amount of imputation.

MI analysis using *ice* consisted of three phases: (i) imputation - creating multiply imputed data, (ii) analysis of imputed datasets, and (iii) combining of estimates from imputed datasets using Rubin's combination rules.

*i) Imputation:*

The original baseline general population sample to draw imputations from was identified after 365 people were omitted from the dataset according to their 3 year exclusion status (308 were excluded prior to mailing, 33 died, 6 changed address, 6 ineligible, 12 ill). Those who were not restricted in self-care at baseline formed the sample to impute for the *onset of self-care restriction* at 3 years (n=6571). Those who were restricted in self-care at baseline formed the sample to impute for the *persistence of self-care restriction* at 3 years (n=796). The unknown missing data were replaced by 20 independent simulated sets of values drawn from the predictive distribution of the missing outcome data, conditional on the observed parsimonious variables data to create multiply imputed datasets at 3 years using *ICE*.

*ii) Analysis of imputed datasets:*

Following the completion of imputations, each imputed dataset for the onset and persistence of self-care restriction at 3 years were analysed separately. The regression coefficients were estimated from each imputed dataset with 95%



confidence intervals. The results of these  $m$  imputations were examined to ensure that the distributions looked reasonable after imputation, and that they were not out of range of the original values.

*iii) Combining of estimates from imputed datasets:*

In the last step the multiply imputed estimates were combined into an overall estimate with 95% confidence interval using Rubin's rule of combination which allows the variance of estimates between different imputed datasets to be taken into account. The combined variance incorporates both within-imputation variability (uncertainty about the results from one imputed dataset) and between imputation variability (reflecting the uncertainty due to the missing information) (White et al., 2011).


## 6.4 Results

### 6.4.1 Onset self-care restriction at 3 years: complete-case analysis

Based on complete case analysis, there were 4089 potentially eligible respondents at 3 year follow-up, of whom 3770 were free of self-care restriction at baseline (Figure 6.1).

250 (6.6%; 95 % CI 5.9, 7.5) reported the onset of self-care restriction at 3 years (Fig: 6.1). There was a linear trend with increasing age ( $p \leq 0.001$ ) but no difference between men and women (Table 6.2).

**Figure 6.1** Flowchart of the onset and persistence of self-care restriction in the general population over a 3 years period



**Table 6.2 Onset of restricted self-care at 3 years; overall and stratified by age and gender**

		<b>N</b>	<b>Onset n (%)</b>	<b>P-value</b>
<b>Overall</b>		3770	250 (6.6)	
<b>Gender</b>				
	Female	2070	141 (6.8)	p=0.336
	Male	1700	109 (6.4)	
<b>Age group</b>				
	50-59	1452	61 (4.2)	p≤0.001
	60-69	1272	79 (6.2)	
	70-79	828	74 (8.9)	
	80+	218	36 (16.5)	

#### 6.4.2 Persistence of self-care restriction at 3 years

Of 319 eligible respondents who reported self-care restriction at baseline and were successfully followed up at 3 years, 124 (38.9%; 95% CI 33.7, 44.3) reported self-care restriction again at 3 years follow-up (Fig: 6.1). Persistent self-care restrictions were higher in men (Table 6.3). The risk of persistent self-care restriction at 3 years was unexpectedly low in 80+ year olds, resulting in no overall trend with age.

**Table 6.3 Persistence of restricted self-care at 3 years; overall and stratified by age and gender**

		<b>N</b>	<b>Persistence n (%)</b>	<b>P-value</b>
<b>Overall</b>		319	124 (38.9)	
<b>Gender</b>				
	Female	202	70 (34.7)	p=0.042
	Male	117	54 (46.2)	
<b>Age group</b>				
	50-59	86	33 (38.4)	p=0.536
	60-69	103	42 (40.8)	
	70-79	83	35 (42.2)	
	80+	47	14 (29.8)	

### 6.4.3 Association with baseline socio-economic and health characteristics

#### *Onset of self-care restriction*

The crude risk of onset of self-care restriction at 3 years was associated with lower socioeconomic status - as measured by educational attainment and perceived financial strain - at baseline. For example, the risk of onset was four times higher among individuals who felt that they were under financial strain, compared to those who rated themselves as 'comfortably off' (Table 6.4). Those who experienced the onset of self-care restriction at 3 years tended to have lower (unadjusted) physical and mental health scores at baseline than those who remained free of self-care restriction at follow-up (Mean (SD) SF-Physical Component Score (0-100): 32 (11) vs 42 (12),  $p<0.001$ ; Mean (SD) SF-Mental Component Score (0-100): 44 (12) vs 50 (11),  $p<0.001$ ).

**Table 6.4 Onset of self-care restriction at 3 years follow-up, by socio-economic status at baseline**

		<b>N</b>	<b>Onset n (%)</b>	<b>P-value</b>
<b>Overall</b>		3770	250(6.6)	
<b>Adequacy of Income</b>				
	Comfortable	445	17 (3.8)	
	Little difficulty	1183	76 (6.4)	
	Be careful	1101	105 (9.5)	
	Strain	91	11 (12.1)	$p\leq 0.001$
<b>Higher educational attainment</b>	<b>No</b>	2318	182 (7.9)	
	<b>Yes</b>	1395	60 (4.3)	$p\leq 0.001$

### *Persistence of self-care restriction*

Similar to the onset analysis, the crude risk of persistence of self-care restriction at 3 years was associated with the perceived financial strain at baseline. The risk of persistence was 2.5 times higher among individuals who stated financial strain than those who described themselves as comfortable, but there was no relationship with educational attainment (Table 6.5). Those who had persistent self-care restriction at 3 years had lower (unadjusted) physical health scores at baseline than those who remained free of self-care restriction at follow-up, but there was no significant difference in mental health scores between those recovered and those had persistent self-care restriction at 3 years (Mean (SD) SF-Physical Component Score (0-100): 27 (10) vs 29 (9),  $p=0.040$ ; Mean (SD) SF-Mental Component Score (0-100): 38 (11) vs 41 (12),  $p=0.105$ ).

**Table 6.5      Distribution of the persistence of self-care restriction at 3 years by socio-demographic and health characteristics**

		<b>N</b>	<b>Persistence n (%)</b>	<b>P-value</b>
<b>Overall</b>		319	124 (38.9)	
<b>Adequacy of Income</b>				
	Comfortable	16	4 (25.0)	
	Little difficulty	69	23 (33.3)	
	Be careful	168	65 (38.7)	
	Strain	26	16 (61.5)	$p=0.050$
<b>Higher educational attainment</b>				
	No	251	97 (38.6)	
	Yes	56	20 (35.7)	$p=0.402$

#### **6.4.4 Association of the onset of self-care restriction with joint pain at baseline**

Individuals with pain at each of the selected peripheral joint areas were at higher risk of onset of self-care restriction at 3 years compared to those without that pain (Hand pain= Crude OR 1.7 95% CI 1.3, 2.3; Hip pain= OR 2.1 CI 1.6, 2.6; Knee pain= OR 1.9 CI: 1.4, 2.4; Foot pain= OR 2.0; CI 1.5, 2.6) (Table 6.6).

The number of painful peripheral joint areas at baseline was associated with the onset of self-care restriction within each category (Table 6.7). For example, the odds of self-care restriction at 3 years was 3.3 times higher in individuals reporting pain in the hand, hip, knee and feet at baseline compared with people reporting pain in none of these areas (Four painful joint areas = OR 3.3; CI 2.2, 4.9).

Pattern of joint pain at baseline was also associated with the onset of self-care restriction at 3 years in each category level but only for those with patterns involving multiple joint areas (Table 6.8). Neither isolated hand pain nor single-site lower limb pain were associated with an increased risk of self-care restriction at 3 years (Hand pain only= OR 1.1; CI 0.6, 2.2; Single lower limb pain= OR 1.0; CI 0.6, 1.6).

**Table 6.6 Onset of self-care restriction at 3 years and site of joint pain at baseline**

3 years response to KAP self-care item of those who were not restricted at baseline		Onset of self-care restriction at 3 years		
n= 3770*				
		No n	Yes n	OR (95%CI)
Hand pain				
	No	1938	103	1
	Yes	1513	141	1.7 (1.3, 2.3)
Hip pain				
	No	2349	123	1
	Yes	1111	120	2.1 (1.6, 2.6)
Knee pain				
	No	1723	85	1
	Yes	1739	160	1.9 (1.4, 2.4)
Foot pain				
	No	2218	115	1
	Yes	1243	129	2.0 (1.5, 2.6)

\* Cells do not add to 3770 due to missing data

**Table 6.7 Onset of self-care restriction at 3 years and number of painful joint areas at baseline**

3 years response to KAP self-care item of those who were not restricted at baseline		Onset of self-care restriction at 3 years		
n= 3770				
		No	Yes	OR (95%CI)
		n	n	
Number of painful joint areas				
	None	832	32	1
	One	867	44	1.0 (0.7, 1.6)
	Two	791	49	1.3 (0.8, 1.9)
	Three	556	56	2.0 (1.4, 3.0)
	Four	329	53	3.3 (2.2, 4.9)

\* Cells do not add to 3770 due to missing data

**Table 6.8 Onset of self-care restriction at 3 years and pattern of joint pain at baseline**

3 years response to KAP3 of those who were not restricted at baseline		Onset of self-care restriction at 3 years		
n= 3770				
		No	Yes	OR (95%CI)
		n	n	
Pattern of joint pain				
No pain	977	48	1	
Hand pain only	199	11	1.1 (0.6, 2.2)	
Single lower limb	668	33	1.0 (0.6, 1.6)	
Multiple lower limb	409	36	1.8 (1.1, 2.8)	
Hand and lower limb	1267	122	2.0 (1.4, 2.8)	

\* Cells do not add to 3770 due to missing data

#### **6.4.5 Association of the persistence of self-care restriction with joint pain at baseline**

Individuals with hip, knee, or foot pain at baseline appeared to have a slightly higher risk of persistent self-care at 3 years, although this was statistically non-significant (Hip= Crude OR 1.5 95% CI 0.9, 2.5; Foot= OR 1.4; CI 0.9, 2.3) (Table 6.9).

People with the four painful peripheral joint areas at baseline had higher odds of reporting self-care restriction at 3 years compared with people reporting no painful joint areas. However the relationship was statistically non-significant (Table 6.10).



The pattern of joint pain at baseline was associated with persistence of self-care restriction at 3 years, but only for those with patterns involving multiple joint areas. However, again these associations were not statistically significant due to small sample size (Table 6.11).

**Table 6.9 Persistence of self-care restriction at 3 years and site of joint pain at baseline**

<b>3 years response to KAP3 of those who were restricted at baseline</b>		<b>Persistence of self-care restriction at 3 years</b>		
<b>n= 319</b>				
		<b>No</b>	<b>Yes</b>	<b>OR (95%CI)</b>
		<b>n</b>	<b>n</b>	
<b>Hand pain</b>				
	No	64	41	1
	Yes	124	80	1.0 (0.6, 1.6)
<b>Hip pain</b>				
	No	87	42	1
	Yes	101	76	1.5 (0.9, 2.5)
<b>Knee pain</b>				
	No	51	26	1
	Yes	138	94	1.3 (0.8, 2.3)
<b>Foot pain</b>				
	No	74	37	1
	Yes	116	84	1.4 (0.9, 2.3)

\* Cells do not add to 3770 due to missing data

**Table 6.10 Persistence of self-care restriction at 3 years and number of painful joint areas at baseline**

3 years response to KAP3 of those who were restricted at baseline		Persistence of self-care restriction at 3 years		
n= 319				
		No	Yes	OR (95%CI)
		n	n	
<b>Number of painful joint areas</b>				
	None	16	7	1
	One	27	15	1.2 (0.4, 3.8)
	Two	33	22	1.5 (0.5, 4.3)
	Three	55	28	1.2 (0.4, 3.1)
	Four	50	45	2.0 (0.8, 5.4)

\* Cells do not add to 3770 due to missing data

**Table 6.11 Persistence of self-care restriction at 3 years and pattern of joint pain at baseline**

3 years response to KAP3 of those who were restricted at baseline		Persistence of self-care restriction at 3 years		
n= 319				
		No	Yes	OR (95%CI)
		n	n	
<b>Pattern of joint pain</b>				
	No pain	30	14	1
	Hand pain only	6	3	1.1 (0.2, 4.9)
	Single lower limb	21	12	1.2 (0.5, 3.2)
	Multiple lower limb	26	22	1.8 (0.8, 4.2)
	Hand and lower limb	112	73	1.4 (0.7, 2.8)

\* Cells do not add to 3770 due to missing data

#### **6.4.6 Sensitivity analysis**

Analyses of each imputed set demonstrated that the distribution of data was reasonable after imputation and they were not out of range of the original values. When 20 sets of imputations were combined using Rubin's rules, the estimated frequency of onset of restricted self-care at 3 years was 7.9% (95% CI 7.1; 8.6) (compared to the observed estimate of 6.6%). The estimated frequency of persistence of self-care restriction at 3 years was 39.8% (CI 35.6; 44.0) (compared to the observed estimate of 38.9%).

#### **6.4.7 Summary of results**

In this study, the incidence of self-care restriction at 3 years in people aged 50 years and over was estimated to be 6.6%. In those already reporting restriction at baseline, 38.9% reported self-care restriction again 3 years later, implying persistent restriction. The net effect of these changes on the prevalence of self-care restriction over 3 years within this cohort was a small increase (7.8% at baseline to 10.9% at 3 years follow-up). An increase in the prevalence of self-care restriction over time is consistent with the observed cross-sectional association between self-care restriction and increasing age reported in Chapter 5. However, selective loss to follow-up appears likely in the current study and both onset and persistence are likely to be under-estimated in the complete-case analysis.

Individuals who were older, of lower socioeconomic status, and who were reporting worse physical and mental health at baseline were at increased risk of onset of self-care restriction at follow-up. The presence of peripheral joint pain at

baseline, particularly multiple joint pains, was also associated with an increased risk of onset of self-care restriction, but there were no significant associations with persistence. The risk of persistent self-care restriction at 3 years appeared to be higher among men, those with poorer physical health, and those under financial strain. But persistence was not related to age, educational attainment, or mental health. Lower numbers were available for the analysis of persistence than the analysis of onset, limiting the precision of the former estimates.

## **6.5 Discussion**

### *Critical comparison against previous studies*

In the study where Balzi and colleagues measured factors potentially associated with high risk of ADL disability at baseline and 3 years follow-up, the ‘incidence’ of disability was defined as a new ADL disability at 3 years follow-up in those participants without ADL disability at baseline. This is consistent with the risk of onset of self-care restriction at 3 years in the study undertaken in this chapter. Their estimated onset of ADL disability was 8.4% (Balzi and colleagues 2010), somewhat similar to the estimated onset of self-care restriction in this study (6.6%). However, it is important to bear in mind that the outcome of interest in Balzi and colleague’s study was ADL disability (reported need for help of another person in performing at least one ADL), and was not person-perceived participation in self-care (self-care needs met as and when wanted with or without help and assistance) as measured in the NorStOP study.

Balzi and colleagues (2010) also reported worsening disability (defined as increased ADL disability in those who already had ADL disability at baseline) in their study of *risk factors for disability in older persons over 3 years follow-up*, which can be compared to the estimation of persistent self-care restriction (defined as continued existence or occurrence of restricted self-care in those reporting self-care restrictions at baseline) in the NorStOP study reported in this chapter. Their estimation of 'worsening disability' was 51.0% over a 3 years follow-up, compared to the NorStOP study estimation of persistent self-care restriction of 38.9%. The persistence of self-care restriction was measured using KAP, an aggregated measure of self-care, which was dichotomised. This dichotomy did not allow accounting for the worsening of the scores over a 3 years follow-up. Also, fewer numbers were available for the analysis of persistence in the follow-up, limiting the precision of the estimates.

Another prospective cohort study investigated ADL function (bathing, dressing, walking and transferring) monthly in a sample of community-dwelling adults aged 70 and older, over a five year period. It reported that 81% recovered within a year and maintained independence for at least 6 months (Hardy and Gill, 2004). The recovery rate from self-care restriction at 3 years in the NorStOP population was 61.1%. However, participants in this study were assessed monthly as opposed to the 3 years interval in the NorStOP study, and were in the study for longer (5 years of 3 years). The NorStOP study may have missed the resolved cases of persistent self-care restriction in this long interval, as this study stated that recovery from disability in essential ADLs among community-dwelling older adults is common,

albeit often brief in duration (Hardy and Gill, 2004). Gill and Hardy's study also took a different approach than the study described in this chapter. They sought to determine the rate of and time to recovery of independent function (defined as not being dependent on carrying out ADL tasks) in community-dwelling older persons who become newly disabled (dependent) in ADLs (bathing, dressing, walking, and transferring). Their measure of ADL included walking as well as self-care tasks, and their data collection methods involved comprehensive home assessments at baseline, 18, and 36 months and monthly telephone interviews for up to 53 months. Comprehensive assessments were completed by trained research nurses using standard instruments. During monthly interviews participants were assessed for each ADL by asking "at the present time, do you need help from another person to perform the task?" Participants who needed help with or were unable to complete one or more of the ADL tasks were considered disabled. Dissimilar to the NorStOP study, participants were not asked about eating, toileting, or grooming on the basis that disability in these ADLs is uncommon without concurrent disability on bathing, dressing, walking, or transferring (Gill et al., 1995; Gill et al., 1995a). However, considering that the methods and approach of this study differ to the NorStOP study, the rates of recovery reported may conceptually explain what is deemed to be an over-optimistic recovery from self-care restriction in this study, as this process is underlined to be transitory.

Other studies that have investigated the level and time course of ADL disability have acknowledged that disability can both increase and decrease over time, and that recovery is quite often followed by decline due to non-linear process of

disability (Nusselder et al., 2006). Therefore recovery or persistence based on two measurements may not reflect permanent change. Nusselder, Caspar and Mackenbach identified 9 trajectories of disability ranging from entirely non-disabled (1) to severely disabled (8 and 9). A person reporting only difficulty in carrying an object would just fit into 'entirely non-disabled' group. Had this person been unable to carry out a task, or had other difficulties, they would then fit into one of the trajectories with disability. Severely disabled persons would be those with at least some difficulty in all disability items (including ADLs), or who had great difficulty, or who was unable to do several items. This study also found significant association between the trajectories of disability and age; those who were younger than 65 years of age were associated with lower risk of almost all trajectories of disability and death.

In terms of the critical comparison with the previous studies which investigated the socio-demographic and health differences and the risk of self-care restriction, Covinsky and colleagues' study (2008) offers interesting comparisons. This prospective longitudinal study examined whether symptomatic arthritis in middle age predicts the earlier onset of difficulties with ADLs (Covinsky et al., 2008). Symptomatic arthritis was measured at baseline according to self-report, by asking participants "Have you ever had, or has a doctor ever told you that you have, arthritis or rheumatism?" The outcome measure was persistent difficulty with ADL tasks and these tasks were bathing, dressing, transferring from a bed to a chair or out of chair, using a toilet, or eating. Those with arthritis were more likely to develop persistent difficulty in ADL function over a 10 year follow-up. Those with

both arthritis and another chronic condition (e.g. depression) were at markedly greater risk for developing difficulties with ADL. Again, the outcome of interest in this study was 'difficulty' in selected self-care activities, which mostly consisted of self-care tasks (other than for transferring) as opposed to 'restriction' in aggregated measure of self-care. The definition of joint pain was also different, as it was restricted to a form of arthritis being presented. Nevertheless, both results suggest that middle aged adults with joint pain are at a greater risk of reporting difficulties and/or restrictions in self-care activities as they age, especially when these adults have a declining mental health.

### *Limitations*

#### *i. Selective loss to follow-up*

Loss to follow-up introduces imprecision and the potential for bias (Hennekens and Buring, 1987). Although the sample size for complete case analysis in the this study was relatively large ( $n=3147$ ), estimates for the onset and persistence of self-care restriction at 3 years were based on 250 and 124 cases respectively, causing imprecision of stratified estimates and odds ratios in multivariable analyses.

Selective loss to follow-up was addressed in this study by sensitivity analysis using multiple imputation. The estimates based on multiple imputation were slightly higher than those from the complete case analysis, suggesting that those lost at follow-up were at higher risk of onset and persistence of self-care restriction. The MI approach was chosen as a suitable method following a set of preliminary



analyses on missing data. This involved the usual exploratory data techniques, such as examining means and standard deviations, and graphing distributions. Also, it was important to understand not just how much data was missing, but the patterns of the missing values. Patterns of missing data were explored using the baseline values. It is important to understand the pattern of missing values as the patterns of missingness may suggest why these values are missing. If a variable that has more missing values than others, it is important to consider why this may be the case. The pattern of missingness influences the type of imputation that can be used. Factors identified in the parsimonious cross-sectional model at baseline (see Chapter 5) were included as covariates in the MI as they were identified as independently associated with self-care restriction at baseline, and were strongly correlated with the missingness in the preliminary analysis. It is suggested that the number of variables entered in the MI increases the accuracy of the estimates (Twisk and Vente, 2002). Inclusion of these variables which were highly correlated with the self-care restriction was made on the assumption that the missingness mechanism was MAR. But if this assumption was not completely true and the data was missing not at random (MNAR), in which the probability of missingness depends upon the value of variables that may themselves be missing (Sterne et al., 2009), then using imputation would still give biased estimates.

Although it is not possible to measure the adequacy of MI in correcting responder bias empirically, it is no longer acceptable to ignore missing values in longitudinal surveys as analysis of complete cases is deemed to be potentially misleading and inefficient. MI is increasingly used and it has been suggested that correctly and

carefully applied imputation methods should reduce bias and increase precision (Spratt et al., 2010). Nevertheless, the author acknowledges that application of MI is not simple, and the MI approach used in this study may not be less biased than the complete case analysis.

#### ii. Long interval – unobserved episodes of self-care restriction

Due to the long interval between baseline and the 3 years follow-up study, it was not possible to observe changes in the participants' self-care status (restricted yes / no) that occur and resolve in this time period. The effect is that the true incidence rate of episodes of self-care restriction is likely to be much higher than the onset at 6.6% at 3 years. The use of the dichotomous definition for self-care restriction meant that it was not possible to further quantify the amount of change in self-care restriction. For example, those with persistent self-care restriction may not be static but actually experience a worsening of the severity of restriction over time.

#### iii. Generalisability

The study population at baseline was representative of the general population in England in terms of age and gender. However, at 3 years follow-up those who were in the older age groups and female had higher rates of loss to follow-up, leaving a cohort that was younger with more males compared to baseline. This means that the estimates of the onset and persistence of self-care restriction is likely to be lower than the true rates in the local and national population, but this does not necessarily mean that the study results are not generalisable. The generalisability of the study findings also depends on factors that determine self-

care restriction (e.g. age, physical limitation, depression, lower socio-economic status), and whether the study population is representative of the local and national population for these characteristics. The sensitivity analysis resulted in a small increase in the estimated risk of onset and persistence and was in agreement with the findings of the wider literature on the onset and persistence of ADL disability and links with joint pain. Thus, study findings are compatible with previous research findings; albeit the estimated rates of the onset and persistence of self-care restriction in the local and national population is expected to be higher.

#### iv. Crude analyses

Analyses that investigated links between the onset and persistence of self-care restriction and pattern, site and number of peripheral joint areas at baseline were crude analyses and established that people with joint pain, in particular with multiple joint pains, have a higher risk of onset and persistence of self-care restriction over the following three years than people without joint pain. The next chapter will examine the association between the onset and persistence of self-care restriction and peripheral joint pain in more detail, considering specific joint characteristics and exploring other factors that modify this relationship.

## **6.6 Conclusions**

Self-care restriction, as measured by the KAP, appears to be an unstable state, with evidence of a substantial majority of people with self-care restriction at one point in time reporting a 'back-transition' 3 years later. Study results suggest that people with joint pain, particularly with multiple joint pain areas, and those with

lower socio-economic backgrounds may be at a higher risk of self-care restriction. To identify factors that predict the onset and persistence of self-care restriction in those with joint pain, further analyses within this group is required. The next chapter describes such analyses.

**Table 6.12     Summary of key findings**

- This chapter described the longitudinal course of self-care restriction in older people in the general population over a three year period, using baseline and three years follow-up data from the NorStOP.
- The onset and persistence of self-care restriction at three years was 6.6% (95% CI: 5.9, 7.5) and 38.9% (CI: 33.7, 44.3) respectively.
- Individuals who were older, of lower socio-economic status, and who were reporting worse physical and mental health at baseline were at increased risk of onset of self-care restriction at follow-up.
- Persistent self-care restriction at three years was higher among men, and those under financial strain.
- The presence of peripheral joint pain, particularly multiple joint pains, was also associated with an increased risk of onset of self-care restriction, but there were no significant associations with persistence.
- In order to identify factors that predict the onset and persistence of self-care restriction in those with joint pain, next chapter describes further analysis within this group.

## **Chapter 7**

# **Potential predictors of the risk of onset and persistence of restricted self-care in older people with joint pain**

### **7.1 Introduction**

The previous chapter (Chapter 6) estimated the risk of onset and persistence of self-care restriction at 3 years in older adults in the general population. The results have shown that older adults with peripheral joint pain, particularly multiple peripheral joint pain, were at a higher risk of reporting self-care restrictions. The next set of analyses focus on those with joint pain to ascertain the potential determinants of restricted self-care in this population.

Persistent pain is commonly experienced by older adults with osteoarthritis, and may result in mobility problems when the pain is severe (Wilkie et al. 2006; Arthritis Research Campaign 2002). The WHO (2003) suggest that eight out of ten people with osteoarthritis experience some kind of physical limitation, and that one in four cannot perform activities of daily living (ADL) due to pain in the small joints of hands and fingers hindering the conduct of these activities. Associations between joint pain/arthritis and future self-care disability were reported in previous studies of ADL (Covinsky et al., 2008; Song et al., 2006; Donald and Foy, 2004; Stuck 1999; Dunlop et al., 1998; Davis et al., 1991; Yelin and Katz, 1990). Earlier studies emphasised the negative influences of joint pain on participation, indicating that pain associated with osteoarthritis is an important predictor of future disability in activities of daily living (Davis et al., 1991). They also highlighted the importance of studying other factors such as psycho-social factors and mental health status to

understand the predictors of functional decrease in those with joint pain (Gignac et al., 2000; Yelin et al., 1987).

A study conducted by Dunlop et al (1998) assessed the longitudinal impact of joint impairment (calculated as presence/absence of tenderness, deformity and/or limitation of lower spine, hips, knees, ankles, feet and upper spine as well as wrist, elbow, shoulder, distal and proximal hand regions) on overall disability and domain-specific thresholds for ADL that are associated with use of long-term care. In a longitudinal study, using 484 adults aged 65 years and older, lower-extremity joint impairment was found as a strong risk factor for future disability associated with the use of long-term care in older adults. One study showed that one-fifth of the population aged over 50 had severe difficulty with activities of daily living due to knee pain. They found strong associations between difficulty in functioning and joint pain duration (Jinks et al., 2002). In another study, in which 4804 participants aged 75 years and over from a UK General Practice (GP) population were followed up for over a year in 1998, it was reported that some degree of joint pain (defined as those pains older people perceived as related to their joints) was reported by 83%, and joint pain was associated with a three-fold increased likelihood of dependency for activities of daily living (Donald and Foy, 2004).

The conceptual model adhered to in this thesis suggests that a range of comorbid health conditions, impairments, and personal and environmental factors may influence the risk of self-care restriction (WHO, 2001). Song and colleagues (2006) examined the population impact of arthritis on disability incidence among older

Americans aged 65 and over using the longitudinal data (1998-2000) from the Health and Retirement Study (HRS); a national probability sample of elderly Americans (n=7758). In this study, older adults who had baseline arthritis had a substantially higher incidence of ADL disability (inability to perform activities of daily living) compared with those without arthritis (9.3% cf 4.5%). Even after adjusting for all other risk factors (demographics, health factors, health behaviors, and medical access) arthritis remained as an independent and significant predictor for developing ADL disability (Song et al., 2006). A subsequent study conducted by Covinsky and colleagues (2008) using the same HRS cohort (n=7543) examined whether symptomatic arthritis (those reporting pain, stiffness and swelling in their joints, were currently taking medications or other treatments for their arthritis or rheumatism, or had seen a doctor for their arthritis or rheumatism in the previous 12 months) in middle age (defined as aged 50+) predicts the earlier onset of ADL difficulties (difficulty with bathing, dressing, transferring from a bed to a chair or out of a chair, using a toilet, or eating). This study also reported that participants with arthritis were more likely to report persistent difficulty with ADL function even after adjusting for other socio-demographic and health factors. Following this, using the same cohort, Covinsky and colleagues (2009) examined the relationship between functional limitations and pain, and reported that participants with significant pain were at much higher risk of reporting functional limitations.

The understanding of the determinants of the relationship between joint pain and ADL disability has evolved with successive studies. In 2008, Machado et al

examined the roles that factors such as activity limitation and depression played on participation restriction in older adults. This study highlighted the importance of making allowances for both physical and psychological consequences of OA, suggesting activity limitation and depressive symptoms mediated the relationship between physical symptoms and subsequent participation restrictions (Machado et al., 2008).

To summarise, previous studies have identified joint specific characteristics (e.g. pain severity) as important risk factors, which increased poor physical functioning and predict future ADL disability in older adults. Other health and psycho-social factors (e.g. depression) could also influence the relationship between joint pain/arthritis and participation restriction, supporting the use of biopsychosocial model when investigating the consequences of health conditions. Thus, this chapter investigates the course of self-care restriction in older people with joint pain. Building on the findings reported in the previous chapter, the analysis will examine the associations between the risk of onset and persistence of self-care restriction and key factors independently associated with restricted self-care in older people at baseline, and test the predictive ability of the parsimonious model to determine whether joint specific characteristics make an additional contribution to the prediction of future self-care restriction.



## **7.2 Aims and Objectives**

The overall aim of this chapter was to identify potential predictors of the onset and persistence of restricted self-care in older adults with joint pain. By investigating the association between joint specific characteristics and the key factors identified as being associated with restricted self-care, this would establish whether the risk profile of those with joint pain differs from the general population. Using baseline and 3-year data from the NorStOP in community-dwelling adults aged 50 years and over with joint pain at baseline the objectives were:

- i. To examine the association between risk of onset and of persistence of self-care restriction at 3 years and joint specific characteristics at baseline, such as pain severity, stiffness and chronicity
- ii. To investigate links between risk of onset and persistence of self-care restriction at 3 years and individual factors that were independently associated with restricted self-care in the cross-sectional analyses at baseline ('the parsimonious model')
- iii. To test the extent to which the parsimonious model predicts the risk of onset and persistence of self-care restriction at 3 years and to determine whether joint-specific characteristics make an additional independent contribution to the prediction of future self-care restriction

## **7.3 Methods**

### **7.3.1 Dataset**

This chapter used NorStOP baseline and 3-year follow-up data. The eligibility criteria were:

- ✓ Responded to NorStOP baseline Health Survey
- ✓ Reported hand, hip, knee or foot pain in the past 12 months
- ✓ Consented to further contact
- ✓ Responded to 3-year follow-up Health Survey
- ✓ Completed the Keele Assessment of Participation (KAP) item at baseline and 3-year follow-up

Of the 4089 participants included in the analysis described in chapter 6, 3147 met these criteria.

### **7.3.2 Outcome**

As in chapter 6, the outcome variables were (i) onset of self-care restriction at 3 years and (ii) persistence of self-care restriction at 3 years. The term 'onset' is used to define the reporting of self-care restriction at 3 years in people who were free of self-care restriction at baseline. 'Persistence' is used to define the reporting of self-care restriction at 3 years in people who were restricted in self-care at baseline.

### 7.3.3 Covariates

#### *Joint-specific characteristics*

For each of the four selected peripheral joint sites, (hand, hip, knee, foot) variables on the chronicity and severity of pain, as well as the severity of stiffness (hand, hip, knee only) from the baseline Regional Pains Survey were used.

The Pain and Stiffness subscales of the Australian/Canadian Osteoarthritis Hand Index (AUSCAN: Bellamy et al., 2002) were used for **severity of hand pain and hand stiffness**. The Pain subscale comprises 5 items (*how much pain do you have in your hands –a) at rest, b) when gripping, c) when lifting, d) when turning, e) when squeezing?*). Each item has the following response options: ‘none’, ‘mild’, ‘moderate’, ‘severe’ or ‘extreme’. Hand stiffness comprises 1 item, (*how severe is stiffness in your hands after first wakening in the morning?*) with the same item response options as for the Pain items. For these analyses, hand pain and stiffness were each dichotomised into ‘non-severe’ (no items rated as ‘severe’ or ‘extreme’) or ‘severe’ (one or more items rated as ‘severe’ or ‘extreme’).

The Pain and Stiffness subscales of the Western Ontario & McMaster Universities Osteoarthritis Index (WOMAC: Bellamy, 1988) were used for **severity of hip pain and severity of hip stiffness**. The Pain subscale comprises 5 items (*walking on a flat surface / going up or downstairs / at night while in bed / sitting or lying / standing upright*). Each item has the following response options: ‘none’, ‘mild’, ‘moderate’, ‘severe’ or ‘extreme’. Hip stiffness comprises 2 items, (*how severe is your stiffness after waking in the morning? / how severe is your stiffness after*

*sitting, lying or resting in the day?*) with the same item response options as for the Pain items. Hip pain and stiffness were each dichotomised into 'non-severe' (no items rated as 'severe' or 'extreme') or 'severe' (one or more items rated as 'severe' or 'extreme').

A separate WOMAC was completed for knee pain, scored and dichotomised as above, giving two variables: **severity of knee pain** (non-severe, severe), **severity of knee stiffness** (non-severe, severe).

**The severity of foot pain** was measured using the Pain Intensity subscale of the Manchester Foot Pain & Disability Index (FPDI: Garrow et al., 2003). This subscale comprises 5 items (*because of pain in my feet: my feet are worse in the morning / my feet are more painful in the evening / I get shooting pains in my feet / I am unable to carry out my previous work / I no longer do all my previous activities*). Each item is rated as 'none of the time', 'on some days', or 'most/every day'. Foot pain was dichotomised into 'non-severe' (no items were rated as 'most/every day') or 'severe' (one or more items were rated as 'most/every day').

Chronicity of hand, hip, knee and foot pain were each defined as pain present for less than 3 months (non-chronic) or 3 months or more (chronic) in the past 12 months.

Each participant was classed as having non-severe or severe joint pain, as having non-severe or severe joint stiffness (hand, hip, knee only), and as having chronic

or non-chronic joint pain. For participants with joint pain in only one site, these values were for that site. For participants with multiple joint pains (e.g. hand and knee joint pain) the highest value from their site-specific measures was used. For example, if hand pain was classed as non-severe from AUSCAN and knee pain was classed as severe from WOMAC, that participant would be classed as having severe pain.

#### *Variables in the parsimonious model*

In the cross-sectional analysis of the baseline data (Chapter 5), a parsimonious model of factors independently associated with restricted self-care in older people in the general population, was derived and validated for use in the general population of adults aged 50 years and over. The factors were:

- i. age: categorised as 50-59, 60-69, 70-79, and 80 $\geq$
- ii. activity limitation: measured using the PF-10 subscale of the SF-36 (0-100: Ware & Sherbourne, 1992) and categorised into tertiles (rather than quartiles as used in the baseline analysis). Categories were highest limitation:  $\geq 35$ ; 2nd highest limitation: 35.1-70.0; and lowest limitation: 70.1-100 respectively. This allowed a better distribution of scores in each category as the population at 3 years follow-up was much smaller than the baseline population due to study attrition
- i. depression: measured using the Hospital Anxiety and Depression Scale (0-21: Zigmond & Snaith, 1981) and categorised into none (0-7), borderline (8-10), and definite (11-21) cases

- ii. cognitive complaint: measured using the cognitive and alertness behaviour subscale of the Functional Limitation Profile (FLP) (0-100: Bergner et al., 1981) and categorised into three groups (no cognitive impairment (0) / high levels of cognitive impairment (22.5-100) / low levels of cognitive impairment (0.1-22.4)) based on distribution of scores
- iii. perceived adequacy of income: a single item with four response options (find it strain to get by from week to week / have to be careful with money / able to manage without much difficulty / quite comfortably off) (Thomas 1999)
- iv. educational attainment: a single item asking about qualifications gained through study as an adult (yes/no).

#### **7.3.4 Statistical Analysis**

##### *Onset and persistence of self-care restriction at 3 years in older adults with joint pain*

In this chapter, the estimates of the risk of onset and persistence of self-care restrictions were based on those with joint pain, which is 76% of the general population sample used in Chapter 6 for the onset analysis (n=2857 cf n=3770), and 91% of the general population sample used in Chapter 6 for the persistence analysis (n=290 cf n=319). Estimates were expressed as percentages.

*Association of the risk of onset and persistence of self-care restriction at 3 years with joint-specific characteristics at baseline*

Cross-tabulations of onset and persistence of self-care restriction at 3 years with joint-specific characteristics (joint-specific pain severity, stiffness severity and chronicity) at baseline were performed. The direction and strength of associations between joint-specific characteristics, the risk of onset, and risk of persistence of self-care restriction were examined using logistic regression to identify which factors at baseline predicted self-care status at 3 years. Results were summarised as crude odds ratios with 95% confidence intervals, and then adjusted for all covariates (factors in the parsimonious model).

*Factors associated with the risk of onset and persistence of restricted self-care*

At first, associations between the risk of onset and persistence of restricted self-care at 3 years and each of the factors from the parsimonious model, and joint specific characteristics were investigated using univariable logistic regression. Then, associations between self-care restriction at 3 years and the factors from the parsimonious model were adjusted for the other factors in the model using multivariable logistic regression. The association between the risk of onset and persistence of restricted self-care at 3 years and the parsimonious model was then adjusted for the joint specific characteristics by including joint pain severity, stiffness, and chronicity. Associations were presented as odd ratios with 95% confidence intervals, first unadjusted then adjusted for the non-pain model, and then fully adjusted for the pain model in the multivariable analysis.

### *Testing the predictive ability of the non-pain and pain models*

The predictive power of the parsimonious model was validated at baseline, using both internal and external validation methods (see Chapter 5). In this study, the extent to which the parsimonious model predicts the risk of self-care restriction at 3 years was investigated using multivariable logistic regression. Furthermore, the effects of the addition of joint specific characteristics to this model were also examined using multivariable logistic regression. Previously it was suggested that adjusting for pain may improve models of disability, strengthening associations once the model accounted for pain that was frequent or severe (Adamson et al., 2003).

An important feature of the prediction model is assessing the goodness of fit. This examines the agreement between observed probabilities and predicted probabilities. It is suggested that a predictive model in which the number of observed cases supports the number of cases predicted by the model, indicates good calibration. Good calibration is crucial for attaining a realistic risk adjustment (Van Houwelingen and Le Cessie, 1990). The calibration method performed in this study to examine how well the method fitted the data was the Pearson Goodness-of-fit test. The Pearson Goodness-of-fit test provides the sum of differences between observed and expected outcome frequencies (counts of observations); each squared and divided by the expectation. The resulting value can be compared to the chi-squared distribution to determine the goodness of fit (Laub and Kuhl, 2005). This calibration method was explained in detail in Chapter 5.



The c-index is a measure of discrimination, which is commonly quantified by a measure of concordance. It has the ability to correctly classify participants into high-risk and low-risk groups (Steyerberg et al., 2001). The c-index is the area under the receiver operating characteristic (ROC) curve for binary outcomes, based on the ranks ranges from 0.5 to 1, with higher values indicating better discrimination (Harrell, 2001 et al., 1996; Pepe, 2003). The c-index measure was explained in detail in Chapter 5.

In order to test the equality of the c-index obtained from applying the un-adjusted and adjusted parsimonious model (for joint specific characteristics) to the same sample, the c-index were compared executing *roccomp* command in Stata 11, using the original outcome variables (onset and persistence of self-care restriction at 3 years) as the reference variable, and the predicted probabilities of the un-adjusted and adjusted parsimonious model as classification variables. When a significant probability value is obtained, it suggests that two models are different in their predictive ability (Steyerberg et al., 2001).

The key difference between calibration and discrimination is that the discrimination reveals the ability of a given model to distinguish a status (with case / without case), while calibration measures how much the estimated values of a predictive model match the observed proportion of the event (Tripepi et al., 2009).

Predictive performances of the un-adjusted and adjusted parsimonious model were tested using multivariable logistic regression. Odds ratios with their

respective 95% confidence intervals were used to summarise the contribution of each predictor in the model. Pearson  $\chi^2$ -statistic and c-index were obtained for each model and for between models using Stata 11.

## **7.4 Results**

### **7.4.1 The risk of onset and persistence of self-care restriction at 3 years in those with joint pain**

Based on 2857 people with joint pain who were not restricted in self-care at baseline, 215 (7.5%; 95% CI 6.6, 8.6) reported the onset of self-care restriction at 3 years and of the 290 people with joint pain and self-care restriction at baseline, 115 (39.7%; CI 34.2, 45.4) reported persistence of self-care restriction at 3 years.

### **7.4.2 Association of the onset of self-care restriction at 3 years with joint specific characteristics at baseline**

In the univariable analysis there were strong associations between the risk of onset of self-care restriction at 3 years and the joint specific characteristics at baseline. The strongest association occurred in those with extreme to severe joint pain. The odds of the onset of self-care restriction at 3 years was 3 times higher in individuals reporting severe to extreme joint pain at baseline compared to those reporting none to moderate pain. This was followed by severe to extreme stiffness of the hand, hip and knee joints. The odds of self-care restriction at 3 years were 2.7 times higher in individuals reporting severe to extreme stiffness at baseline compared to those reporting none to moderate stiffness. Chronicity of joint pain at baseline was also associated with reporting onset of self-care restriction at 3

years. The odds of reporting self-care restriction at 3 years in those with chronic pain at baseline were 1.9 times higher compared to those with duration of joint pain with less than 3 months (Table 7.1).

**Table 7.1 Joint-specific characteristics at baseline and onset of self-care restriction at 3 years**

3 year response to KAP self-care item of those who were not restricted at baseline		Onset of self-care restriction at 3 years		
		No	Yes	
		N	n	OR (95%CI)
<b>Joint pain severity</b>				
	Non-severe	1505	68	1
	Severe	920	124	3.0 (2.1, 4.0)
<b>Joint pain stiffness</b>				
	Non-severe	1975	127	1
	Severe	342	59	2.7 (1.9, 3.7)
<b>Joint pain chronicity</b>				
	<3 months	768	38	1
	3 months+	1619	154	1.9 (1.3, 2.8)

\*cells do not add to 215 due to missing data

### **7.4.3 Factors associated with the onset of self-care restriction at 3 years**

In the univariable analysis, all factors from the parsimonious model at baseline were significantly associated with the risk of onset of self-care restriction at 3 years (Table 7.2). The parsimonious model was a good fit for the data (Pearson's  $\chi^2=350.33$   $p=0.88$ ), and was able to moderately discriminate between those with and without the onset of self-care restriction at 3 years (c-index: 0.76, 95% CI 0.72, 0.80). The adjusted model for the joint specific characteristics was not a good fit for the data (Pearson's  $\chi^2=951.41$   $p=0.01$ ). However, the model was still able to discriminate between cases with or without self-care restriction at 3 years (c-index: 0.73, CI 0.7, 0.80). Therefore, the addition of joint specific characteristics to the parsimonious model did not improve discrimination (Pearson's  $\chi^2=2.34$   $p=0.13$ ).

**Table 7.2 Predicting the onset of self-care restriction at 3 years, multivariable analysis**

		Unadjusted	Adjusted	Fully adjusted
		OR (95%CI)	OR (95%CI) <sup>‡</sup>	OR (95%CI) <sup>‡‡</sup>
<b>Age</b>	50-59	1	1	1
	60-69	1.5 (1.0, 2.1)	1.0 (0.7, 1.6)	1.1 (0.7, 1.7)
	70-79	2.1 (1.4, 3.0)	1.1 (0.7, 1.8)	1.2 (0.7, 2.0)
	80+	4.1 (2.6, 6.7)	1.6 (0.8, 2.9)	2.1 (1.1, 4.1)
<b>Activity limitation score (PF 10)</b>				
	lowest (70.1-100)	1	1	1
	2nd highest (35.1-70.0)	3.6 (2.3, 5.5)	2.6 (1.6, 4.2)	2.2 (1.3, 3.8)
	highest ≥35	8.2 (5.4, 12.4)	4.8 (2.9, 7.9)	3.4 (1.9, 6.1)
<b>Hospital depression (HAD) Score</b>				
	Non cases (0-7)	1	1	1
	Borderline cases (8-10)	2.4 (1.6, 3.5)	1.1 (0.7, 1.7)	1.0 (0.6, 1.7)
	Definite cases (11+)	5.1 (3.4, 7.7)	2.5 (1.5, 4.1)	2.3 (1.0, 4.0)
<b>Cognitive complaint</b>				
	No cognitive impairment	1	1	1
	Low levels	2.3 (1.6, 3.3)	1.7 (1.1, 2.6)	1.9 (1.2, 3.0)
	High levels	3.9 (2.7, 5.5)	1.8 (1.2, 2.7)	1.9 (1.2, 3.1)
<b>Perceived adequacy of income</b>				
	Comfortable	1	1	1
	Little difficulty	1.7 (1.0, 2.9)	1.4 (0.8, 2.6)	1.3 (0.7, 2.4)
	Be careful	2.6 (1.6, 4.5)	1.6 (0.8, 2.8)	1.5 (0.8, 2.7)
	Strain	3.5 (1.6, 7.7)	1.6 (0.6, 3.9)	1.1 (0.4, 3.2)
<b>Higher Educational attainment</b>				
	Yes	1	1	1
	No	2.1 (1.5, 3.0)	1.5 (1.0, 2.1)	1.5 (1.0, 2.4)
<b>Joint pain severity</b>				
	Non-severe	1	-	1
	Severe	3.0 (2.1, 4.0)	-	1.4 (0.9, 2.2)
<b>Joint pain stiffness</b>				
	Non-severe	1	-	1
	Severe	2.7 (1.9, 3.7)	-	1.2 (0.8, 1.9)
<b>Joint pain chronicity</b>				
	<3 months	1	-	1
	3 months+	1.9 (1.3, 2.8)	-	0.9 (0.6, 1.4)
<b>Goodness of fit test</b>				
	Pearson $\chi^2$	-	350.33	951.41
	P value	-	0.88	0.01
	c-index	-	0.76 (0.72, 0.80)	0.76 (0.73, 0.80)

<sup>‡</sup> Adjusted for the parsimonious model factors

<sup>‡‡</sup> Adjusted for the parsimonious model and joint specific characteristics (joint pain severity, joint pain chronicity and hand, hip and knee stiffness)

#### 7.4.4 Association of the persistence of self-care restriction at 3 years with joint specific characteristics at baseline

Associations between the risk of persistence of self-care restriction at 3 years with joint specific characteristics at baseline had similar patterns with the associations of the onset of restricted self-care at 3 years. Again, in the univariable analysis, all joint-specific characteristics at baseline were significantly associated with the persistence of self-care restriction at 3 years. The odds of self-care restriction at 3 years were two times higher in individuals reporting severe to extreme joint pain at baseline compared with people reporting none to moderate pain, 1.7 times higher in those reporting severe to extreme stiffness compared to those with none to moderate stiffness, and 1.4 times higher in those with chronic joint pain at baseline compared to those with pain lasting 3 months or less (Table 7.3).

**Table 7.3 Joint-specific characteristics at baseline and persistence of self-care restriction at 3 years**

3 year response to KAP self-care item of those who were restricted at baseline	Persistence of self- care restriction at 3 years		OR (95%CI)
	No n	Yes n	
Joint pain severity			
Non-severe	36	14	1
Severe	118	90	2.0 (1.0, 3.8)
Joint pain stiffness			
Non-severe	80	41	1
Severe	71	61	1.7 (1.0, 2.8)
Joint pain chronicity			
<3 months	27	14	1
3 months+	124	91	1.4 (1.0, 2.8)

#### **7.4.5 Factors associated with the persistence of self-care restriction at 3 years**

In the univariable analysis most factors from the parsimonious model were associated with the persistence of self-care restriction at 3 years, with the exception of 80+ category within age groups, 2<sup>nd</sup> highest score category in the activity limitation (PF10), and the absence of higher educational attainment (Table 7.4). Although these associations were mostly insignificant, this may be due to small numbers in each category.

In the multivariable analysis, the parsimonious model was a good fit for the data (Pearson's  $X^2 = 124.39$   $p = 0.22$ ). The c-index was 0.68 (95% CI 0.54, 0.70), suggesting that data is consistent with a model that has got a poor discriminative ability. Adjusting the parsimonious model for joint-specific characteristics resulted in a slight improvement in c-index value (0.69; CI 0.62, 0.77), but reduced the goodness of fit (Pearson's  $X^2 = 162.84$   $p = 0.11$ ).

Comparing c-indexes for the unadjusted and adjusted parsimonious model suggested there were no statistical difference between the models in terms of their predictive ability (Pearson's  $X^2 = 1.56$   $p = 0.21$ ).

**Table 7.4 Predicting the persistence of self-care restriction at 3 years, multivariable analysis**

		<b>Unadjusted</b>	<b>Adjusted (non-pain model)</b>	<b>Fully adjusted</b>
		OR (95%CI)	OR (95%CI) <sup>‡</sup>	OR (95%CI) <sup>‡‡</sup>
<b>Age</b>				
	50-59	1	1	1
	60-69	1.1 (0.6, 2.1)	1.2 (0.6, 2.7)	1.1 (0.5, 2.6)
	70-79	1.3 (0.7, 2.5)	1.4 (0.6, 3.2)	1.2 (0.5, 2.9)
	80+	0.7 (0.3, 1.5)	0.5 (0.2, 1.6)	0.6 (0.2, 2.0)
<b>Activity limitation score (PF 10)</b>				
	lowest (70.1-100)	1	1	1
	2nd highest (35.1-70.0)	0.9 (0.3, 2.9)	0.8 (0.2, 3.2)	0.5 (0.1, 2.4)
	highest ≥35	2.2 (0.9, 5.5)	2.0 (0.7, 6.2)	1.1 (0.3, 4.1)
<b>Hospital depression (HAD) score</b>				
	Non cases (0-7)	1	1	1
	Borderline cases (8-10)	1.8 (1.0, 3.4)	1.9 (0.9, 4.0)	2.4 (1.1, 5.5)
	Definite cases (11+)	2.6 (1.4, 4.6)	1.8 (0.9, 4.0)	2.4 (1.1, 5.4)
<b>Cognitive complaint</b>				
	No cognitive impairment	1	1	1
	Low levels	1.1 (0.5, 2.3)	0.6 (0.3, 1.5)	0.7 (0.3, 1.7)
	High levels	1.8 (1.0, 3.4)	0.8 (0.4, 1.9)	0.8 (0.3, 2.0)
<b>Perceived adequacy of income</b>				
	Comfortable	1	1	1
	Little difficulty	1.5 (0.4, 5.2)	1.8 (0.4, 8.1)	2.2 (0.5, 10.3)
	Be careful	1.9 (0.6, 6.1)	1.8 (0.4, 7.8)	1.8 (0.4, 8.2)
	Strain	4.8 (0.2, 19.1)	4.5 (0.8, 24.6)	3.5 (0.6, 20.1)
<b>Higher educational attainment</b>				
	Yes	1	1	1
	No	1.0 (0.6, 1.9)	1.1 (0.5, 2.9)	1.3 (0.6, 2.9)
<b>Joint pain severity</b>				
	Non-severe	1	-	1
	Severe	2.0 (1.0, 3.8)	-	1.2 (0.4, 3.8)
<b>Joint pain stiffness</b>				
	Non-severe	1	-	1
	Severe	1.7 (1.0, 2.8)	-	1.4 (0.7, 2.9)
<b>Joint pain chronicity</b>				
	<3 months	1	-	1
	3 months+	1.4 (1.0, 2.8)	-	0.9 (0.3, 2.4)
<b>Goodness of fit test</b>				
	Pearson $\chi^2$	-	124.39	162.84
	P value	-	0.22	0.11
	c-index (95%CI)	-	0.68 (0.54, 0.70)	0.69 (0.62, 0.77)

<sup>‡</sup> Adjusted for the parsimonious model factors

<sup>‡‡</sup> Adjusted for joint specific characteristics (joint pain severity, joint pain chronicity and hand, hip and knee stiffness)



## **7.5 Summary of results**

All joint specific characteristics were associated with the risk of onset and persistence of self-care restriction at 3 years in the univariable analysis. People with severe pain or stiffness were at a higher risk, as were those with more chronic peripheral joint pain symptoms.

Older age, greater activity limitation, more depressive symptoms, lower educational attainment, and cognitive complaint were independently associated with risk of onset of self-care restriction at 3 years. However, only depressive symptoms were associated with persistence of self-care restriction, which might be explained by the lack of power in the statistical analysis due to smaller sample.

The combination of age, activity limitation, educational attainment, perceived financial strain, cognitive complaint, and depressive symptoms was able to predict the onset of self-care restriction at 3 years with good discriminative ability. However, these were not able to predict the risk of persistence of self-care at 3 years, although again this may be related to loss of power in this analysis. Joint-specific characteristics did not add to the performance of the parsimonious model in predicting the onset or persistence of restricted self-care at 3 years.

## **7.6 Discussion**

### **7.6.1 Principal findings**

Among adults aged 50 years and older with peripheral joint pain, those with more severe, chronic symptoms are at higher risk of onset of self-care restriction, and of that restriction persisting over time. However, these joint-specific characteristics did not provide any added value in predicting the future course of self-care restriction over and above age, educational attainment, perceived financial strain, cognitive complaint, activity limitation, and depressive symptoms. The predictive ability of these latter factors was still only moderate (c-index: 0.76 for onset, 0.69 for persistence).

### **7.6.2 Critical comparison against previous studies**

Previous studies into older adults with osteoarthritis reported strong links between pain, psychological factors and disability (Machado et al., 2008; Ay and Evcik, 2008; Cook et al., 2007; Ferreira and Sherman 2007; Gignac et al., 2006; Donald and Foy, 2004; Reid et al., 2003:a-b; Cardol et al., 2002). A study conducted using 226 community-dwelling adults aged 70 years and over in New Haven, Connecticut, USA, aimed to determine the relationship between psychological factors and the occurrence of disabling musculoskeletal pain in older people using restricted activity as a disability indicator. This study found that depressive symptoms were associated with greater disability in older adults due to joint pain experienced, and identified the presence of depressive symptoms as a key factor associated with the occurrence of disabling musculoskeletal pain in community-

dwelling older people (Reid et al., 2003:b). Reid and colleagues also suggested that the adherence to treatment and rehabilitative recommendations may be lower in older people with musculoskeletal pain and depression than those without depression, which may partially explain the risk for pain-related disability (Reid et al., 2003: b). Although the definitions of pain and restriction used in Reid and colleagues' study is different to this thesis, both studies point to the strong relationship between pain, depression and restricted activity. In the NorStOP sample, those with severe and chronic symptoms were also at a higher risk of self-care disability, but the inclusion of these symptoms in predictive models of self-care restriction did not provide any added value. The odds of reporting the onset and persistence of self-care restriction in those with joint pain at baseline were still twice as high in those with depression than in those without, after the inclusion of joint specific characteristics in the predictive models. This may be due to high correlations between the depressive symptoms and severe/chronic joint pain, as strong associations with these variables were also supported by previous studies (Cook et al., 2007; Donald and Foy, 2004; Reid et al., 2003:b).

Another study that emphasised the importance of the inclusion of psychological factors when measuring the relationship between pain and disability is the Longitudinal Study of Joint Pain conducted by Donald and Foy (2004). As reported in the systematic review (Chapter 2), Donald and Foy (2004) described the natural history of joint pain in a general practice sample of adults aged 75 years and over in the UK. Study results suggested that joint pain was associated with a threefold increase in the likelihood of dependency for ADL (i.e. dressing dependency and

personal hygiene) and psychological distress. This study concluded that the frequency of joint pain fluctuates over time, and is strongly linked to psychological factors. This study also highlighted that, even though the joint destruction from arthritis would be permanent once developed, the severity and frequency of joint pain may be intermittent and that the severity of joint pain can be influenced by psychological factors (Donald and Foy, 2004). Psychological factors such as depression and cognitive impairment were key factors associated with self-care restriction in this study, and adjusting the associations between these factors and self-care restriction for severe and chronic pain did not alter these relationships. Given the results of the previous studies done by Donald and Foy (2004) and others, these results may suggest that those with long lasting severe pain may be more depressed than those who do not experience such symptoms, which then may result in a negative appraisal of their needs for self-care being met, as and when wanted. For example, an individual with chronic severe pain may experience activity limitations due to pain, as a result, become depressed, and may not be motivated to carry out the personal care activities such as washing and dressing, and caring for body parts up to their normal standard, thus perceive themselves as being restricted in self-care.

Other studies also support strong links between depression and disability. A prospective cohort study of older adults conducted by Yang Yang and George (2005) suggested that both onset and persistence of functional disability are significantly associated with increased depressive symptoms over time; recovery on the other hand is a marker of decreased depressive symptoms. As it was

observed in both Chapter 6 and 7, a substantial number of adults with restricted self-care at baseline had recovered at the time of the 3 year follow-up study. Given the previous studies of self-care disability this was an unusual finding. According to these studies, self-care disability was one of the last disabilities experienced by older people, and a marker of becoming dependent on others, thus such recovery in 3 years was unexpected. However, restriction in self-care was measured from the person's perception of their self-care needs being met 'as and when wanted' in this study, therefore the recovery did not mean that the limitations were no longer there, but it meant that the person did not perceive themselves as restricted in self-care anymore as their needs were being met as and when they had wanted. This 'changeable nature' of person-perceived participation restriction may be partially explained by the healthier cohort at 3 years, as participants had reported better physical and mental health and less depression compared with those lost to follow-up at 3 years. Thus, the back-transitions in those with person-perceived restricted self-care may be due to improved psychological factors, and / or change in the environmental factors (e.g. receipt of help and assistance or use of aids for self-care activities).

### **7.6.3 Limitations**

The sample used in this study was 77% of the general population sample used in Chapter 6. Thus, limitations discussed in the previous chapter with regards to selective loss to follow-up, and generalisability of the findings also apply to the study conducted in this chapter.

The definition of joint pain used in this study was self-reporting of joint pain. As the results have shown that more severe pain is associated with a higher risk of onset and persistence of self-care restriction at 3 years, current estimates for the risk of onset and persistence might be lower than for samples with radiographic osteoarthritis. However, this does not necessarily mean that the patterns of associations would be different for those with radiographic osteoarthritis.

## **7.7 Conclusions**

Among older adults with peripheral joint pain, those with more chronic, severe pain are at increased risk of developing self-care restriction and of this persisting. However, this appears to be due to the relationship between pain and other socio-demographic and health factors, rather than being purely the result of the symptoms of joint specific characteristics. These findings indicate that self-care restriction in those with joint pain are due to a number of reasons and may be explained by complex interactions between biopsychological factors.

Having found that restriction in self-care is multifactorial, the next chapter examines the role of environmental factors on self-care in older adults with joint pain.

**Table 7.5      Summary of key findings**

- This chapter investigated the course of self-care restriction in older people with joint pain, building on the findings reported in the previous chapter. The analyses examined the associations between the risk of onset and persistence of self-care restriction and key factors independently associated with restricted self-care in older people at baseline, and test the predictive ability of the parsimonious model to determine whether joint specific characteristics make an additional contribution to the prediction of future self-care restriction.
- The onset and persistence of self-care restriction at three years in people with joint pain was 7.5% (95% CI: 6.6, 8.6) and 39.7% (CI: 34.2, 45.4) respectively.
- Chronic joint pain, severe pain and stiffness were associated with onset and persistence of self-care restriction at 3 years (i.e. for onset, severe vs. non-severe pain: crude OR 3.0 (2.1, 4.0), chronic vs. non-chronic pain (<3 months): 1.9 (1.3, 2.8), severe vs. non-severe stiffness: 2.7 (1.9, 3.7); for persistence, 2.0 (1.0, 3.8); 1.4 (1.0, 2.8); 1.7 (1.0, 2.8) respectively).
- For all joint pain factors, adjustment for the factors associated with self-care restriction in the general population attenuated all associations. Being aged 80 years and over, lower educational attainment, higher levels of activity limitation, depression and cognitive impairment were independently associated with onset (Aged ≥80 vs. 50-59: Adj. OR 2.1 (1.1, 4.1), lower vs. higher educational attainment: 1.5 (1.0, 2.4), highest (≤35) vs. lowest activity limitation (70.1-100) 3.4 (1.9, 6.1), definite (11+) vs. non cases of depression (0-7): 2.3 (1.0, 4.0), high levels of cognitive impairment vs. no cognitive impairment: 1.9 (1.2, 3.1).
- Only depression was associated with persistence of self-care restriction (adj. OR 2.4 (1.1, 5.4)) after adjustment for all other covariates. However, this might be explained by the lack of power in the statistical analysis due to smaller sample.
- Perceived self-care restriction, as assessed by the KAP, changes over time. Older adults with chronic and severe peripheral joint pain have an increased risk of developing self-care restriction and of this persisting. In a multivariable model these joint-specific characteristics were found not to be independent predictors, suggesting that, in the absence of significant bias or confounding, the effect of joint pain on self-care restriction may be through its effects on activity limitation, depression, and cognitive complaint.
- These analyses confirm that restriction in self-care is determined by multiple factors. Therefore the next chapter considers the role environmental factors play in the relationship between self-care and joint pain.

## Chapter 8

### **The concordance between activities and participation and the impact of environmental factors on self-care**

#### **8.1 Introduction**

Throughout this thesis, self-care restriction in community-dwelling older adults is measured using an aggregated self-care item from the Keele Assessment of Participation (KAP) (Wilkie, 2005). Response to this item involves aggregation of participation in a number of tasks, where individuals interact with their environment to look after themselves “as and when they want”; this includes personal care tasks such as washing, toileting, dressing, feeding and maintaining health. Some of these specific tasks have been referred to as basic activities of daily living (ADL) (Naik et al., 2004; Gignac et al., 2000; Gill et al., 1998) and traditionally have been measured in epidemiological studies with reference to *difficulty* (e.g. none, some, a lot, unable) and *dependence* (requiring help from another person). This is in contrast to the measurement of person-perceived self-care restriction by the KAP which measures performance rather than capacity and whether ones’ needs were met “as and when they have wanted”. Within this concept of self-care, it is therefore plausible for an individual to have difficulty with a task or even to depend on others for its completion, but still to feel as though their self-care needs have been met “as and when they have wanted” (i.e. they have limited capacity (activity limitation) but they continue to participate). Throughout this thesis, similarities and differences between person-perceived self-care restriction and task-specific



activity limitation have been discussed, but the level of agreement between these two constructs has not been examined empirically.

The ICF model encourages consideration of the impact of social and environmental factors on all components of functioning and restriction, and it is important to take these factors into account when examining the occurrence and determinants of self-care restriction. Contextual factors - both personal and environmental - may help to explain the difference in the occurrence of task specific activity limitation and self-care restriction. Although some contextual factors were included in the analyses described in the previous chapters (for example occupational class, educational attainment, structural aspects of social networks, perceived adequacy of income), these are classified as “personal” factors in the ICF and are not specific to the domain of self-care. Environmental factors particular to self-care were not measured in the baseline survey or at three year follow-up. However, data on environmental factors that link specifically to self-care were collected in the 6-year follow-up study and allow examination of their role on restriction.

This chapter investigates the level of agreement between the measures of task-specific activity limitation and person-perceived self-care restriction, and studies the impact that environmental factors have on this relationship in order to gain a better understanding of the differences between self-care restriction measured by KAP, and self-care disability measured using traditional ADL measures.

## **8.2 Aims and Objectives**

The overall aims of this chapter are to establish the level of agreement between measures of task-specific activity limitation and person-perceived self-care restriction, and the impact environmental factors have on this relationship. The specific objectives of this study are:

- i. To compare the frequency and age and gender distribution of task-specific activity limitations, multiple activity limitation, and relative severity of each limitation in relation to frequency and age and gender distribution of person perceived self-care restriction
- ii. To estimate the level of agreement between estimates of person-perceived self-care restriction and task-specific activity limitation
- iii. To determine and compare the risk profiles of person-perceived self-care restriction and task-specific activity limitations
- iv. To determine whether environmental factors moderate the relationship between restricted self-care and task-specific activity limitation
- v. To investigate the associations between task-specific activity limitation and person-perceived participation restriction by the need for, and use of, environmental facilitators
- vi. To examine the socio-demographic, health and joint specific characteristics of the study sample to elucidate the wider factors associated with the need for, and use of, environmental facilitators in older adults with joint pain

## **8.3 Methods**

### **8.3.1 Dataset**

This chapter used data collected through the Health Survey Questionnaire of the NorStOP project at 6-year follow-up. This constrained the analysis to cross-sectional data from surviving respondents of the NorStOP cohort at 6 years. However, the content of the Health Survey Questionnaire at 6-year follow-up had been supplemented by a series of detailed questions on activity limitation on individual self-care tasks, as well as information on use of key environmental facilitators for self-care, notably personal help and assistance and aids and adaptations. The eligible sample were adults aged 50 years and over, reporting joint pain at 6 years follow-up, who responded to the KAP self-care item (person-perceived self-care restriction), and to at least one of the task-specific activities items (washing and bathing, dressing, caring for body parts, eating and drinking and toileting).

### **8.3.2 Outcome**

There were two outcomes of this study, (i) self-care restriction at 6 years; which was measured using the KAP self-care item (restricted in self-care/ not restricted in self-care) and (ii) task-specific activity limitation; which was measured by a single item asking participants “*Does your health now limit you in these activities?, If so, how much*” the answer options were ‘Yes, limited a lot / Yes, limited a little / No, not limited at all.’ For the purpose of the selected analyses, this item was either used as it is, or dichotomised by categorising those who were limited a little

and limited a lot into the “limited” category and those who were not limited at all into the “not limited” category.

### **8.3.3 Covariates**

The ICF framework was used to categorise variables (Table 8.1).

#### *Environmental factors*

In the NorStOP 6-year follow-up questionnaire, environmental factors were measured under the sections of ‘help and assistance’ and ‘aids and appliances’ as summarised below:

#### Help and Assistance (*Support and relationships*)

The perceived need for, and actual receipt of help and assistance were measured for each of the following task-specific activities: washing and bathing; looking after your skin, teeth, hair and nails; putting on and taking off clothes; eating and drinking and using a toilet. Participants were asked “*During the past 4 weeks have you had help from anyone else, such as family (including your spouse), friends, a health professional (e.g. nurse, occupational therapist, physiotherapist) or someone from social services (e.g. home help, care worker), to do the following activities?*” Response options were ‘No I did not need any help / No, but I have needed help / Yes, I have had help from family or friends / Yes, I have had help from a health professional or from social services’ (Table 8.1).

#### Aids and adaptations (*Products and technology for personal use in daily living*)

The use of aids and adaptations to help with selected task-specific activities was measured by asking the participants “*During the past 4 weeks have you used any of the following aids and appliances to help with your daily activities?*” Examples were; aids to help you wash and dress / aids to help you with cooking or eating / raised chair or bed / special stool / raised toilet seat / hand rails in your bathroom / bath seat / shower seat and modified bath or shower’ (Table 8.1). Response options were yes/no.

#### *Demographic, socioeconomic and health factors*

Other covariates used in the analysis to describe the characteristics of those who used environmental facilitators were chosen according to their level of association with restricted self-care (i.e. those variables from the parsimonious model), and to summarise the physical and mental health status (SF-12) of the participants (see Chapter 4 for details for these measures) (Table 8.2).

**Table 8.1 Key variables on self-care restriction, task-specific self-care activity limitation, and environmental facilitators/barriers and their mapping to ICF**

Concept	Domain/subdomain	Questionnaire item	Response options
Participation restriction	d5 Self-care	<p><i>'During the past 4 weeks, my self-care needs (examples are washing, toileting, dressing, feeding, maintaining health) have been met, <b>as and when I have wanted</b>'</i></p> <p><b>NB.</b> This item was dichotomised as restricted/ not restricted (see chapter 5 for methods)</p>	<p>All the time</p> <p>Most of the time</p> <p>Some of the time</p> <p>A little of the time</p> <p>None of the time</p>
Activity limitation	d510 Washing oneself d520 Caring for body parts d530 Toileting d540 Dressing d550 Eating / d560 Drinking	<p><i>'Does your health now limit you in these activities? If so, how much?'</i></p> <ul style="list-style-type: none"> <li>- Washing and bathing</li> <li>- Looking after your skin, teeth, hair and nails</li> <li>- Using a toilet</li> <li>- Putting on and taking off clothes</li> <li>- Eating and drinking</li> </ul>	<p>Yes, limited a lot</p> <p>Yes, limited a little</p> <p>No, not limited at all</p>
Environmental factors	e3 Support and relationships / e575 General social support services, systems and policies / e580 Health services, systems and policies	<p><i>During the past 4 weeks have you had help from anyone else, such as family (including your spouse), friends, a health professional (e.g. a nurse, occupational therapist, and physiotherapist) or someone from social services (e.g. home help, care worker) to do the following activities?</i></p> <ul style="list-style-type: none"> <li>- Washing and bathing</li> <li>- Looking after your skin, teeth, hair and nails</li> <li>- Using a toilet</li> <li>- Putting on and taking off clothes</li> <li>- Eating and drinking</li> </ul>	<p>No, I didn't need help</p> <p>No, but I have needed help</p> <p>Yes, I have had help from family or friends</p> <p>Yes, I have had help from a health professional or from social services</p>
Environmental factors	e115 Products and technology for personal use in daily living / e155 Design, construction and building products and technology of buildings for private use	<p><i>'During the past 4 weeks have you used any of the following aids and appliances to help with your daily activities?'</i></p> <ul style="list-style-type: none"> <li>- Aids to help you wash and dress</li> <li>- Aids to help you with cooking or eating</li> <li>- Raised chair or bed/special stool</li> <li>- Hand-rails in your bathroom/bath seat/shower seat</li> <li>- Modified bath or shower</li> </ul>	<p>Yes</p> <p>No</p>

**Table 8.2 Other covariates used in this chapter**

<b>Construct</b>	<b>Measure</b>	<b>Scoring</b>	<b>Reference</b>
<u>Demographic</u>	Age	Years	
	Gender	Male, Female	
<u>Socioeconomic</u>	Educational attainment	Yes/ No	
	Perceived adequacy of income	Comfortable Little difficulty Be careful Strain	(Thomas, 1999)
<u>Health Factors</u>			
Depression	HADS: Depression	0-21 (Mean: SD) +	(Zigmond & Snaith,1983)
Cognition and alertness	Functional Limitations Profile – Cognitive and Alertness behaviour subscale	No impairment Low levels High levels	(Bergner et al.1981)
Joint pain	No of peripheral joint pain	0-4	
Joint specific characteristics	Hand pain intensity Hip pain intensity Knee pain intensity Foot pain intensity	0-10*	(Von Korff et al.1992)
Physical health	SF-12 PH**	0-100	(Ware et al. 1998)
Mental health	SF-12 MH++	0-100	

+ 0 indicate no depression and 21 indicate severe depression

\* 0 indicates no pain 10 indicate pain as bad as could be

\*\* Medical Outcomes Study Short-Form 12 for physical health

++Medical Outcomes Study Short-Form 12 for mental health

#### **8.3.4 Level of agreement between activities and participation**

In previous chapters, the prevalence estimates of person-perceived self-care restriction were compared to reported estimates of task-specific activity limitation in the wider literature to discuss the implications of the study findings. Through these comparisons it was noted that similar prevalence estimates can be obtained from activity and participation measures, even though they differ conceptually. Therefore, in this analysis, this overlap is examined empirically to establish whether these are two distinct concepts in measuring self-care disability in older people.

The 'agreement' between measures of activities and participation were long debated in the previous literature and a few studies attempted to empirically distinguish between the two concepts (Badley, 2008; Whiteneck, 2006; Schuntermann 2005; Barral, 2004; Jette et al., 2003; Nordenfelt, 2003; Simeonsson et al. 2003). Together, using different methodologies, these studies provided conceptual and empirical arguments as to why activities and participation were two distinct concepts; emphasising how everyday tasks involve physical activity, and participation required individuals to interact with complex biopsychosocial factors. However, this thesis is not just another conceptual/empirical argument into differentiating between activities and participation. This thesis provides empirical support that self-care should be considered as a form of participation, and occurs as a result of complex interactions with other personal and environmental factors in the context of an



individual's life. It also suggests that participation restriction should be more appropriately measured against an individual's own standards (i.e. person-perceived "needs have met as and when I have wanted"), and not against the population norms, which may significantly vary across different socio-demographic and economic groups.

When evaluating a diagnostic test we assume that participants can ultimately be categorised as cases or non-cases. This differentiation is made by a standard test that calculates the sensitivity, specificity, and positive and negative predictive values to ascertain whether the test identifies those with or without the disease accurately (Lang and Secic, 1997). If at all possible, the diagnostic test should yield, 100% sensitive and 100% specific for the disease in question and should be applied to all participants in the study. The test should identify all cases of the disease that are of significance, but should not identify any cases that are of no significance. However, such a perfect reference standard probably does not exist. Therefore, defining the reference standard usually involves a degree of compromise. The reference standard used in this study was person-perceived self-care restriction. The analysis examines whether this captures task-specific activity limitation by task (e.g. washing and bathing limitation) and count (one or more limitations, two or more limitations), with the aim of establishing whether what the KAP measures differs to limitation in specific self-care tasks.

Introducing bias through the application of diagnostic tools may result in inflated estimates, which produce over-optimistic results within sensitivity and specificity analyses. In order to avoid bias it should be ensured that the test is evaluated in a cohort of participants presenting with suspected disease, and that the test is compared to an independent reference standard (Knottnerus et al., 2002).

It is recommended that, to estimate the potential role of chance in diagnostic tools, results should be reported with 95% confidence intervals (Porta, 2008). It is also understood that the prevalence of the disease is just as important as the larger sample size to provide more precise estimates. This is due to the fact that in diseases where the expected prevalence is low, there will be fewer people with the disease to estimate sensitivity, even though precise estimates of specificity can still be obtained (Porta, 2008).

### **8.3.5 Statistical Analysis**

#### *Main hypothesis tested*

This thesis puts forward the hypothesis that participation restriction in self-care is not the same as task-specific activity limitation in self-care. Measuring self-care disability from the perspective of person-perceived participation restriction provides estimates of the prevalence, distribution and associated factors that contrast with those obtained from more traditional self-care disability measures. To test this hypothesis, analyses were undertaken in this chapter to address specific objectives set out below.

**Objective 1:** To compare person-perceived self-care restriction and task-specific activity limitation by age and gender

This analysis aimed to summarise the prevalence of person-perceived participation restriction and task-specific activity limitation in self-care to identify similarities and differences in the occurrence and pattern of self-care disability measured by these two different constructs.

#### *Descriptive Analysis*

The relative frequency and age and gender distribution of task specific activity limitation (e.g. washing oneself, dressing oneself), multiple activity limitation (e.g. one or more, two or more...), and relative severities (limited and limited a lot) of each limitation were cross-tabulated against the relative frequency and age and gender distribution of person-perceived self-care restriction. This was conducted using frequencies and percentages to observe differences in occurrence in terms of prevalence, difficulty threshold and pattern of distribution. Due to small numbers available for analysis, differences in groups were not tested for statistical significance.

**Objective 2:** To determine the level of agreement between measures of task specific activity limitation and self-care restriction

The results of the analysis undertaken to meet the first objective may provide some evidence on the possible differences in the measurement of participation restriction and task-specific activity limitation in self-care. However, additional

analysis is required to estimate the level of agreement between these two measures, as the same prevalence estimates can be obtained from two measures even if they describe completely different people. Thus, to understand how strongly person-perceived participation restriction in self-care is related to the number, severity, and type of task-specific activity limitation, it is necessary to look at individuals.

### *Statistical Analysis*

The agreement between the two measurement constructs (KAP self-care item and task-specific self-care restriction) was tested by cross-tabulating those with task-specific activity limitation and person-perceived self-care restriction. Sensitivity and specificity analyses were then conducted as below, stratified by individual limitations (e.g. washing oneself, dressing oneself), and count of task-specific activity limitation (e.g. limitation in one or more, two or more activities) for two different cut-off points; (i) limited: those who answered as *limited a little* + *limited a lot*) and (ii) limited a lot: those who responded as *limited a lot*.

Sensitivity and specificity of the KAP measure was tested as below;

<b>Self-care restriction measured by KAP</b>	<b>Task-specific activity limitation</b>	
	limited	not limited
restricted	<b>a</b>	<b>b</b>
not restricted	<b>c</b>	<b>d</b>

a= true positive	(individuals with task-specific activity limitation detected by the test as having person-perceived self-care restriction)
b= false positive	(individuals with no task-specific activity limitation detected by the test as having person-perceived self-care restriction)
c= false negative	(individuals with task-specific activity limitation detected by the test as not having person-perceived self-care restriction)
d= true negative	(individuals with no task-specific activity limitation detected by the test as not having person-perceived self-care restriction)

Sensitivity =  $a / a+c$

*True positive/ true positive + false negative*

Specificity =  $d / b+d$

*True negative/ false positive + true negative*

Predictive value of a positive test result (PPV) =  $a / a+b$

*True positive/ true positive + false positive*

Predictive value of a negative test result (NPV) =  $d / c+d$

*True negative/ false negative + true negative*

Percentage agreement is  $(a+d) / (a+b+c+d) * 100$

**Objective 3:** To identify the risk profiles of person-perceived self-care restriction and task-specific activity limitation

Following the estimation of the level of agreement between the two constructs (objective 2), this part of the analysis aims to establish how different the risk factors are for person-perceived participation restriction and task-specific activity

limitation in self-care, and to determine whether environmental factors moderate the relationship between these two constructs.

### *Statistical analysis*

Cross-sectional associations were examined for person-perceived self-care restriction and for each factor in the parsimonious model (plus gender). Crude associations were adjusted for age and gender using multivariable logistic regression (column (a)). This analysis was repeated using task-specific activity limitation (limited in one or more task-specific activity) as the dependent variable (column (b)) to allow comparisons. Following these analysis, associations between person-perceived restricted self-care (a) and parsimonious model variables were adjusted using multivariable logistic regression; first for task-specific activity limitations (b) (as activity limitation in the parsimonious model was previously measured using PF-10 – a conceptually different tool to task-specific activity limitation in self-care as it actually measures ‘physical limitation’), secondly for the environmental factors (c), and finally both for task-specific activity limitation and the environmental factors (receipt of help and assistance and/or use of aids and adaptations) to examine factors moderating this relationship.

**Objective 4:** To determine if environmental factors moderate the relationship between task specific limitation and self-care restriction

First of all, associations between task specific activity limitation, person-perceived self-care restriction and environmental factors were assessed. Then, to

understand what determines the need for, and the use of, environmental facilitators, such as aids and assistance, the analyses aimed to identify factors associated with the use of environmental facilitators in those with task-specific activity limitation.

### *Statistical analysis*

#### Associations between those who are limited/restricted in self-care

Cross-tabulation was used to examine the associations (using percentages) between task specific activity limitation and person perceived self-care restriction. This was performed, first overall and then separately for: (a) those with no expressed need for assistance; unmet need in assistance (expressed need but non-receipt of help); and those who are in receipt of help (expressed need and receipt of help from friends and family / health professionals / social services), and (b) those who use aids/assistance and those who do not.

#### Socio-demographic, health and joint-specific characteristics

The socio-demographic, health and joint-specific characteristics of those who are limited in task-specific activities (e.g. limited in washing oneself, dressing) were cross-tabulated with help and assistance status. Frequencies and percentages to were used to allow comparisons between different groups. Differences between the groups were tested using *chi-squared* test for non-parametric variables and the *independent samples t-test* for parametric variables, and expressed as p-values.

## 8.4 Results

### 8.4.1 Comparison of person-perceived self-care restriction and task-specific activity limitation by age and gender

Of those responders with joint pain at 6 years follow-up (n=2313), 2282 (99%) had completed KAP self-care and task-specific activity limitation items; of those 58% were female and the mean age was 69 (SD: 8.9). Overall, 208 (9%) had reported person-perceived self-care restriction, and 65% of those were female with a mean age of 73 (SD: 9.1). When stratified by gender, the prevalence of restricted self-care appeared to be higher in women than men (10% cf 8%). This remained the same across the age groups, increasing with age, with the exception of the 60-69 age group category, where the gender distribution was more equal. Men reported slightly higher restriction in self-care than women (Age groups 50-59 (8% cf 4%); 60-69 (5% cf 6%); 70-79 (14% cf 9%); 80+ (20% cf 15%)).

#### ADL limitations, by count

Of 2282 responders with joint pain, 730 (32%) had task-specific activity limitation in one or more self-care activities, and 167 of those had reported severe limitations. Overall, the prevalence of task-specific activity limitation was slightly higher in women (33% cf 30%), and more women reported severe limitations (8% cf 6%). The difference between genders in reporting task-specific activity limitation narrowed with increasing number of limitations (e.g. any two or more; three or more limitations...), as well as the severity of these limitations. However the smaller numbers of participants in these groups may have affected the prevalence



estimates. When the prevalence of task-specific activity limitation was stratified by age groups and gender, similar patterns to person-perceived self-care restriction were observed. The prevalence of task-specific activity limitation increased with increasing age and was higher in women, with the exception of the 60-69 age group, where limitations were higher in men, although women reported higher rates of severe limitation (Table 8.3).

#### ADL limitations, by each ADL

In those with joint pain at 6 years, 605 (27%) had reported limitation in washing; 524 (23%) had reported limitation in dressing; 378 (17%) had reported limitation in caring for body parts; 135 (6%) had reported limitation in eating and drinking and 182 (8%) had reported limitation in toileting. Washing limitation was higher in women (28% cf 25%), and they reported more severe limitations (7% cf 4%), but there were no differences in between genders for other ADL limitations. Again, women appeared to experience more severe limitations in caring for body parts than men, whilst men reported more severe limitations in toileting (1% cf 2%).

Similar to the person-perceived self-care restriction and task-specific activity limitation by count, limitations in each task-specific task increased with age, and were higher in women in the 80 years and older age group (Table 8.3b and Figure 8.1).

**Table 8.3 Comparative occurrence and distribution by age and gender of self-care restriction and self-care activity limitation**

	All ages n=2282			50-59 n=322		60-69 n=949		70-79 n=722		80+ n=289	
	n	Female 1331 (58)	Male 951 (42)	Female 186 (58)	Male 136 (42)	Female 557 (59)	Male 392 (41)	Female 407 (56)	Male 315 (44)	Female 181 (63)	Male 108 (37)
<b>Person-perceived participation restriction n %</b>											
Self-care restriction	208	136 (10)	72 (8)	15 (8)	5 (4)	28 (5)	24 (6)	56 (14)	27 (9)	37 (20)	16 (15)
<b>Activity limitation by count n %</b>											
any 1+more	730	442 (33)	288 (30)	44 (24)	32 (24)	119 (21)	102 (26)	158 (38)	102 (32)	121 (63)	52 (46)
limited a lot	167	113 (8)	54 (6)	9 (5)	5 (4)	27 (5)	16 (4)	41 (10)	23 (7)	36 (19)	10 (9)
any 2+more											
limited	500	292 (25)	208 (24)	30 (17)	21 (17)	78 (15)	77 (21)	104 (29)	72 (25)	80 (53)	38 (39)
limited a lot	71	43 (3)	28 (3)	5 (3)	3 (2)	11 (2)	8 (2)	14 (4)	12 (4)	13 (8)	5 (5)
any 3+more											
limited	336	198 (18)	138 (17)	18 (11)	17 (14)	54 (11)	48 (14)	68 (21)	50 (19)	58 (45)	23 (28)
limited a lot	38	22 (2)	16 (2)	3 (3)	2 (2)	4 (1)	3 (1)	8 (2)	6 (2)	7 (4)	5 (5)
any 4+more											
limited	175	94 (9)	81 (11)	10 (7)	9 (8)	31 (7)	29 (9)	29 (10)	29 (12)	24 (26)	14 (19)
limited a lot	24	10 (1)	14 (2)	1 (1)	2 (2)	2 (0.4)	3 (1)	3 (1)	5 (2)	4 (3)	4 (4)
all 5											
limited	83	46 (5)	37 (5)	7 (5)	5 (5)	12 (3)	13 (4)	14 (5)	3 (6)	13 (16)	6 (9)
limited a lot	11	5 (0.4)	6 (1)	1 (1)	2 (2)	2 (0.4)	1 (0.3)	1 (0.3)	2 (1)	1 (1)	1 (1)

limited: represents those who are limited a little + limited a lot

limited a lot: only represents those who are limited a lot

columns do not add due to missing data

**Table 8.3b Comparative occurrence and distribution by age and gender of self-care restriction and self-care activity limitation**


		All ages n=2282			50-59 n=322		60-69 n=949		70-79 n=722		80+ n=289	
		Total n=	Female 1331 (58)	Male 951 (42)	Female 186 (58)	Male 136 (42)	Female 557 (59)	Male 392 (41)	Female 407 (56)	Male 315 (44)	Female 181 (63)	Male 108 (37)
<b>Person-perceived participation restriction n %</b>												
Self-care restriction		208	136 (10)	72 (8)	15 (8)	5 (4)	28 (5)	24 (6)	56 (14)	27 (9)	37 (20)	16 (15)
<b>Activity limitation by task n %</b>												
Washing												
limited		605	373 (28)	232 (25)	32 (17)	24 (18)	94 (17)	79 (20)	134 (33)	86 (28)	113 (61)	43 (38)
limited a lot		133	94 (7)	39 (4)	7 (4)	3 (2)	22 (4)	14 (4)	34 (8)	15 (5)	31 (17)	7 (6)
Dressing												
limited		524	305 (23)	219 (23)	34 (18)	27 (20)	93 (17)	84 (22)	106 (26)	73 (23)	72 (39)	35 (32)
limited a lot		72	40 (3)	32 (3)	5 (3)	3 (2)	7 (1)	7 (2)	15 (4)	14 (5)	13 (7)	8 (7)
Caring for body parts												
limited		378	219 (17)	159 (17)	19 (10)	15 (11)	58 (11)	54 (14)	76 (19)	57 (18)	66 (36)	33 (30)
limited a lot		60	37 (3)	23 (2)	5 (3)	3 (2)	9 (2)	6 (2)	12 (3)	10 (3)	11 (6)	4 (4)
Eating and drinking												
limited		135	74 (6)	61 (6)	9 (5)	9 (7)	21 (4)	20 (5)	26 (6)	23 (7)	18 (10)	9 (8)
limited a lot		19	9 (1)	10 (1)	1 (1)	3 (2)	4 (1)	1 (0.3)	2 (1)	4 (1)	2 (1)	2 (2)
Toileting												
limited		182	101 (8)	81 (9)	15 (8)	9 (7)	28 (5)	32 (8)	31 (7)	27 (9)	27 (15)	13 (12)
limited a lot		27	13 (1)	14 (2)	1 (1)	2 (2)	4 (1)	3 (1)	4 (1)	5 (2)	4 (2)	4 (4)

limited: represents those who are limited a little + limited a lot

limited a lot: only represents those who are limited a lot

columns do not add due to missing data

**Figure 8.1 Comparison of age-related trends in prevalence of person-perceived self-care restriction (KAP restriction – solid bold black line) and number and severity of ADL limitations**



#### Summary of comparisons for Figure 8.1

- Both person-perceived self-care restriction measured by KAP and count of ADL limitations shows a general increase in prevalence with age and higher prevalence in females than males
- The KAP appears to show an age-related increase in prevalence that is not as steep as for ADL limitation, particularly for females
- In both males and females, the prevalence estimates from KAP appear similar to those reporting limitation on any 4 or more ADLs or a lot of limitation in at least one ADL. Therefore, KAP prevalence estimates may be most comparable with ADL definitions requiring either, a high threshold of limitation in any one activity or some limitation across multiple activities (Fig 8.1).

#### **8.4.2 Self-care disability in the individual: examining the level of agreement between the constructs of activity and participation**

##### **Sensitivity**

Sensitivity answers the question “if the responder has the disease, how likely is it to have a positive test?” (Lang and Secic 1997). In this analysis the sensitivity of the KAP self-care measure was first tested in those with a number of task-specific activity limitations, and then in those with domain specific activity limitations to measure the level of agreement between the two constructs.

The sensitivity of the KAP self-care item increased with the increase in number of activity limitations (e.g. any two or more, any three or more) in those who were limited in task-specific activities (defined as limited a little + limited a lot). For those who were limited in any one or more task-specific activity, only 23% (95% CI 19.7, 25.8) had reported restricted self-care. However, of those who were limited in all five activities, 56% (CI 45.3, 66.6) also reported restriction in self-care. The level of agreement between participation and activities followed a different pattern for those who were ‘limited a lot’ in a number of activities. It was observed that in this group the restriction in self-care did not increase with the number of task-specific limitations. Instead, in those with severe limitation in any one or more activities, 39% (CI 31.9, 46.7) also reported restriction in self-care and in those with severe limitations in any two or more activity limitations nearly half (49% CI 37.8, 60.8) reported restriction. However, the wide confidence intervals suggested a greater variability in reporting self-care restrictions for those with a number of severe limitations (Table 8.4).

When the KAP self-care item was tested in those with specific activity limitations (e.g. in those with washing oneself limitation), it was more sensitive in detecting those with severe limitations in washing, dressing and caring for body parts. For example, one in two who were limited a lot in dressing also reported self-care restriction, whereas only 28% (CI 24.4, 32.1) of those who were 'limited' were also restricted in self-care. However, this gap was not as big in those with eating and drinking and toileting limitations. 48% (CI 39.4, 56.2) of those with eating and drinking limitations were also reporting self-care restriction as opposed to 32% (CI 15.4, 54.0) of those who were severely limited in eating and drinking. Also, 43% (CI 36.1, 50.6) of those who were limited in toileting were also restricted in self-care, and the proportion of those who were severely limited in toileting and restricted in self-care followed similar patterns (41% CI 24.5, 59.3) (Table 8.4).

### Specificity

Specificity is the probability that a person without the disease will be correctly identified as not having the disease by the test (Porta, 2008). In this study this is interpreted as the probability of a person without the task-specific activity limitation to be correctly identified as 'not-restricted' in self-care by the KAP self-care item.

When the specificity of the KAP self-care item was tested on those with number of activity limitations, the specificity rates amongst *limited* or *limited a lot*, stratified by number of limitations, did not differ due to the definition of 'not limited' being the same for all different cut-offs. The specificity test showed that 97% (for *limited*

*group*; CI 96.0, 97.0) cf 93% (for *limited a lot group*; CI 92.0, 94.2) of those who reported *no limitations* in any one or more activity had also reported *no restriction* in self-care.

The specificity rates of the KAP self-care item remained satisfactory in those with task-specific activity limitations. The specificity rates were 91% and above for those who were limited and limited a lot in dressing, caring for body parts, eating and drinking and toileting limitations (e.g. specificity for dressing limitation; 96% of those who were *not limited* in dressing were also *not restricted* in self-care). In those with a 'washing oneself' limitation, the KAP self-care item had slightly lower rate of specificity; 77% (for *limited group*; CI 70.4, 81.7) of those who were *not limited* in washing oneself were also *not restricted* in self-care as opposed to 93% (for *limited a lot group*; CI 91.5, 93.7).

#### Positive predictive value (PPV)

The positive predictive value is the probability that a person with a positive test result is a true positive, or in other words, that they do have the disease (Porta, 2008). However, in this study, this analysis was conducted to examine the level of agreement between the measurement of activity and participation. Therefore, the PPV corresponds to the proportion who report limitation in one or more task-specific activity limitation and report self-care restriction.

As the number of activity limitations increased, the PPV decreased from 77% in those with one or more activity limitations (in limited group) (95% CI 71.3, 82.6), to

49% in those with all five activity limitations (CI 38.9, 59.0). The PPV were even lower in those who were *limited a lot*, and also decreased with increasing number of limitations. For example, 31% (CI 24.9, 37.3) of those who were restricted in self-care were also limited a lot in one or more activities, and 3% (CI 1.1, 6.7) were limited a lot in all five task-specific activities.

Highest PPV was attained in those with washing limitation for both who were limited, and limited a lot (in limited; 76% CI 69.6, 81.2 cf in limited a lot; 25% CI 19.3, 30.9). The PPV values were generally higher in those who were *limited* in task-specific activity limitations compared to those who were *limited a lot*. For example, 69% (CI 62.7, 75.1) of those who were restricted in self-care were also limited in dressing oneself, compared to 17% (CI 12.4, 22.5) who were limited a lot.

#### Negative predictive value (NPV)

The negative predictive value is defined as the probability that a person with negative test result is a true negative (Porta, 2008). Thus, in this study this means, of people reporting no self-care restriction, the proportion who reports no limitation in any task-specific activity in self-care.

73% (95% CI 71.5, 75.3) of those who reported no self-care restriction were also not limited in one or more task-specific activities, and the NPV values increased for those who were *limited* in any four or more activities (NPV: 94% CI 93.2, 95.4). For those who were *limited a lot*, NPV values varied between 95% (CI 64.1, 96.0)



for those who were limited in any one or more activities to 99% (CI 98.8, 99.6) for those who were limited in any four or more activities.

When stratified by each task, NPVs were again higher in those who were *limited a lot*, compared to those who were *limited*. 79% (CI 77.1, 80.6) of those who reported no self-care restriction were also not limited in washing oneself in the *limited* group, as opposed to 96% (CI 95.2, 96.9) in those who were *limited a lot*.

**Table 8.4 Level of agreement between person-perceived self-care restriction and task-specific activity limitation**

		% (95%CI)			
		Sn	Sp	PPV	NPV
<i>Activity limitation by count</i>					
any one or more					
	limited	23 (19.7, 25.8)	97 (96.0, 97.0)	77 (71.3, 82.6)	73 (71.5, 75.3)
	limited a lot	39 (31.9, 46.7)	93 (92.0, 94.2)	31 (24.9, 37.3)	95 (64.1, 96.0)
any two or more					
	limited	29 (25.5, 33.6)	97 (96.0, 97.0)	75 (68.7, 80.9)	82 (79.8, 83.3)
	limited a lot	49 (37.8, 60.8)	93 (92.0, 94.2)	19 (14.0, 25.0)	98 (97.6, 98.7)
any three or more					
	limited	37 (32.1, 42.5)	97 (96.0, 97.0)	72 (64.8, 78.3)	88 (86.5, 89.6)
	limited a lot	46 (31.0, 61.6)	93 (92.0, 94.2)	11 (6.7, 16.3)	99 (98.5, 99.3)
any four or more					
	limited	47 (39.7, 54.5)	97 (96.0, 97.0)	63 (54.3, 70.9)	94 (93.2, 95.4)
	limited a lot	42 (24.5, 61.2)	93 (92.0, 94.2)	7 (3.6, 11.5)	99 (98.8, 99.6)
all five activities					
	limited	56 (45.3, 66.6)	97 (96.0, 97.0)	49 (38.9, 59.0)	98 (96.9, 98.4)
	limited a lot	36 (15.2, 64.6)	93 (92.0, 94.2)	3 (1.1, 6.7)	97 (99.3, 99.8)
<i>Activity limitation by task</i>					
Washing oneself					
	limited	20 (17.3, 24.0)	77 (70.4, 81.7)	76 (69.6, 81.2)	79 (77.1, 80.6)
	limited a lot	39 (31.3, 47.8)	93 (91.5, 93.7)	25 (19.3, 30.9)	96 (95.2, 96.9)
Dressing oneself					
	limited	28 (24.4, 32.1)	96 (95.3, 97.1)	69 (62.7, 75.1)	82 (80.2, 83.5)
	limited a lot	50 (38.6, 61.4)	92 (90.8, 93.1)	17 (12.4, 22.5)	98 (97.6, 98.8)
Caring for body parts					
	limited	30 (25.7, 35.0)	95 (93.9, 95.9)	54 (47.1, 60.6)	86 (86.0, 88.9)
	limited a lot	44 (32.2, 56.7)	92 (90.6, 92.9)	13 (8.8, 18.0)	98 (97.7, 98.8)
Eating and drinking					
	limited	48 (39.4, 56.2)	93 (92.2, 94.4)	31 (25.1, 37.7)	97 (95.8, 97.3)
	limited a lot	32 (15.4, 54.0)	91 (89.9, 92.3)	3 (1.4, 6.3)	99 (98.9, 99.6)
Toileting					
	limited	43 (36.1, 50.6)	94 (92.7, 94.8)	37 (30.9, 44.1)	95 (94.1, 96.0)
	limited a lot	41 (24.5, 59.3)	91 (90.1, 92.4)	5 (3.0, 9.4)	99 (98.7, 99.5)

Sn Sensitivity; Sp Specificity; PPV Positive predictive value; NPV Negative predictive value (in each case person-perceived participation restriction is the 'reference standard') % 95% CI: Percentage with 95 percent confidence interval

#### **8.4.3 The risk profile of person-perceived self-care restriction and task-specific activity limitation**

Self-care restriction was associated with all of the variables in the parsimonious model. The strongest associations were with high levels of depression (Adj. OR 11.9; 95% CI 8.0, 17.9), perceived financial strain (8.2; CI 3.6, 18.8), and high levels of cognitive impairment (4.9; CI 3.3, 7.2) (Table 8.5; column (a)).

To compare the risk profiles, associations between limitation in one or more task-specific activities and the parsimonious model were examined. Similar to the analysis conducted using person-perceived self-care restriction as the dependent variable, age (for those aged 70 years and older), depression, cognitive impairment, perceived adequacy of income and not having adult qualifications were all associated with task-specific activity limitation, before and after adjusting for the parsimonious model in the multivariable analysis. Again, the highest associations were with those who had definite cases of depression (Adj. ORs 10.1; CI 6.8, 15.0), who found their income a 'strain' (9.1; CI 5.2, 15.7), and for those with high levels of cognitive impairment (5.6; CI 4.4, 7.1). Gender was not associated with task-specific activity limitation once adjusted for the parsimonious model (Table 8.5; column (b)).

In addition to adjustment for the parsimonious factors, associations were further adjusted for: (i) task-specific activity limitation in any one or more activities, (ii) aids and assistance and (iii) both in the multivariable analysis. Adjustment for activity limitation led to attenuation of the point estimates; female gender, cognitive

impairment and not having adult qualifications were no longer associated with restricted self-care (Table 8.5; column (a:b)). When the model was adjusted for 'aids and help and assistance', age, as well as gender, cognitive impairment and not having adult qualifications also dropped out of the model (Table 8.5; column (a:c)). Adjusting the model for both task-specific activity limitation and 'aids and help and assistance' meant that only depression and the 'being careful' category of the perceived adequacy of income remained in the parsimonious model of factors associated with restriction in self-care in those with joint pain at 6 years.

**Table 8.5 Comparison of risk profiles of person-perceived self-care restriction and task-specific activity limitation**

overall n= 2282		(a) Dependent variable: person-perceived self-care restriction++		(b) Dependent variable: task-specific activity limitation*		(a) adjusted for:		
		Crude	Adj. OR †	Crude OR	Adj. OR †	(b) Adj. OR †	(c) Adj. OR †	(b) + (c) Adj. OR †
<b>Age groups</b>								
	56-59	1	1	1	1	1	1	1
	60-69	0.9 (0.5, 1.5)	0.9 (0.5, 1.5)	1.0 (0.7, 1.3)	1.0 (0.7, 1.3)	1.1 (0.3, 2.2)	1.1 (0.6, 2.3)	1.1 (0.5, 2.1)
	70-79	2.0 (1.2, 3.3)	2.0 (1.2, 3.3)	1.8 (1.3, 2.4)	1.8 (1.3, 2.4)	1.8 (1.0, 3.6)	1.7 (0.8, 3.3)	1.6 (0.8, 3.1)
	80+	3.4 (2.0, 5.8)	3.3 (1.9, 5.7)	4.3 (3.0, 6.1)	4.3 (3.1, 6.1)	2.2 (1.0, 4.6)	1.8 (0.8, 3.7)	1.6 (0.8, 3.5)
<b>Gender</b>								
	Male	1	1	1	1	1	1	1
	Female	1.4 (1.0, 1.9)	1.4 (1.0, 1.9)	1.1 (1.0, 1.4)	1.1 (0.9, 1.3)	1.4 (0.9, 2.0)	1.2 (0.8, 1.8)	1.2 (0.8, 1.8)
<b>Depression (HAD)</b>								
	Non cases	1	1	1	1	1	1	1
	Borderline cases	3.9 (2.7, 5.8)	3.8 (2.6, 5.6)	5.4 (4.1, 7.1)	5.5 (4.1, 7.3)	2.1 (1.3, 3.4)	2.4 (1.5, 3.9)	2.1 (1.3, 3.4)
	Definite cases	11.6 (7.8, 17.1)	11.9 (8.0, 17.9)	9.5 (6.5, 14.0)	10.1 (6.8, 15.0)	5.2 (3.1, 8.6)	5.3 (3.1, 9.1)	4.6 (2.7, 7.9)
<b>Cognitive complaint</b>								
	No cognitive complaint	1	1	1	1	1	1	1
	Low levels	2.7 (1.7, 4.2)	2.5 (1.6, 4.0)	2.7 (2.1, 3.5)	2.6 (2.0, 3.4)	1.4 (0.8, 2.3)	1.3 (0.7, 2.2)	1.2 (0.7, 2.1)
	High levels	5.4 (3.7, 7.9)	4.9 (3.3, 7.2)	5.9 (4.7, 7.4)	5.6 (4.4, 7.1)	1.4 (0.8, 2.2)	1.2 (0.7, 2.1)	1.1 (0.7, 1.9)
<b>Perceived adequacy of income</b>								
	Comfortable	1	1	1	1	1	1	1
	Little difficulty	1.7 (0.9, 3.2)	1.7 (0.9, 3.2)	1.5 (1.1, 2.1)	1.5 (1.1, 2.2)	1.7 (0.7, 3.7)	1.6 (0.7, 3.8)	1.7 (0.9, 3.8)
	Be careful	3.3 (1.8, 6.1)	3.4 (1.8, 6.3)	2.8 (2.0, 3.8)	3.0 (2.2, 4.2)	2.3 (1.0, 5.2)	2.2 (1.0, 5.0)	2.2 (1.0, 4.9)
	Strain	5.8 (2.6, 13.1)	8.2 (3.6, 18.8)	6.4 (3.8, 10.8)	9.1 (5.2, 15.7)	3.2 (1.2, 9.1)	2.8 (1.0, 8.2)	2.7 (0.9, 7.8)
<b>Higher educational attainment</b>								
	Yes	1	1	1	1	1	1	1
	No	2.2 (1.6, 3.1)	1.8 (1.3, 2.5)	1.7 (1.4, 2.1)	1.4 (1.2, 1.7)	1.4 (0.9, 2.1)	1.3 (0.9, 2.0)	1.3 (0.9, 2.0)

++ Measured by KAP self-care item \* Defined as limited on at least one activity from washing, dressing, caring for body parts, eating/drinking and toileting; \*\* Defined as those who use an aid and/ help and assistance in any of the self-care tasks; †OR adjusted for age and gender

#### **8.4.4 Use of environmental facilitators**

##### Associations between ADL limitation and perceived self-care restriction

Of 730 people who were limited in one or more activities of daily living relating to self-care, 161 (22%) reported that their self-care needs were not met “as and when I wanted” (i.e. restricted self-care). Of 1552 people who were not limited in any self-care activities, 3% had reported self-care restriction (Table 8.6).

For those who were limited in at least one or more task-specific activities, restriction was greater for those who needed and received help (29%). However, 19% of those who needed help and did not receive it (unmet need) had also reported self-care restriction. In those who were not limited in any task-specific self-care activity, self-care restriction was greater in those who needed help but did not receive it (8%), compared to those who received help (5%), and those who did not need help (2%) (Table 8.6b).

Self-care restriction was greater in those who used aids and appliances, for both those who reported limitations and those who reported no limitations in task-specific self-care activities (limited: 26%; not limited: 8%), when compared to those who did not use aids and appliances (limited: 10%; not limited: 2%) (Table 8.6c).

**Table 8.6 Cross-sectional association between restricted self-care and task-specific activity limitation in at least one self-care activity or more**

N%		Restricted self-care+		
ADL++ limitations	<u>Limited in 1+ ADL</u> <u>Not limited</u>	Restricted	Not restricted	
		161	569	730
		47	1505	1552
		208	2074	2282

+ Measured by KAP self-care item

++ Defined as limited on at least one activity from washing, dressing, caring for body parts, eating/drinking and toileting

**Table 8.6b Associations between restricted self-care and task-specific activity limitation; stratified by help and assistance status**

**(i) Help & assistance = No need**

N%		Restricted self-care		
ADL limitations	<u>Limited in 1+ ADL</u> <u>Not limited</u>	Restricted	Not restricted	
		5 (4)	99	114
		27 (2)	1186	1213
		32	1285	1327

**(ii) Help & assistance = Need help but do not receive it**

N%		Restricted self-care		
ADL limitations	<u>Limited in 1+ ADL</u> <u>Not limited</u>	Restricted	Not restricted	
		32 (19)	137	169
		8 (8)	91	99
		40	228	268

**(iii) Help & assistance = Needs and receives help**

N%		Restricted self-care		
ADL limitations	<u>Limited in 1+ ADL</u> <u>Not limited</u>	Restricted	Not restricted	
		123 (29)	300	423
		10(5)	185	195
		133	485	618

**Table 8.6c Associations between restricted self-care and task-specific activity limitation; stratified by use of aids and appliances**

**(i) Do not use aids & appliances**

N%		Restricted self-care		
		Restricted	Not restricted	
<b>ADL limitations</b>	<u>Limited in 1+ ADL</u>	22 (2)	1218	1240
	<u>Not limited</u>	16 (10)	141	157
		38	1359	1397

**(ii) Uses aids & appliances**

N%		Restricted self-care		
		Restricted	Not restricted	
<b>ADL limitations</b>	<u>Limited in 1+ ADL</u>	23 (8)	259	282
	<u>Not limited</u>	141 (26)	399	540
		164	658	822

*Socio-demographic, health and joint-specific characteristics*

The next stage was to examine the socio-demographic and health characteristics of those with joint pain at 6 years, for each of the five task-specific activity limitations individually. The need and receipt of help for washing and bathing, dressing, caring for body parts, eating and drinking, and toileting was considered separately at each domain. Although the need for, and receipt of, help for each activity limitation varied, there was no notable difference across the 'help and assistance' status of people with these limitations. This might be due to the high correlation in occurrence of these limitations in people with joint pain. It was observed that there was a significant overlap between the limitations in these task-specific activities; 73% of those with washing oneself limitation also reported limitations in dressing, and 56% reported limitations in caring for oneself. Likewise,



of those who were limited in eating and drinking, 87% were limited in washing and dressing, 79% were limited in caring for body parts and 67% were limited in toileting. Thus, for the purpose of this analysis (i.e. providing descriptive characteristics of those who use environmental facilitators), and to avoid the repetitive reporting of results relating to each one of the task-specific activity limitations, only the characteristics of those who were limited in *washing oneself* is reported in this chapter.

The mean age of participants who were limited in washing oneself ranged from 72 (SD: 8.8) in those who did not need help, 73 (SD: 9.3) for those with unmet need in help, and 73 (SD: 10.1) in those who received help. There were no significant differences across the help status ( $p=0.202$ ). The group of those who needed help had a significantly higher proportion of females (unmet need: 70%; needed and received help: 67%) than the group who did not need help (55%) ( $p\leq 0.005$ ). 24% of those who were restricted in self-care and limited in washing oneself did not report the need for help and assistance, but 46% who were restricted in self-care did require and received help, although 30% had unmet needs ( $p\leq 0.001$ ). There was also a significant difference between the amount of limitation reported and the help status of the participants. Of those who 'did not need help' 91% were limited a little. However, 75% of those with unmet needs in help and 58% of those who received help were also limited a little in washing oneself ( $p\leq 0.001$ ).

There was no statistical difference in different groups in terms of their perceived adequacy of income ( $p=0.066$ ). However, those with adult qualifications needed

and used less help (not needed help; 34%, those with unmet needs; 23%, those in receipt of help; 25%) when compared to those without adults qualifications (p=0.049) (Table 8.7).

**Table 8.7      Socio-demographic characteristics of those who report limitation in washing oneself; and their help and assistance status**

Overall n=588 ▲	Not needed help n=309	Unmet need n=104	Received help <sup>†</sup> n=175	P value
Age (years): Mean (SD <sup>††</sup> )	71.9 (8.8)	73.2 (9.3)	73.2 (10.1)	p=0.202
Female (n %)	170 (55)	73 (70)	118 (67)	p≤0.005
Restricted self-care (n %)	35 (24)	45 (30)	68 (46)	p≤0.001
Limited a lot (n %)	27 (9)	26 (25)	73 (42)	p≤0.001
Perceived Adequacy of income (n %)				
Comfortable	29 (10)	7 (7)	11 (7)	
Little difficulty	108 (38)	26 (26)	54 (33)	
Be careful	124 (43)	59 (60)	89 (55)	
Strain	27 (1)	7 (7)	9 (5)	p=0.066
Higher Educational attainment (n %)				
NO	204 (66)	77 (77)	128 (75)	p=0.049

▲ Columns do not add to 588 due to missing data in covariates

† Help received from family and/ friends + health professionals and/ or social services

### *Health characteristics*

Significant differences occurred in most health indicators. Those who required help and assistance (those with unmet needs and those who received help) had higher depression scores than those who reported no need for help (Unmet need: 7.9 (3.9); Needed and received help: 8.0 (3.9) cf Not needed help: 6.5 (3.4)) ( $p \leq 0.001$ ). Those who required help also had higher levels of cognitive impairment than those who did not need help ( $p = 0.019$ ).

There were also significant differences in physical and mental health scores across the help and assistance categories. Those who did not need help had higher physical and mental health scores compared to those who required help ( $p \leq 0.001$ ). Those with unmet needs reported better physical health than those who were in receipt of help, and by contrast, those who received help and assistance reported worse mental health than those who needed help, but their needs were unmet (Table 8.8).

**Table 8.8 Health characteristics of those who report limitation in washing oneself; and their help and assistance status**

	Not needed help	Unmet need n (%)	Received help <sup>†</sup> n (%)	P value
HAD+ depression score 0-21:				
Mean (SD) <sup>††</sup>	6.5 (3.4)	7.9 (3.9)	8.0 (3.9)	p≤0.001
Cognitive Alertness (n %)				
No impairment	87 (32)	18 (19)	28 (18)	
Low levels	63 (23)	21 (23)	41 (27)	
High levels	125 (45)	53 (58)	83 (55)	p=0.019
*SF-12 Physical health 0-100:				
Mean (SD) <sup>††</sup>	29.2 (7.4)	27.5 (5.7)	25.7 (5.9)	p≤0.001
*SF-12 Mental health 0-100:				
Mean (SD) <sup>††</sup>	43.6 (11.4)	38.6 (10.3)	40.1 (11.0)	p≤0.001

<sup>†</sup> Help received from family and/ friends + health professionals and/ or social services

<sup>††</sup> Standard deviation; \* Medical Outcomes Study Short-Form 12; + Hospital Anxiety and Depression.

### *Joint specific characteristics*

Those who received help reported higher hand pain intensity than those who did not need help and those who had unmet needs (p≤0.001). Hip pain intensity did not differ significantly by the need for, or receipt of help and assistance. Those with unmet needs in help reported the highest knee pain intensity (M=6.5 SD: 3.1) and this group were significantly different to those with unmet needs and no need for help (p≤0.001). In those with foot pain, those who received help had the most intense pain (p≤0.001). There was no significant difference across the help status of those with a number of peripheral joint pains (Table 8.9).

**Table 8.9 Joint specific characteristics of those who report limitation in washing oneself; and their help and assistance status**

	Not needed help	Unmet need n (%)	Received help <sup>†</sup> n (%)	P value
Hand pain intensity Mean (SD) <sup>††</sup>	3.6 (3.1)	4.5 (3.3)	4.9 (3.4)	p≤0.001
Hip pain intensity Mean (SD) <sup>††</sup>	4.1 (3.5)	4.2 (3.8)	4.7 (3.7)	p=0.239
Knee pain intensity Mean (SD) <sup>††</sup>	5.0 (3.4)	6.5 (3.1)	5.9 (3.5)	p≤0.001
Foot pain intensity Mean (SD) <sup>††</sup>	4.0 (3.4)	4.8 (3.4)	5.4 (3.6)	p≤0.001
No. of peripheral joint pain (n %)				
1	38 (13)	4 (4)	18 (11)	
2	65 (22)	27 (27)	31 (18)	
3	88 (29)	26 (26)	43 (25)	
4	109 (36)	44 (43)	76 (46)	p=0.102

<sup>†</sup> Help received from family and/ friends + health professionals and/ or social services

<sup>††</sup> Standard deviation

## 8.5 Summary of results

Among a community sample of older adults with joint pain, person-perceived self-care restriction, as measured by the KAP tool, showed broadly similar age-related trends and prevalence as limitation across four or more self-care tasks or severe limitation in at least one self-care task. The prevalence of limitation in one or more activities of self-care was much higher than for self-care restriction (32% cf 9%).

Very few individuals who reported no limitation in self-care tasks felt that their self-care needs were not met “as and when I have wanted” (i.e. participation

restriction). However, one in five individuals reporting participation restriction reported no limitation in any of the five self-care ADLs. Conversely, only 47% of individuals reporting limitation in four or more self-care tasks reported participation restriction.

While the pattern of associations with socio-demographic and health-related variables was broadly similar for KAP and individual ADL items, there were several notable exceptions. Reporting participation restriction on the KAP appeared more likely to be associated with female gender, higher levels of depression and greater perceived inadequacy of income.

Lastly, it was shown that 28% of individuals in this sample of symptomatic older adults received help or assistance for self-care tasks, while a further 12% expressed an unmet need for such help. Joint pain intensity, cognitive function, general health, severity of limitation, and self-care restriction were generally worse among those receiving or expressing a need for help and assistance for self-care, as well as among those using aids and adaptations.

## **8.6 Discussion**

### **8.6.1 Principal findings**

The prevalence of person-perceived self-care restriction was much lower than the prevalence of task-specific activity limitation, but the patterns of occurrence in terms of age and gender distribution in older adults with joint pain was very similar.

This could be explained by the strong associations found in the previous chapters between self-care restriction and activity limitation, and from a simple analysis which suggested that activity limitation mediates the relationship between joint pain and self-care restriction (Chapter 5).

The level of agreement between the constructs of activity and participation was tested using a method designed to measure the diagnostic accuracy. Ultimately, this approach was taken to test the hypothesis that the KAP self-care item was not just an aggregated measure of self-care tasks. The convergent validity of the KAP was previously tested against other measures of participation (Wilkie et al., 2005) and it was reported that KAP had high levels of agreement with items from the Reintegration to Normal Living Index (RNL) (Wood-Dauphinee et al., 1988), and the Impact of Participation and Autonomy (IPA) (Cardol et al., 2001); two different tools that included items of participation restriction within their constructs. In this study, the diagnostic accuracy test was related to the measurement of agreement between the limitation in self-care tasks and restriction in self-care, rather than confirming the construct validity of the KAP. As expected, the sensitivity of the KAP measure increased with the increasing activity limitations, and the measure was more sensitive in detecting those with severe limitations. This finding strengthens the previous findings of strong associations between activity limitation and self-care restriction. As the number of limitations increase in task-specific activities, such as washing and dressing, and these limitations become severe, the risk of reporting self-care restriction also increases. Nevertheless, the sensitivity of KAP self-care item did not exceed 56% (limited in all five activities), suggesting

that even in those adults with multiple limitations in self-care tasks, restriction in self-care occurred in one in two people; hence the measure was not overly sensitive (e.g. it did not highlight all those with severe/multiple limitations as those with self-care restrictions, and was unable to differentiate between the two concepts). The relatively high proportion of specificity rates suggested that restriction in self-care did not occur without limitation in self-care tasks. However, the measure was less specific for those with the 'washing oneself' limitation. Again, this might suggest that those with no limitation in washing oneself might be restricted in self-care due to severe limitation in other self-care tasks, such as dressing and/or caring for body parts. The level of agreement between the constructs of person-perceived self-care restriction and task-specific activity limitation suggests there is discordance, and that measures of activity limitation yield different results to that of *person-perceived* participation restriction in self-care. That factors such as depression and perceived adequacy of income remained in the parsimonious model of factors associated with restricted self-care, after adjusting for task-specific limitations and environmental factors, suggests that these factors may impact the appraisal of needs in self-care being met as and when wanted. These further explain the changeable state of self-care restrictions in older people with joint pain as reported in Chapter 7.

The risk profiles of those with person-perceived self-care restriction and task-specific activity limitation were similar. The parsimonious model variables had similar levels of associations, with both task-specific activity limitation and self-care restriction in the univariable and multivariable analyses. This may be due to high



levels of correlation between the two constructs, as discussed earlier. When the associations between the parsimonious model of key factors associated with restricted self-care and person-perceived self-care restriction at 6 years (column (a) Table 8.5) was further adjusted for task-specific activity limitation (column (b) Table 8.5), associations with these factors were weakened and became statistically insignificant for gender, cognitive impairment and educational attainment. This could be explained by a high correlation between activity limitations and participation restriction in self-care, but could also be interpreted as evidence that activity limitation mediates the relationship between key factors associated with self-care participation, as suggested in chapter 5. Further adjustment of this relationship for aids and assistance (column (c) Table 8.5) resulted in eliminating age from the model also, removing the previously strong association between increasing age and self-care restriction. This suggests that the older people are, the more likely they are to receive aids and assistance, and despite this they are still restricted. Adjusting the model for both the task-specific activity limitation and 'aids and assistance' (column (b) + (c)) (Table 8.5) resulted in the borderline and definite cases of depression, and 'being careful' category of the 'perceived adequacy of income', remaining in the parsimonious model of factors associated with restricted self-care at 6 years. Perceived financial strain was also independently associated with restricted self-care at the baseline cross-sectional analysis and was an important predictor in estimating the onset of self-care restriction at 3 years. It was discussed in great length in previous chapters that responders with poorer backgrounds and depression may generally be less satisfied with their quality of life (Moore et al., 2005) (Chapter 5 & 7). Further more

that depression is associated with a disruption of normal emotional experience that can result in a negative bias. This means that the participant has a tendency to appraise both forthcoming and past events negatively (Henry et al., 2010). What the current analyses establish is that this may be particularly likely to happen to KAP responses compared to conventional ADL items.

Two percent of those who were restricted in self-care were not limited in any self-care activities, and neither expressed a need for help, nor used aids or adaptations. There may be a number of explanations for this discordance. One explanation for this may be the measurement error within the KAP self-care item, which may have resulted in the misclassification of those who are not restricted in self-care as restricted. The five task-specific activity limitations measured in this study may have not comprehensively captured all of the individual self-care tasks. Limitation in the non-measured tasks could lead to the report of not participating “as and when they want”. As the concept of self-care that the KAP measures is person-perceived participation, people who do not define themselves as limited in task-specific activities could still perceive themselves as being restricted in self-care. This may be due to other factors such as depression, cognitive impairment and low socio-economic income affecting the appraisal of their situation.

The analysis of the dependence on task-specific activities demonstrated that there were no significant differences with age and help status, but women needed more help and assistance for washing. The poor physical health reported by female

responders compared to those of males could explain the increased dependency in washing (PCS mean score 28.5 cf 29.4  $p=0.028$ ).

One in three people with restricted self-care and who expressed a need for help had unmet needs. The majority of those with unmet needs in self-care were those who were limited a little, whereas nearly half of those who received help were severely limited in washing oneself. This may suggest that people experience limitation in self-care tasks over a certain period of time and only get help when their limitations and/or health worsen. Those who received help and assistance were more depressed compared to those who did not need help or had unmet needs in help. This could be due to becoming dependent on others, or again as a result of negative appraisal, as those who received help also had the poorest physical and mental health compared to those who did not need help, and those with unmet needs. Again, there were significant differences between help status for washing limitation and hand, knee and foot pain. Those who received help reported more intense pain in the hand and hip joints. However, for those with knee pain, those who needed, but did not receive help (unmet need), reported higher pain intensity than others. Due to the study design it was not possible to determine the direction of this association; thus it is unknown whether the severity of joint pain in those who receive help is due to restricted activity or whether they receive help due to the severity of pain and its disabling effects.

To summarise, those who received help were older, predominantly female, with severe limitations in washing, had higher levels of cognitive impairment, and worse

physical health than those who did not receive help and assistance. It was not possible to analyse the reasons why those who were in receipt of help and assistance were still restricted in self-care with the available data in this study. One explanation might be that once the individual's health deteriorates, even if their self-care needs have been met by means of help and assistance, they may still see themselves as *dependent* on others/services as the help may not be available/accessible to them '*as and when they have wanted it*', which results in perceiving themselves as 'restricted in self-care'.

### **8.6.2 Previous studies of activities and participation**

There are no other studies known to the author which have looked at the 'agreement' between measures of activities and participation in a population sample using a diagnostic accuracy test at the time this thesis written. However, there have been considerable debates regarding the conceptual differences concerning the development of ICF, and empirical attempts to distinguish between activities and participation.

In 2003, Jette and colleagues suggested that distinct activity and participation dimensions of the ICF could be identified using physical functioning items drawn from the Late Life Function and Disability Instrument (Late Life FDI) to test. Items in the Late Life FDI are similar to ones found in ICF Chapter 4 Mobility, Chapter 5 Self-care and Chapter 6 Domestic Life (Jette et al., 2003, p:146). This was a cross-sectional analysis of a sample of 150 community-dwelling adults aged 60 years and older living in Boston, Massachusetts, USA. The survey encompassed a

wide range of physical actions, daily activities and tasks outlined in the ICF manual, such as changing and maintaining body positions; carrying, moving and handling objects; mobility and travel; and ADLs. Similarly to the NorStOP 6 years cross-sectional study, responders had a mean age of 76 years, were predominately white (84%), and female (77%). The hypothesis of the existence of a single unified physical functioning domain underlying the 48 survey items was rejected in the initial factor analysis. The variance in the sample was explained by 3-factor solution that best fit the data, and these were “Mobility Activity” (perceived difficulty in performing vigorous physical actions such as walking a mile or getting up from the floor), “Daily Activities” (physical actions involved in basic and instrumental activities of daily life), and “Social Participation” (complex behaviours such as going out to public places and working at a volunteer job). This study concluded that contrary to qualifications and concerns noted within the ICF Handbook concerning physical functioning, differentiating between Activities and Participation on the basis of their domain content could be accomplished empirically (Jette et al., 2003, p:146). Although this study considered ‘social participation’ as opposed to ‘participating in self-care’, which is comparatively one form of participation that involves interaction with the environment, it highlighted the differences between the two constructs, and emphasised that participation required more complex behaviours than the physical actions involved in conducting daily activities. The results provide empirical support for self-care being multifactorial, determined by biopsychosocial factors and can be considered as one form of participation.


Schuntermann (2005) argued that the concept of activity is only intuitively introduced in the ICF and not theoretically founded. One theory, which is likened by Schuntermann to the activity concept of the ICF, is the action theory of Nordenfelt (Nordenfelt, 2003). This theory suggests that an *action* is what people actually do (i.e. performance) in their environment, and there are three preconditions for an action; (i) the individual must have the *capacity* to perform the action, (ii) they must objectively have the *opportunity* to transform their capacity into the respective action, (iii) and they must have the *will* to perform the action. Schuntermann suggests that the concept of activity is introduced without these internal factors, and describes this as a clear deficiency in the ICF (Schuntermann, 2005). However, one way to defend the ICF's approach to defining activity as simply "*the execution of a task or action by an individual*" (WHO, 2001), would be to highlight that the ICF uses activity to define these tasks and take these internal/external factors into account under *participation* to allow the breakdown, hence thorough analysis of such restrictions is taking place. Thus, the ICF considers difficulty in executing actions at an individual (activity limitation) and a societal level (participation restriction). Consequently, three preconditions suggested in this theory, which underpin the action, fit with the findings of this thesis. For example, those with pain and impairment have a reduced capacity to perform *activities*, and the use of environmental facilitators may create the *opportunity* to transform the intrinsic capacity into a respective action by enabling those with limitations to self-care. Again, Nordenfelt's theory suggests that the *will* to perform the action is also important, thus the strong association between depression and self-care restriction may be due to lack of motivation, such as

individuals not having the will to perform these self-care tasks, or seek out ‘help and assistance’ as and when they need it. Hence, Nordenfelt’s approach brings further support for the argument that self-care is a form of participation, and restrictions in self-care cannot be determined by assessing difficulty threshold and dependency in performing these personal tasks only.

Further attempts were made by Badley (2008) to enhance the conceptual clarity of the activity and participation components of the ICF, by dividing activities and participation into three domains. These were defined as (i) acts (general things that a person can do independent of context or purpose), (ii) tasks (purposeful things that people do in daily life in a specific context), and (iii) societal involvement (individual in the context of societally defined and acknowledged areas of human endeavour). Badley suggested that there are reciprocal relationships between contextual factors acting as scene-setters, contextual factors acting as facilitators or barriers, societal involvement, tasks, and acts (Figure 8.2 and Table 8.10). However, Badley’s attempt to improve the conceptual clarity of the ICF resulted in introducing further complexity in the use of terms when the original aim of the ICF was to simplify terms, allowing flexibility for researchers / health professionals / policy makers and stake holders to adapt the ICF concepts for their own purpose”. Proposed clarification of the ICF’s activities and participation domain in Badley’s paper (2008) would not have necessarily allowed the study of person-perceived self-care restriction as the author defines the societal involvement as *“The individual in the context of societally defined and acknowledged areas of human endeavour”*, thus depicting participation from a

societal norm point of view, rather than person-perceived restriction. Furthermore, it suggests that tasks are not defined culturally. However, it could be argued that a practicing Muslim might need to wash using only the left hand when toileting, thus experience activity limitation in caring for body parts due to an impairment on one hand, whereas others would have made the use of right hand. Therefore, Badley's proposed model further complicates the classification of the simple concepts offered by ICF.

**Figure 8.2** The disability model illustrating relationship between contextual factors as scene-setters, contextual factors as facilitators and barriers, societal involvement, tasks, and acts



**Source:** Badley, 2008



**Table 8.10 Distinguishing the components of functioning disability**

Distinguishing characteristic	Body structures and functions	Activity and participation		
		Acts	Tasks	Societal involvement
Relates to	Body systems and parts	General things that a person can do independent of context or purpose.	The purposeful things people do in daily life in a specific context	Defined by the social role
Exhibited by	Involvement of 1 or more body systems	Synchrony of functioning/ perhaps with other actions	Comprises synchronized acts or other (sub) tasks	The individual as a player in socially/culturally influenced areas of life
Origin of problems	Where >1 system involved, failure of any one can result in impairment	A problem in a contributing system need not result in a problem as there may be leeway	A difficulty need not result in a problem because of personal and environmental factors	May be trade-off between opportunities and personal preferences in choice of area
Individual effects <u>cognitive</u>	No cognitive component	Cognitive contribution (but often habitual or automatic processing)	Cognitive component	On-going appraisal and evaluation of the wider environment
<u>volitional</u>	Minimal voluntary control	Voluntary (and volitional)	Purposive, usually with a specific objective	Depends external factors /personal preference
Contextual scene-setters	No, except where integral to functioning	Context implied by what is done	Defines the nature of the tasks and how they are accomplished	Physical, cultural and social setting
Cultural variation	Does not vary by culture	Does not vary by culture	Nature and accomplishment dependent on setting	Only at conceptual level
Involvement of others <u>for acquisition</u>	No	Generally acquired with maturation: practice or skill may contribute	Acquired through informal or formal learning	Acquired through informal or formal processes
<u>for performance</u>	No	No	Incidental	Integral
Contextual facilitators or barriers	No, only if integral to structure or function	Can modify somewhat	Can modify manner and extent of accomplishment	May affect choices and degree of involvement

**Source:** Adapted from Badley, 2008

In a more recent study, Pollard and colleagues (2011) explored the relationships between ICF constructs of impairment, activity limitation and participation restriction in people with osteoarthritis (OA) prior to joint replacement. These authors highlighted that only a few studies have empirically explored the relationships between the constructs in the ICF model by condition (e.g. OA). The study sample was a geographical cohort of patients with osteoarthritis from the Tayside Joint Replacement (TJR) cohort who were about to undergo hip or knee joint replacement surgery at Hospital in Scotland (n=413). Participants completed measures of impairment, activity and participation prior to the surgery at home. Confirmatory factor analysis was used to test the three factors measurement model, and structural equation modelling (SEM) was used to explore the impairment, activity and participation pathways in the ICF model (Pollard et al., 2011). The study concluded that significant ICF pathways for OA prior to joint replacement were found to be between impairment and activity, and between activity and participation, but not between impairment and participation, suggesting that activity might fully mediate the relationship between impairment and participation. Although the sample used in this study was those with severe osteoarthritis requiring surgery, as opposed to those with joint pain in the general population, the findings were similar to those reported in this thesis.

Using similar methods to Pollard et al (2011) to derive instruments, Machado and colleagues (2008) also tested a model which hypothesized that the relationship between physical symptoms and later participation restrictions among older adults with osteoarthritis is mediated by activity limitations and depressive symptoms. This was a longitudinal study of 184 community-dwelling older adults aged 55

years and over, residing in the Greater Toronto area of Ontario, Canada. Thirteen items on activity limitations and participation restrictions were derived from a set of questions about difficulty with activities of everyday life. Respondents were asked to think about performing the activity *“without any help from another person or from a special gadget or piece of equipment.”* Responses were rated on a 4-point Likert scale (where 0= no difficulty, 1= some difficulty, 2= a lot of difficulty, and 3= not able to do the activity). The study reported that physical symptoms, depressive symptoms, activity limitations, and participation restrictions were significantly correlated with one another. Physical symptoms were correlated with depressive symptoms, activity limitations and participation restrictions, and depressive symptoms were correlated with activity limitations and participation restrictions. The highest correlation occurred between activity limitations and participation restrictions (Machado et al., 2008). The findings of this study are also comparable to the study reported in this chapter in terms of highlighting the importance of taking both activity limitations and depressive symptoms into account as consequences of osteoarthritis/joint pain when establishing a pathway to the understanding of the association between impairment and an individual's participation.

In summary, previous studies provide empirical support that activities and participation are two distinct concepts; activities refer to the level of the individual and an individual's capacity and participation requires the individual to interact within the context of their environment and a multitude of other factors. The results in this analysis suggest that self-care is a complex concept determined by various

socio-demographic, health and environmental factors and that self-care is more than an individual's capacity in specific activities related to personal tasks.

### **8.6.3 Limitations**

Although a generous population sample of 2282 responders was included in the 6 years cross-sectional analysis, responders were only the mere 29% of the NorStOP baseline study sample; in which the prevalence and factors associated with restricted self-care in the general population was established in Chapter 5. Responders at 6 years were younger, predominantly women and had better physical and mental health as well as improved socioeconomic status compared to those who were lost at follow-up (Chapter 4). As self-care restriction is associated with increasing age, poor health and lower socio-economic status, the study attrition may have biased the results, thus resulted in an underestimation of the frequencies of limitation, dependency, use of aids and adaptations and associations with restricted self-care.

At 6 years, some variables were measured at different time periods than others. Limitations in selected self-care activities (e.g. washing and bathing, dressing, caring for body parts) were measured at the present time (Does your health now limit you in these activities); environmental facilitators such as the receipt of help and assistance were measured during the past month (During the past 4 weeks have you had help from anyone else to do the following activities?); depression (HADS scale) was measured in the past week (How you have been feeling in the past week?); cognitive alertness was measured in the present time; presence of joint pain was measured in the last year; and joint pain intensity was measured in

the last 6 months. Different time periods may have under or overestimated the associations found.

The environmental facilitators studied in this chapter were only measured in the 6 years follow-up. Thus, it was not possible to test the cross-sectional associations with restricted self-care longitudinally to establish the direction of these associations, and whether environmental facilitators were associated with the onset and persistence of self-care restriction in older people with joint pain or not. There were also limitations to the cross-sectional associations developed. The study sample was a healthy cohort of older people compared to baseline and 3 years, which means those with severe self-care limitations / restrictions were already lost to the study. Also, the study sample was further reduced due to limited numbers of people having specific limitations in task-specific activities, such as washing and bathing, as the use of environmental facilitators such as 'help and assistance' and 'aids and adaptations' had to be measured against these specific limitations (e.g. receiving help for washing and bathing/ using modified bath or shower). Due to small numbers in each category, those who were in receipt of help and assistance from family or friends and/or health professionals or social services were collapsed into one category (i.e. needed and received help and assistance), which meant that it was not possible to analyse whether there were any differences in these groups.

## **8.7 Conclusions**

This study provided empirical evidence to support the assumption that task-specific activity limitation and person-perceived participation restriction in self-care, measured by the KAP, are two distinct outcomes. However, these two concepts are highly correlated and thus have similar associations with known risk factors such as age, gender, depression, physical limitation, cognitive impairment, perceived adequacy of income and lower educational attainment. To an extent, the relationship between self-care restriction and known risk factors is moderated by environmental facilitators. However, the receipt of help and assistance was not sufficient for those with depression and inadequate incomes to report that their self-care needs were met 'as and when they have wanted.'

Among older adults with joint pain and difficulties in self-care tasks, there is a high level of expressed need for personal help and assistance and considerable proportion of these needs are unmet. The receipt of 'help and assistance' and use of 'aids and appliances' contributes to having their self-care needs met 'as and when they want', but is still insufficient in many cases. Additional provision of personal assistance may reduce self-care restriction. However, further investigation into the reasons for person-perceived self-care restriction is required to explain why it occurs when the need for personal help is met.

**Table 8.11      Summary of key findings**

- This chapter investigated the level of agreement between the measures of task-specific activity limitation and person-perceived self-care restriction, and studied the impact that environmental factors have on this relationship. The rationale behind this was to gain a better understanding of the differences between self-care restriction measured by KAP, and self-care disability measured using traditional ADL measures.
- Both self-care restriction measured by KAP and count of ADL limitations increased in prevalence with age. The prevalence was higher in women.
- The prevalence of self-care restriction measured by KAP was most similar with the prevalence of ADL where there was a higher threshold of limitation in any one or more activity, or some limitation across multiple activities.
- There was discordance between self-care restriction measured by the KAP, and task-specific activity limitation, suggesting these are different constructs.
- The pattern of associations with socio-demographic and health related factors was broadly similar for self-care restriction measured by KAP and individual ADL items. However, participation restriction in self-care measured by KAP was more likely to be associated with female gender, higher levels of depression and greater inadequacy of income.
- Amongst this sample of symptomatic older adults there was a high level of need for personal help and assistance, and self-care restriction was greater in those who needed help but did not receive it.
- Self-care restriction was also greater in those who used aids and appliances, compared to those who did not.
- Joint pain intensity, cognitive function, general health, severity of limitation and self-care restriction was generally worse among those receiving or expressing a need for help and assistance for self-care, as well as among those using aids and appliances.

## **Chapter 9**

### **Summary and discussion**

#### **9.1 Introduction**

The impetus for this research came from the need to prepare health and social care for the increasing number of older adults in the United Kingdom who may suffer from health conditions and disability. The number of older adults is expected to increase further over the next two decades, and with this there will be more older people with musculoskeletal conditions. Musculoskeletal conditions have been linked to increased dependency in self-care activities, which is associated with future morbidity and mortality in community-dwelling older adults. However, it is unknown whether self-care restriction is an inevitable consequence of joint pain or whether people with joint pain report more restriction in self-care than those without. This chapter presents a brief review of the principal findings in this thesis, critical discussion of key issues and reflection of the implications for future research and clinical practice.

#### **9.2 Summary of key findings**

A wide range of terms have been used to describe self-care disability which has tended to focus on the individual's capacity, with no consideration of the role of social and environmental factors. Differences in definitions of self-care disability have made comparison across studies difficult. Following a systematic review, the links between joint pain and self-care disability were still unclear and the determinants of self-care restrictions in older adults with joint pain were unknown.



The ICF was proposed as a framework that could provide a standard language to facilitate the understanding of health and its consequences (WHO, 2003). It provided a conceptual framework to define self-care restrictions in older people and investigate factors associated with restriction using a biopsychosocial model. In addition, self-care restriction was measured using a person-perceived approach in which individuals judge whether their self-care needs are met “as and when I have wanted”. The estimated prevalence of self-care restriction in adults aged 50 years and over in the general population was 11.5% (95% Confidence Interval (CI) 10.8; 12.2). Restrictions increased with age and were higher in women (% difference with 95% CI: 2.3 % 0.8, 2.7). Key factors associated with self-care restriction were identified as age, activity limitation, depression, cognitive impairment, perceived inadequacy of income and not having adult qualifications. Those with joint pain were more likely to report self-care restriction than those who did not have joint pain. Much of this relationship may be explained by other factors; for example, a simple mediation analysis indicated that activity limitation may be on the causal pathway between pain and self-care restriction, in line with the proposed conceptual model of the ICF. Of those who were free of self-care restriction at baseline, 6.6% (CI 5.9; 7.5) reported the onset of self-care restriction at 3 years. The risk of reporting onset of self-care restriction increased with age, and in those who reported lower socio-economic background, worse physical and mental health at baseline. Of those who were restricted in self-care at baseline, 38.9% (CI 33.7; 44.3) reported persistence at 3 years. The risk of reporting persistent self-care restriction was higher in men and in those who reported poor physical health and perceived financial strain at baseline. Peripheral joint pain, particularly multiple joint pains at baseline increased the risk of the onset of self-

care restriction 3 years later. Looking at older adults with joint pain, those who reported severe pain, stiffness and chronic pain (lasting >3 months) at baseline were at a higher risk of reporting both the onset and persistence of self-care restriction at 3 years. Key factors identified at baseline were also predictors of the onset of self-care restrictions in older people with joint pain at 3 years, but only depression at baseline predicted the persistence of self-care restriction.

The level of discordance in the measurements of activities and participation in self-care confirmed that these were two distinct constructs. There was a high level of expressed need for personal help and assistance among older adults with joint pain and difficulties in self-care tasks. Considerable proportions of these needs were unmet. The receipt of help and assistance and use of aids and appliances contributed to having their self-care needs met 'as and when they want' to an extent, but remained insufficient in many cases.

To conclude, self-care restriction is common in community-dwelling older people and strongly associated with joint pain. The relationship between joint pain and self-care restriction is linked by other health, socio-demographic and environmental factors. Self-care restriction measured by KAP is a changeable state, with the majority of people with self-care restrictions reporting 'back transitions' 3 years later (Table 9.1).

**Table 9.1      Key findings from the thesis**

- Self-care restriction occurs in one in ten community-dwelling older adults in the general population
- Self-care restriction increases with age and is more common in women
- Key factors associated with restricted self-care in older people in the general population are age, activity limitation, depression, cognitive impairment, inadequacy of income and not having adult qualifications.
- Joint pain is strongly associated with restricted self-care in community-dwelling older people, but the relationship is not independent from other factors
- One in twenty community-dwelling adults aged 50 years and over is at risk of reporting the onset of restricted self-care 3 years later. Onset is associated with increasing age, poor physical and mental health, multiple joint pains and low socio-economic status.
- 39% of older adults in the general population report persistent self-care restriction, which is higher in men, and is associated with poor physical health, and financial strain.
- Six out of ten adults with person-perceived self-care restriction report recovery 3 years later.
- In older adults with joint pain, severity, stiffness and pain duration are associated with both the risk of onset and persistence of self-care restriction.
- In older adults with joint pain the key factors associated with restricted self-care at baseline (i.e. age, activity limitation, depression, cognitive impairment, inadequacy of income, lower educational attainment) are also predictive of the onset of self-care restriction at 3 years. However, only baseline depression is associated with persistent self-care restriction.
- The level of discordance between person-perceived self-care restriction and task-specific activity limitation suggests these constructs measure different concepts. KAP self-care measure is able to distinguish between participation restriction in self-care, and task-specific activity limitation in self-care activities in those with joint pain.
- Amongst older adults with joint pain and difficulties in self-care tasks, there is a high level of expressed need for personal help and assistance and considerable proportion of these needs are unmet.
- The receipt of help and assistance and use of aids and appliances contributes to having a person's self-care needs met 'as and when they want' to an extent, but remains insufficient in many cases.

### **9.3 General discussion**

This section discusses the key decisions taken in this thesis and potential issues with the conceptual arguments, the systematic review, and the suitability of the use of NorStOP for secondary analysis along with the interpretation of certain findings.

#### **9.3.1 The systematic review**

The systematic review aimed to identify the existing studies of self-care restriction and joint pain in the literature and was conducted at the beginning of this PhD project in 2009. In order to identify keywords, an eliminatory literature review was carried out to find alternative descriptions to terms such as 'self-care' and 'restriction'. Two issues arose from using this approach. First of all, 'self-care' was a widely used term in the literature for various concepts such as that of personal activities of daily living (ADL). Thus, in addition to the studies of limitation in ADL, the search yielded studies of self-maintenance/management (e.g. taking your medication, looking after your garden, taking exercise, seeing your friends and staying at work, taking control of your life and how to avoid being ill) as well as psychosocial studies of self-care of one's wellbeing through the use of mediums such as psychological therapy and meditation. This meant that most studies generated using these terms were not relevant to this systematic review. The large array of terms used to define self-care showed the need to use a common language to allow comparison across studies.

Although there were a number of studies which considered limitation and dependency in individual items of self-care (e.g. bathing) in community-dwelling older people, these studies were not included in the review if there were no references made to links with musculoskeletal conditions. Nevertheless, these studies were a good source for identifying potential determinants of self-care restrictions in older people in the general population, and were used to draw comparisons with the results of the analyses when possible.

The systematic review was conducted using 11 bibliographic databases from the time of their inception to 2009. However, this search did not cover grey literature (i.e. research either unpublished or has been published in non-commercial form, such as government or organisational reports), or contact any experts with regards to unpublished studies or thesis. There could potentially be relevant working papers, government reports or policy statements issued that were unknown to the author at the time.

#### *New research on self-care*

Stineman and colleagues (2012) published a paper on staging activity limitation and participation restriction in elderly community-dwelling persons, according to difficulties in self-care and domestic life functioning. This study aimed to describe the conceptual foundation and development of activity limitation, and participation restriction staging system for community-dwelling older adults aged 70 years or older, according to the severity and types of self-care and domestic life limitations experienced. Combined stages defined 29 patterns of activity limitations

expressing the individual's potential for participating in life situations pertinent to self-care and independent community life. This study concluded that ADL and IADL stages can serve to distinguish between groups of people according to both the severity and the types of limitations experienced (Stineman et al., 2012). Similar to the traditional measurement of self-care, this study examined task-specific activities such as bathing and dressing, and included 'walking' as one of these ADL. Despite references to ICF domains and qualifiers, the definition of participation restriction and the rationale behind the choice of personal and environmental factors included in the analysis were not clearly defined. Thus, although this study builds up on the existing literature of ADL hierarchy, by providing new evidence on how to stage these activities and combine with IADL to distinguish between groups of people with limitations, the study makes no reference to the determinants of self-care, other than briefly reporting on associations between ADL stage, and other 'person and environment' related factors.

A small scale study conducted by Josephine (2010) in Australia, described activity limitations and participation restrictions in a sample of older patients (n=57) from an acute health setting. This study identified any differences in activity limitations and participation restrictions between patient sub-groups within the sample, using Functional Independence Measure (FIM<sup>TM</sup>) (Uniform Data System for Medical Rehabilitation, 1999) and the Personal Care Participation Assessment and Resource Tool (PC-PART) (Vertesi et al., 2000). The mean age of the participants were 80.3 (SD: 7.5) and 60% were female. Of 57 patients, 16 (28%) reported

mobility impairment, 21 (37%) reported frailty, and 20 (35%) reported cognitive impairment. The percentages for those reporting task-specific activity limitations in personal care, and requiring assisted supervision were: 81% in dressing; 78% in toileting; 83% in bathing; 39% in eating; and 60% in grooming. Overall, 65% of mobility impairment group and 95% of the cognitive impairment group had participation restrictions in hygiene, clothing and mobility domains. The FIM<sup>TM</sup> and PC-PART reflected distinct activity limitations and participation restrictions of participants with different medical conditions. This study recommended the use of both measures of activity and participation to enable an accurate assessment of the older patient's overall functioning at home (Josephine, 2010). This study had a small sample size and the sample was derived from a hospital site, which could potentially have resulted in the increased estimates of activity limitations and participation restriction. Also the mean age of the participants was 80; an age where these restrictions are expected to be present at an even higher rate. Unfortunately the published data on this study was limited and it was not possible to contact the author as the detailed source of this study was not available through the common search engines. Nevertheless, this study considered participation restriction in personal care activities akin to the definition of self-care in this thesis, and highlighted the importance of making the conceptual distinction between measuring activity limitation and participation restriction in assessing function in older people, supporting the findings of this thesis.

Meanwhile some studies reported 'self-neglect' which is broadly defined as the inability to perform essential self-care tasks threatening a person's health and

safety. Although this concept is somewhat closer to self-management behaviour, it also covers issues with personal hygiene (e.g. washing and bathing, dressing, grooming) thus the findings could be used to compare and contrast with the findings of this thesis. According to Papaioannou et al. (2012) the exact prevalence of self-neglect in community-based older adults is unknown. Papaioannou and colleagues (2012) suggest that cognitive impairment is the most important predisposing factor of self-neglect, alongside a number of other factors such as psychiatric diseases, pre-morbid personality, alcohol abuse, poor physical ability, lack of social support and a low socioeconomic situation. The risk factors mentioned in this overview are similar to the key factors associated with restricted self-care in older people in the general population in this thesis (i.e. age, activity limitation, depression, cognitive impairment, inadequacy of income and not having adult qualifications).

Dong et al., (2012) estimated the prevalence of self-neglect across gender, race, and socioeconomic status in older people. This study showed that black older adults had a significantly higher prevalence of self-neglect than white older adults. The prevalence of self-neglect was higher in those with less than high school education, and for those with an annual income of less than \$15,000. It was not possible to determine differences across races using the NorStOP data, due to its homogeneity, thus comparisons cannot be made for these findings. However, the results of this study have shown associations between self-neglect and socioeconomic factors that are very similar to the determinants of self-care restrictions in community-dwelling older people in the NorStOP sample. Therefore, the author



suggests that it is plausible that, conceptually, there may be associations between self-care restrictions and self-neglect in older people. However, specifically designed studies need to examine the concordance between these concepts before any conclusions can be drawn from the similarities in risk factors.

### *Summary of the studies of self-care*

The systematic review of the literature identified three epidemiological studies of which two studies explored the relationship between limitation in self-care activities and joint pain, and one study considered the determinants of self-care restriction in community-dwelling older adults with joint pain. However, even in this small sample, there was variation in how these studies defined self-care, disability and joint pain. Therefore, although these studies indicated some links between joint pain and limitations in self-care activities, and pointed at the determinants of self-care disability in older people with joint pain, it was not possible to draw empirical evidence to support these links due to methodological issues which were discussed in detail in chapter 2. The results of the systematic review indicated the need for clear and consistent definitions of self-care and the need for empirical studies to investigate the extent and determinants of self-care restriction. The current research into self-care remains sporadic in terms of the concepts and definitions used to define self-care disability. This thesis is an important addition to the self-care and disability literature, by highlighting the ICF as a model that allows a common approach and helps the understanding of the occurrence, course, and determinants of self-care disability in older people with joint pain.

### 9.3.2 The concept of person-perceived self-care restriction

Most health measurement instruments do not differentiate between participation restriction and activity limitation (Wilkie et al., 2005 & 2004; Weigl et al., 2003; Fransen et al., 2002). Self-care involves the interaction between an individual and their environment. This is inherently different than the task-specific activity limitation in self-care (e.g. limitation in bathing), which measures individual's capacity to execute a task regardless of the context of the individual's situation or environmental surroundings. The WHO defines participation restriction as performance-based and context dependent, and promotes the assessment of participation restriction against an accepted population standard, that refers to what is expected for someone of the same age, gender and culture (WHO, 2001). The participation in self-care in this thesis was measured using one item from the Keele Assessment of Participation (KAP) tool that was developed to specifically measure participation restriction in population surveys (Wilkie et al., 2005). The conceptual model of participation restriction in KAP advocates that *"the experience of participation restriction is specific to each person due to the variability of roles and influencing factors; thus judgement on the occurrence of participation restriction is best perceived by the individual who experiences it"* (Wilkie et al., 2005: p.1890). Thus, the KAP does not measure participation restriction by describing the deviation from an established normative standard. Person-perceived self-care restriction measures whether the individual's needs in self-care are being met 'as and when wanted'. Therefore, the KAP measures participation restriction as perceived by the individual.

The estimated prevalence of person-perceived self-care restriction in community-dwelling older people at baseline was significantly lower than the published prevalence estimates of ADL limitations as discussed in Chapter 5. This was also true for the onset and persistence of self-care restriction at 3 years. Persistence estimates suggested that 60% of those with self-care restriction at baseline recover at 3 years. Considering that self-care disability increases with age, such a higher rate of recovery might suggest that self-care measured by KAP is an unstable state, with evidence of a substantial majority of people with self-care restriction at one point in time reporting a 'back-transition' 3 years later. Thus, the limitation of this subjective approach to participation restriction lies in the perceptual biases. This may explain the strong associations with other self-reported health and socio-demographic factors, such as depression and financial strain. A perceptual appraisal of one factor may influence the similar appraisal of another factor. The perception of a situation can be the cause of a negative psychological reaction, rather than the situation itself. To explain further, the perceived restriction may be part of a wider negative appraisal of themselves and their situation (Pincus and Morley, 2001). This can result in the underestimation of the potential determinants of self-care, and importance of such perception to actual behaviour cannot be assumed (Wilkie, 2005). Nevertheless, every measure has its limitations and subjective responses should not be considered inaccurate due to subjective factors impacting the person's appraisal of a situation.

When the levels of concordance between the task-specific activity limitation and person-perceived participation restriction were examined, analysis provided

empirical evidence to show that these are two distinct health outcomes. Nevertheless these two concepts are highly correlated, and thus have similar risk profiles. This raises the question of what the estimates of participation restriction tell us that is different from the estimates of activity limitation. Looking at these concepts at a population level, traditional methods of measuring capacity and performance in personal ADL only provide estimates of limitation and dependency; thus still refers to those who self-care with the help of environmental facilitators as disabled. This contradicts the modern understanding of disablement, which supports the measurement of actual disability (a person's ability to carry out tasks with environmental facilitators) as disability is not a personal characteristic, but is instead a gap between personal capability and environmental demand (Verbrugge, 1994). For example, the difference between the intrinsic and actual disability is not made in the Stanford Health Assessment Questionnaire (HAQ) Disability Scale (Fries et al., 1980), which is commonly used in epidemiological studies as a measure of function and disability. In this scale, higher scores indicate a greater disability. The use of environmental facilitators such as 'help and assistance' and 'aids and assistance' results in a higher score in HAQ, suggesting even those who are enabled to perform through the use of these facilitators are disabled.

The broad measure of self-care restriction provided by the KAP self-care item may be inadequate for use in studies where the outcomes require more detailed information on the nature of self-care restriction (i.e. restriction in washing and bathing or toileting specifically) for clinical or rehabilitation purposes. The dichotomous version of the KAP self-care measure does not provide information

on the levels of participation, or give any information as to whether the participation restriction occurs with or without the use of environmental facilitators. However, this is because the KAP was designed to be a crude measure of participation restriction, with the aim of being a short and easy to complete questionnaire that can be used in population surveys. There are various tools available to measure task-specific activity limitations in the literature, as discussed in previous chapters, and these tools may provide clinicians and researchers with more details on specific limitations than the KAP self-care item could offer.

### **9.3.3 The use of ICF as a framework**

The ICF framework was used in this study as a guide to define self-care, and examine the associations with health conditions, impairments, activity limitation and the impact of environmental factors in this relationship. One of the reasons behind the use of ICF as a framework in this thesis was its capacity to provide a standard method for collating and communicating health information. The ICF framework was mainly used to provide a definition of self-care restriction (hence eliminating the ambiguity of self-care) and to organise the potential determinants of self-care restriction in the community-dwelling older adults in this thesis. As discussed in the earlier chapters, problems with the definition of self-care are evident both in the current literature and policies and legislations governing services for people with health conditions. Thus, attempts to define the extent and determinants of self-care restrictions have been hindered by the absence of common language to describe health status of people with chronic conditions such as osteoarthritis. The ICF has the potential to standardise the information on

participation and osteoarthritis as a widely accepted model. It was employed in this study to provide clear guidance on methods of administering data.

The ICF framework was helpful for organising variables under different components (i.e. impairment, activities, participation and contextual factors), despite the previous criticisms made in the literature about the lack of distinction between activities and participation (Badley 2008, Imrie, 2004; Jette et al., 2003; Fougereyrollas and Beauregard, 2001). These criticisms were discussed in the previous chapter (Chapter 8).

#### *Is there a way forward?*

Although the use of the ICF as a framework in this thesis was an attempt to clearly define self-care restriction in older adults using a common language, it is evident from the new studies that the use of the ICF is still limited amongst studies of self-care disability in older adults. Criticisms of the ICF's definitions of activities and participation may be one explanation for why disability researchers are not increasingly using the ICF as a framework to inform their studies. However, the use of ADL scales to describe disability in self-care tasks is common amongst the studies of community-dwelling older people. There is a need for a culture shift in self-care disability research. It needs to change from the measure of 'disability', which only considers the individual's intrinsic capacity and dependency on others, to a measure of 'self-care' on a societal level, where interactions between the person, their health outcome, and environmental factors are considered thoroughly.

#### **9.3.4 The use of the NorStOP dataset**

NorStOP provided a practical and cost-effective alternative to primary data collection and access to a large dataset allowed investigation of self-care restrictions and joint pain in older adults. The data contained information on self-care, physical functioning, joint pain, and other important demographic, health, and socio-economic factors that were necessary to address the main objectives of this thesis. Problems associated with secondary analysis of population health data usually relate to issues such as the difficulty in locating the required data, incongruity of primary and secondary research objectives, and ensuring the data quality (Bibbs, 2007). As this thesis was undertaken in the Arthritis Research UK Primary Care Centre at Keele University, where the KAP measure was developed and the NorStOP was designed, there were no difficulties in locating the required data. As capturing joint pain, participation restriction, and associated factors in the general population of older adults were amongst the primary objectives of NorStOP, there were no issues with regards to secondary research objectives meeting the primary objectives (i.e. investigating links between joint pain and self-care restriction). The statistical analyses of the NorStOP dataset was approached in the same manner as that for the primary analysis; the data was organised and cleaned, and the quality of the data for each variable was examined by the author.

There was attrition in the NorStOP cohorts at 3 and 6-year follow-up which resulted in a healthier cohort compared to those at baseline. Responders compared to those lost at follow-up at 3 years were younger, and had better physical and mental health, better physical functioning, less depression and lower

levels of cognitive impairment. There were also more people with adult qualifications, and fewer people who found their income inadequate. Similarly, responders at 6 years follow-up were healthier than that lost-at-follow-up for each variable, and those with adult qualifications and reporting adequate income were significantly higher. In comparison to the national population of England, the NorStOP population had a higher proportion of females, and those aged 60 to 69 were over represented in the sample. However, this was more likely to be related to the slightly different population structure of the NorStOP, in terms of age and socio-economic status compared to the national population, rather than entirely due to the response bias. The NorStOP sample was derived from a population sample in the North Staffordshire region, which is one of the 10 most deprived areas in England (Indices of Deprivation, 2010). Using a population that is high in females and individuals from lower socio-economic backgrounds may have biased the results, as this can result in reports of higher levels of ill health (DoH, 2003).

Ideally, a secondary analysis of a national, or even an international dataset would have provided more generalisable results for this PhD. The generalisability of the results is an important aspect of epidemiological research, particularly when aiming to establish the commonality, and determinants of a certain health outcome, such as 'self-care restriction in community-dwelling older people' as in this thesis. Nevertheless, there was not a database available nationally or internationally that included necessary measures, such as measuring self-care as a form of participation, as defined by ICF framework at the time this study was undertaken. This thesis measured self-care at a domain level, using a novel



person-perceived participation restriction measure, which was only available through the use of NorStOP dataset. Also, the time and financial constraints of this full-time PhD would not allow conducting a primary study to include all the necessary measures in a new population study of older adults in the community and a longitudinal follow-up of this population to observe changes in their self-care status.

## **9.4 Implications**

### **9.4.1 Implications for practice (health and social care)**

This study provided an estimate of the frequency, risk of onset, persistence, and factors associated with self-care restriction in community dwelling older adults in the general population, with a particular focus on those with joint pain. As the population is ageing both in the UK and other developed countries, the number of adults with osteoarthritis is expected to rise (Verzijl et al., 2003). A better understanding of self-care restriction and its determinants is important to (i) identify the need to reduce the burden, (ii) aid the development of preventative strategies in health and social care, and (iii) identify the potential for social and clinical interventions to help those with presenting self-care restrictions.

This study identified that one in ten adults aged 50 years and over in the general population experience restriction in self-care, and those who are older, depressed, cognitively impaired, don't have adult qualifications and have inadequate incomes, are at increased risk of reporting self-care restriction. The sheer number of older adults means that there is a need to manage this form of disability. Interventions

targeted at these specific groups of older adults may help to prevent and/or reduce the restriction in self-care, which may result in decreased use of health and social care services and hospital admissions in the longer term.

The results suggest that the association between joint pain and self-care restrictions is through activity limitation. Thus, there is a need for the use of effective strategies to prevent/reduce activity limitation in community-dwelling older adults. The research evidence suggests that regular physical activity in middle-aged and older adults reduces the risk of functional limitations and disability in older age (Paterson and Warburton, 2010). It is recommended that older adults should reduce sedentary behaviour, carry out muscle-strengthening activities similar to resistance exercises which use the major muscle groups, and involve themselves in a moderate-intensity aerobic physical activity for a minimum of 30 minutes on five days each week, or vigorous-intensity aerobic activity for a minimum of 20 minutes on three days each week to promote and maintain their health (Nelson et al., 2007). Thus, early identification of older people with joint pain and activity limitations in primary care and provision of patient education may help to improve functional capabilities in this population group. Furthermore, utilisation of needs assessments in older people with joint pain to identify needs for assistance in ADL could help in provision of appropriate rehabilitative solutions. This may be through personal assistance and / or aids and adaptations to manage activity limitation in older adults before self-care restrictions occurred. Unfortunately it was not possible to examine the influence of environmental facilitators on participation restriction in self-care in this study longitudinally as the

data for help and assistance and aids and adaptations were only collected at 6 years follow-up study.

Depression was also identified as strongly associated with restricted self-care, and as a predictor of its onset and persistence over 3 years. Strong associations were also observed between depression and task-specific activity limitation in older adults with joint pain. Previous research also reports links between joint pain and depression (Yang and George, 2005; Donald and Foy, 2004). Depression in later life ( $\geq 65$  years) is associated with disability and poorer outcomes from physical illness (Manthorpe and Liffie, 2010). Evidence suggests that chronic pain and social isolation are risk factors for depression in older people (Rodda et al., 2011). Studies of depression in older people put forward that psychological interventions and antidepressants provide an effective solution for the treatment of depression in older people (Chew-Graham et al., 2011). An earlier study found that in a population of older adults with arthritis and comorbid depression, benefits of improved depression go further than reduced depressive symptoms, and result in decreased pain and improved functional status and quality of life (Lin et al., 2003). Despite the availability of effective interventions, it may be problematic to identify the older people presenting with depressive symptoms in the community. Most people with mild mental health problems such as depression are managed in primary care (Age Concern, 2006). However older people can present with non-specific symptoms such as malaise, tiredness or insomnia which may go undiagnosed (Rabins, 1996). Furthermore, the symptoms of depression may be put down to 'just getting older' (Burroughs et al., 2006). Therefore, it is important

that primary care practitioners identify older people presenting with depression when consulted for joint pain and functional limitations to support people with depression. Interventions to increase social participation, physical activity, and provision of psychological support may help to overcome depression, which may result in increased participation in self-care.

Socio-economic factors such as not having adult qualifications and inadequacy of income was also amongst the key factors associated with restricted self-care. Socio-economic status, whether assessed by income, education, or occupation, is linked to a wide range of health problems, including arthritis (Pamuk et al., 1998). Education is identified as a key to health inequality, thus, policies encouraging more years of schooling and supporting early childhood education, may have health benefits in later life (Adler and Newman, 2002). While the association between income and health is stronger in those with lower incomes, income effects persist above the poverty level. This reflects a relative status for those in the upper side of the distribution, and may be more linked to absolute deprivation for those who are at the lower end of this distribution (Backlund et al., 1999). There is growing evidence to suggest inequalities in both the distribution of health (i.e. health outcomes) and access to healthcare both internationally and in the UK (WHO, 2007). It is suggested that some health care services may not be available to certain population groups, or clinicians may have different tendencies to offer/not offer treatment to patients from different population groups, even where they have identical needs (National Audit Office, 2010; Woods et al., 2005; Goddard and Smith, 2001). Inadequacy of income was strongly associated with

restricted self-care in this study, even after adjusting for environmental facilitators. One explanation of this might be that the provision of help and assistance was inadequately distributed for those with poorer backgrounds due to the aforementioned reasons. Therefore, increasing the availability of the support from the health and social care services for those with lower socio-economic backgrounds may result in better health outcomes and thus help to increase participation in self-care.

Preliminary analysis, conducted using NorStOP 6 years cross-sectional dataset, highlighted 588 adults as having 'washing limitation'. This showed that amongst older adults with joint pain and difficulties in self-care tasks, there was a high level of expressed need for personal help/assistance which, when received, is provided largely by informal networks of friends and family rather than formal health or social care (Greenhill et al., 2011). Although it did not emerge as one of the key factors associated with restricted self-care in the multivariable analysis, social networks were significantly associated with restricted self-care even after adjusting for age and gender (adj. OR: 2.7; 95% CI 1.9, 2.8). The high levels of missing data (7.7%) in Berkman and Syme Social Network Index variable may also have influenced the significance level of these associations. Implications of this finding might be that those who are older and socially isolated would be more vulnerable, as they could not depend on the support provided by family and friends. Thus, support provided by health care and/or social services should prioritise those individuals with less social support available to them.

As discussed in the introductory chapter, occupational therapists (OT) are best placed for the holistic assessment of individuals with functional problems to develop an effective intervention plan. The main goal of the OT is to help people with activity limitations to further their skill and function, or at least maintain their current ability and prevent the deterioration of their quality of life (Turner et al., 2002). Hypothetically, the older adults that present to primary care for joint pain/osteoarthritis could be screened for early signs of task-specific activity limitation in self-care tasks (e.g. limitations in washing and bathing, toileting...) and other health factors such as depression and cognitive impairment to identify those at risk of developing restricted self-care. Those who present with the early signs of poor outcome in these measures could be referred to the community OTs to conduct needs assessments. Following a holistic assessment, OTs could provide the support and patient education required for those with pain and fatigue, and prescribe help and assistance and/or aids and adaptations needed to enable these people to overcome their limitations, thus self-care as and when they have wanted. This could help the community-dwelling older people with joint pain to cope with pain and limitations and become more independent. Consequently this would reduce primary care consultations and burden on social and health care provision within the NHS. According to the statutory legislation, local authorities in the UK provide equipment and minor works for vulnerable and disabled people (Department of Health (DoH), 2005). Prior to 2003, most local authorities used a locally developed eligibility criteria for their service. However, this method was criticised for not providing equal opportunities. Therefore in April 2003 the first national eligibility framework for adult services, namely 'Fair Access

to Care Services (FACS)' (DoH, 2002), was put in place. This involved the assessment of an individual's autonomy, health and/or safety, daily activities and ability to maintain meaningful participation in family or community life (Sackley and Lett, 2007). However, it is suggested that this service has not been implemented widely, and local authority budgets continue to determine the level of provision, with some authorities choosing not to fund specific services and equipment (Sackley and Lett, 2007). Thus, access to available OT services may be an issue for community-dwelling older adults with joint pain, depending on the provision of these services in the local area they live. Implementation of FACS nationally may help to ensure a fair distribution of these services and reduce the self-care disability in older people living in the community.

### **9.3.2 Implications for research**

The cohort suffered from response bias at the 3 and 6 years follow-up. Therefore, the analyses concerning the onset and persistence of restricted self-care and factors predicting restricted self-care in those with joint pain were conducted on a healthier cohort than those at baseline, which were representative of the general population. This might have resulted in the underestimation of the onset and persistence of self-care restriction in older people in the general population. Multiple imputation (MI) technique was used to attempt to understand and control this possible bias by examining whether the estimates based on partial and complete datasets had differed; the results of the MI analysis showed that the onset and persistence estimates of self-care restriction were not vastly different

than the estimates obtained using the original data. However, as discussed earlier in Chapter 6, the application of MI is not simple, and the approach to imputation taken in this study may not be less biased than the complete case analysis. Therefore, while participant attrition is inevitable in longitudinal studies where people are followed-up over long periods, future longitudinal studies of self-care based on a sample of older adults representative of the general population at follow-ups would increase the generalisability of the findings .

The use of a person-perceived measure of participation restriction in self-care was a novel approach for this thesis. The prevalence estimates of self-care disability were much lower compared to the previous studies' estimates which were acquired through the measure of limitation and/or dependency in self-care items. This could be due to one of two reasons. First of all, participation in self-care is a complex phenomenon which includes the interaction of biopsychosocial factors; thus people with limitation/dependency in self-care may still be 'not restricted in self-care' if their needs are met by environmental facilitators such as help and assistance or aids and adaptations. Secondly, person-perceived self-care restriction may be affected by the context they find themselves in, which may affect the judgement of their needs in self-care being met 'as and when I have wanted'. For example, an individual with depression may appraise their situation differently than a highly motivated individual just as restricted. Therefore, further studies are required to corroborate or refute the findings from this study. Further testing of the KAP self-care measure in other populations would also allow comparisons of the results and validation of what this tool really measures. The



concept of person-perceived participation restriction should also be evaluated by qualitative and quantitative studies to establish its appropriateness in use of population surveys. Further quantitative studies are also required to investigate factors mediating the relationship between joint pain and self-care restriction to build on the findings of this thesis.

Prior to this study the links between self-care restriction and joint pain in the general population of older adults were unknown, and the determinants of self-care restrictions in older people with joint pain were unclear. This study established factors that are potentially modifiable and amenable to interventions to enhance self-care in older adults, providing a basis for future studies to investigate the potential for interventions to enhance self-care in older adults with joint pain.

## References

- Adamson, J., Hunt, K. & Ebrahim, S. 2003, "Association between measures of morbidity and locomotor disability: diagnosis alone is not enough", *Social science & medicine* (1982), vol. 57, no. 8, pp. 1355-1360.
- Adamson, J., Lawlor, D.A. & Ebrahim, S. 2004. "Chronic diseases, locomotor activity limitation and social participation in older women: cross sectional survey of British Women's Heart and Health Study". *Age and Ageing*, vol. 33, no. 3, pp. 293-298.
- Adler, N.E. & Newman, K. 2002. "Socioeconomic disparities in health: pathways and policies", *Health affairs (Project Hope)*, vol. 21, no. 2, pp. 60-76.
- Age Concern 2006. [Online] Available: [http://www.ageuk.org.uk/ /health-and-wellbeing/improving\\_services\\_and\\_support\\_for\\_older\\_people\\_with\\_mental\\_health\\_problems](http://www.ageuk.org.uk/ /health-and-wellbeing/improving_services_and_support_for_older_people_with_mental_health_problems) [2012, 9/12/2012].
- Age UK. 2012. *Later life in the United Kingdom.*, Age UK. London.
- Age-standardised rate 2005. [Homepage of AIHW Knowledgebase] [Online] Available:<http://meteor.aihw.gov.au/content/item.phtml?itemId=273787&nodeId=file41fd72fab2993&fn=Age-standardised> rate, version 1, Derived DE, NHDD, NHIMG, Superseded 01/03/2005.pdf [2012, 9/9/2012].
- Age standardization by direct method. 1984. *The Journal of the Royal College of General Practitioners*, vol. 26, no. 7, pp. 7.
- Agresti, A. 2007. "An Introduction to Categorical Data Analysis ".
- Ahern, K. & Le Brocque, R. 2005. "Methodological Issues in the Effects of Attrition: Simple Solutions for Social Scientists". *Field Methods*, vol. 17, no. 1, pp. 53-69.
- Ahluwalia, S.C., Gill, T.M., Baker, D.I. & Fried, T.R. 2010. "Perspectives of older persons on bathing and bathing disability: a qualitative study" *Journal of the American Geriatrics Society*, vol. 58, no. 3, pp. 450-456.

Albrecht, G.L. 2010. "The sociology of disability: Historical foundations and future directions" in , eds. C.E. Bird, P. Conrad, A.M. Fremont & S. Timmermans, Vanderbilt University Press, Nashville, TN US, pp. 192-209.

Allan, C.M., Campbell, W.N., Guptill, C.A., Stephenson, F.F. & Campbell, K.E. 2006. "A conceptual model for interprofessional education: the international classification of functioning, disability and health (ICF)" *Journal of interprofessional care*, vol. 20, no. 3, pp. 235-245.

Allen, K.D., Jordan, J.M., Renner, J.B., Kraus, V.B. 2006. Validity, factor structure, and clinical relevance of the AUSCAN osteoarthritis hand index. *Arthritis & Rheumatism*, Volume 54, Issue 2, pages 551–556.

Andersen, R.E., Crespo, C.J., Ling, S.M., Bathon, J.M. & Bartlett, S.J. 1999. "Prevalence of significant knee pain among older Americans: results from the Third National Health and Nutrition Examination Survey" *Journal of the American Geriatrics Society*, vol. 47, no. 12, pp. 1435-1438.

Andrianakos, A., Trontzas, P., Christoyannis, F., Dantis, P., Voudouris, C., Georgountzos, A., Kaziolas, G., Vafiadou, E., Pantelidou, K., Karamitsos, D., Kontelis, L., Krachtis, P., Nikolia, Z., Kaskani, E., Tavaniotou, E., Antoniadis, C., Karanikolas, G. & Kontoyanni, A. 2003. "Prevalence of rheumatic diseases in Greece: a cross-sectional population based epidemiological study. The ESORDIG Study" *The Journal of Rheumatology*, vol. 30, no. 7, pp. 1589-1601.

Archenholtz, B. & Dellhag, B. 2008. "Validity and reliability of the instrument Performance and Satisfaction in Activities of Daily Living (PS-ADL) and its clinical applicability to adults with rheumatoid arthritis" *Scandinavian Journal of Occupational Therapy*, vol. 15, no. 1, pp. 13-22.

ARMA Statement on Supporting People with Long Term Conditions report. Arthritis and Musculoskeletal Alliance (ARMA). 2005. *Talkback Magazine*, pp. 6-6.

Armitage, P. & Berry, G. 1994. *Statistical Methods in Medical Research*, 2nd edn, Blackwell.

Arola, H.M., Nicholls, E., Mallen, C. & Thomas, E. 2010. "Self-reported pain interference and symptoms of anxiety and depression in community-dwelling older adults: can a temporal relationship be determined?" *European journal of pain* (London, England), vol. 14, no. 9, pp. 966-971.

Arthritis Research Campaign. 2002. Arthritis the big picture.

Arthritis Care : Managing OA in primary care: maximising patient consultation time 2004. [Homepage of Arthritis Care], [Online]. Available: <http://www.arthritiscare.org.uk/@3235/Forhealthprofessionals/10MinuteManager> [2012, 9/12/2012].

Arthritis and Musculoskeletal Alliance. Standards of care for people with osteoarthritis. [Homepage of Arthritis and Musculoskeletal Alliance.], [Online]. Available: <http://arma.uk.net/local-groups/care-documents/osteoarthritis/> [2012, September 4th.].

Atwal, A. & JONES, M.B.N. 2009. *Preparing for Professional Practice in Health and Social Care* Wiley-Blackwell.

Ay, S. & Evcik, D. 2008. "Effectiveness of pain, disease severity and radiological grading on disability of daily living activities in knee osteoarthritis [Turkish]" *Romatizma/Rheumatism*, vol. 23, no. 1, pp. 14-17.

Ayis, S., Arden, N., Doherty, M., Pollard, B., Johnston, M. & Dieppe, P. 2010. "Applying the impairment, activity limitation, and participation restriction constructs of the ICF model to osteoarthritis and low back pain trials: a reanalysis" *Journal of Rheumatology*, vol. 37, no. 9, pp. 1923-1931.

Backlund, E., Sorlie, P.D. & Johnson, N.J. 1999. "A comparison of the relationships of education and income with mortality: the National Longitudinal Mortality Study" *Social science & medicine*, vol. 49, no. 10, pp. 1373-1384.

Badley, E.M. & Tennant, A. 1992. "Changing profile of joint disorders with age: findings from a postal survey of the population of Calderdale, West Yorkshire, United Kingdom", *Annals of the Rheumatic Diseases*, vol. 51, no. 3, pp. 366-371.

- Badley, E.M. 2008. "Enhancing the conceptual clarity of the activity and participation components of the International Classification of Functioning, Disability, and Health". *Social science & medicine*, vol. 66, no. 11, pp. 2335-2345.
- Badley, E.M. & Ibanez, D. 1994. "Socioeconomic risk factors and musculoskeletal disability" *The Journal of rheumatology*, vol. 21, no. 3, pp. 515-522.
- Bagge, E., Bjelle, A., Valkenburg, H.A. & Svanborg, A. 1992. "Prevalence of radiographic osteoarthritis in two elderly European populations" *Rheumatology international*, vol. 12, no. 1, pp. 33-38.
- Bailey, K.M. 1998. *Learning about language assessment: Dilemmas, decisions, and directions*. Heinle & Heinle Publishers, Pacific Grove, CA.
- Ballantyne, P. & Marshall, V.W. 2001. "Subjective income security of (middle) aging and elderly Canadians" *Canadian Journal on Aging*, vol. 20, no. 2, pp. 151-173.
- Balzi, D., Lauretani, F., Barchielli, A., Ferrucci, L., Bandinelli, S., Buiatti, E., Milaneschi, Y. & Guralnik, J.M. 2010. "Risk factors for disability in older persons over 3 years follow-up", *Age and Ageing*, vol. 39, no. 1, pp. 92-98.
- Barber J, Muller S, Whitehurst T, Hay E. Measuring morbidity: self-report or health care records? *Fam Pract*. 2010 Feb;27(1):25-30. doi: 10.1093/fampra/cmp098. Epub 2009 Dec 17.
- Barberger-Gateau, P., Rainville, C., Letenneur, L. & Dartigues, J. 2000. "Hierarchical model of domains of disablement in the elderly: a longitudinal approach", *Disability and rehabilitation*, vol. 22, no. 7, pp. 308-317.
- Baron, R.M. & Kenny, D.A. 1986. "The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations", *Journal of personality and social psychology*, vol. 51, no. 6, pp. 1173-1182.

Barral, C. 2004. ICF: Internal and external obstacles to implementation, Presentation at the 10th Annual NACC Conference on ICF: Halifax, Nova Scotia, Canada, June 1–4.

Bates, S.M. & Rogstad, K.E. 2000. "Postal research: too many problems?", Sexually transmitted infections, vol. 76, no. 5, pp. 332-334.

Baum, C.M. 2002. "Creating partnerships: Constructing our future ", Australian Occupational Therapy Journal, vol. 49, no. 2, pp. 58-62.

Bayona, M. & Olsen, C. , Observational Studies and Bias in Epidemiology 2004. Available:[http://www.collegeboard.com/prod\\_downloads/yes/4297\\_MODULE\\_19.pdf](http://www.collegeboard.com/prod_downloads/yes/4297_MODULE_19.pdf) [2012, 9/13/2012].

Bedson, J. & Croft, P.R. 2008. "The discordance between clinical and radiographic knee osteoarthritis: A systematic search and summary of the literature", BMC Musculoskeletal Disorders, vol. 9, pp. 116-116.

Bedson, J., Mottram, S., Thomas, E. & Peat, G. 2007. "Knee pain and osteoarthritis in the general population: what influences patients to consult?", Family practice, vol. 24, no. 5, pp. 443-453.

Bellamy, N., Buchanan, W.W., Goldsmith, C.H., Campbell, J. & Stitt, L.W. 1988. "Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee", The Journal of rheumatology, vol. 15, no. 12, pp. 1833-1840.

Bellamy N. WOMAC Osteoarthritis Index User Guide. London Health Services. Centre; McMaster University: Ontario; 1996.

Bellamy N, Kirwan J, Boers M, Brooks P, Strand V, Tugwell P, Altman R, Brandt K, Dougados M, Lequesne M. Recommendations for a core set of outcome measures for future phase III clinical trials in knee, hip, and hand osteoarthritis. Consensus development at OMERACT III. J Rheumatol. 1997(a) Apr;24(4):799–802.

Bellamy, N., Campbell, J., Stevens, J., Pilch, L., Stewart, C. & Mahmood, Z. 1997(b). "Validation study of a computerized version of the Western Ontario and McMaster Universities VA3.0 Osteoarthritis Index", *The Journal of rheumatology*, vol. 24, no. 12, pp. 2413-2415.

Bellamy, N., Campbell, J., Haraoui, B., Buchbinder, R., Hobby, K., Roth, J.H. & MacDermid, J.C. 2002. "Dimensionality and clinical importance of pain and disability in hand osteoarthritis: development of the Australian/Canadian (AUSCAN) Osteoarthritis Hand Index", *OSTEOARTHRITIS AND CARTILAGE*, vol. 10, no. 11, pp. 855.

Benyon, K., Hill, S., Zadurian, N. & Mallen, C. 2010. "Coping strategies and self-efficacy as predictors of outcome in osteoarthritis: a systematic review", *Musculoskeletal care*.

Berger, B.D., Greenberg, D.A., Hodge, S.E. & Mendell, N.R. 1993. "The application of jackknife statistics to estimates of the recombination fraction", *Genetic epidemiology*, vol. 10, no. 6, pp. 471-476.

Bergner, M., Bobbitt, R.A., Carter, W.B. & Gilson, B.S. 1981. "The Sickness Impact Profile: development and final revision of a health status measure", *Medical care*, vol. 19, no. 8, pp. 787-805.

Berkman, L.F. & Syme, S.L. 1979. "Social networks, host resistance, and mortality: a nine-year follow-up study of Alameda County residents", *American Journal of Epidemiology*, vol. 109, no. 2, pp. 186-204.

Berkman, L.F. & Syme, S.L. 1994. "Social networks, host resistance, and mortality: A nine year follow-up study of Alameda County residents" in , eds. A. Steptoe & J. Wardle, Cambridge University Press, New York, NY US, pp. 43-67.

Berlau, D.J., Corrada, M.M., Peltz, C.B. & Kawas, C.H. 2012. "Disability in the oldest-old: Incidence and risk factors in The 90+ study", *The American Journal of Geriatric Psychiatry*, vol. 20, no. 2, pp. 159-168.

Bewick, V., Cheek, L. & Ball, J. 2005. "Statistics review 14: Logistic regression", *Critical Care (London, England)*, vol. 9, no. 1, pp. 112-118.

Bibb, S. 2007. "Clinical methods. Issues associated with secondary analysis of population health data", *Applied Nursing Research*, vol. 20, no. 2, pp. 94-99.

Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res.* 2002 Feb;52(2):69-77.

Bickenbach, J.E., Chatterji, S., Badley, E.M. & Ustun, T.B. 1999. "Models of disablement, universalism and the International Classification of Impairments, Disabilities and Handicaps", *Social science & medicine*, vol. 48, no. 9, pp. 1173-1187.

Blood, I. 2010 (a). Older people with high support needs: how can we empower them to enjoy a better life. Joseph Rowntree, York.

Blood, I. & Bamford, S.M. 2010 (b). Equality and Diversity and Older People with High Support Needs. Y, Joseph Rowntree Foundation, York.

Blyth,, F, M, March,, L, M, Brnabic,, A, J, Jorm,, L, R, Williamson, , M & Cousins,, M, J 2001. "Chronic pain in Australia: a prevalence study", *Pain* (03043959), vol. 89, no. 2-3, pp. 127-134.

Bone and Joint Decade. 1998. [Online]. Available:  
<http://www.boneandjointdecade.org> [2012, 9/12/2012].

Bonita, R., Beaglehole, T. & Kjellström, R. 2006. *Basic Epidemiology*, Second Edition, World Health Organisation.

Boorse, C. 1975. "On the Distinction between Disease and Illness", *Philosophy and Public Affairs*, vol. 5, no. 1 (Autumn, 1975), pp. 49-68.

Bornman, J. 2004. "The World Health Organisation's terminology and classification: application to severe disability ", *Disability & Rehabilitation*, vol. 26, no. 3, pp. 182 - 188.

Bowling, A. 2005. "Mode of questionnaire administration can have serious effects on data quality", *Journal of Public Health*, vol. 27, no. 3, pp. 281-291.



Bowling, A. (ed) 2002. Research methods in health: investigating health and health services, 2nd edn, Open University Press, Maidenhead, UK.

Bowling, A. 1997. Research methods in health: Investigating health and health services. Open University Press. Buckingham.

Busija L, Osborne RH, Nilsson A, Buchbinder R, Roos EM. Magnitude and meaningfulness of change in SF-36 scores in four types of orthopedic surgery. Health Qual Life Outcomes. 2008 Jul 31;6:55

Bradburn, M.J., Deeks, J.J., Berlin, J.A. & Russell Localio, A. 2007. "Much ado about nothing: a comparison of the performance of meta-analytical methods with rare events", Statistics in medicine, vol. 26, no. 1, pp. 53-77.

Bradley, P. & Burls, A.J. 1999. "Critical Appraisal Skills Programme: a project in critical appraisal skills teaching to improve the quality in health care", Journal of Clinical Governance, vol. 7, no. 2, pp. 88-91.

Brandt, E.N. & Pope, A.M. 1997. Enabling America: assessing the role of rehabilitation science and engineering National Academy Press, Washington, D.C.

Brandt, K.D., Dieppe, P. & Radin, E.L. 2008. "Etiopathogenesis of osteoarthritis", Rheumatic diseases clinics of North America, vol. 34, no. 3, pp. 531-559.

Brazier, J.E., Harper, R., Jones, N.M., O'Cathain, A., Thomas, K.J., Usherwood, T. & Westlake, L. 1992. "Validating the SF-36 health survey questionnaire: new outcome measure for primary care", BMJ (Clinical research ed.), vol. 305, no. 6846, pp. 160-164.

Brazier JE, Harper R, Munro J, Walters SJ, Snaith ML. Generic and condition-specific outcome measures for people with osteoarthritis of the knee. *Rheumatology (Oxford)*. 1999; 38(9):870-877.

Breivik, H., Collett, B., Ventafridda, V., Cohen, R. & Gallacher, D. 2006. "Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment", European journal of pain (London, England), vol. 10, no. 4, pp. 287-333.

- Breslin, F.C., Gnam, W., Franche, R.L., Mustard, C. & Lin, E. 2006. "Depression and activity limitations: examining gender differences in the general population", *Social psychiatry and psychiatric epidemiology*, vol. 41, no. 8, pp. 648-655.
- Brock, C. & Dost, A. 2005. *Self care a real choice*. Department of Health, London.
- Bruce, B. & Fries, J.F. 2003. "The Stanford Health Assessment Questionnaire: a review of its history, issues, progress, and documentation", *Journal of Rheumatology*, vol. 30, no. 1, pp. 167-178.
- Buchner, D.M. & Larson, E.B. 1988. "Transfer bias and the association of cognitive impairment with falls", *Journal of general internal medicine*, vol. 3, no. 3, pp. 254-259.
- Burholt, V. & Windle, G. 2006. *The Material Resources and Well-Being* [http://books.google.co.uk/books/about/The\\_Material\\_Resources\\_and\\_Well\\_Being\\_of.html?id=SsHkAQAACAAJ&redir\\_esc=y](http://books.google.co.uk/books/about/The_Material_Resources_and_Well_Being_of.html?id=SsHkAQAACAAJ&redir_esc=y) Joseph Rowntree Foundation.
- Burholt, V. & Windle, G. 2006. *Findings informing change: the material resources and well-being of older people*. Joseph Rowntree foundation, York.
- Burroughs, H., Lovell, K., Morley, M., Baldwin, R., Burns, A. & Chew-Graham, C. 2006. "'Justifiable depression': how primary care professionals and patients view late-life depression? A qualitative study", *Family practice*, vol. 23, no. 3, pp. 369-377.
- Bury, M. 2000. "A comment on the ICIDH2", *Disability and Society*, vol. 15, no. 7, pp. 1073.
- Cambridge CHPR 2010. Available: <http://www.jrf.org.uk/sites/files/jrf/supporting-older-people-summary.pdf> [2012, 9/16/2012].
- Cameron, J., Worrall-Carter, L., Page, K., Riegel, B., Lo, S.K. & Stewart, S. 2010. "Does cognitive impairment predict poor self-care in patients with heart failure?", *European journal of heart failure : journal of the Working Group on Heart Failure of the European Society of Cardiology*, vol. 12, no. 5, pp. 508-515.

Cardol, M., de Haan, R., de Jong, B., GAM & de Groot, I. 2001. "Psychometric properties of the Impact on Participation and Autonomy Questionnaire", Archives of Physical Medicine & Rehabilitation, vol. 82, no. 2, pp. 210-216.

Cardol, M., de Jong, B.A., van den Bos, G.A., Beelem, A., de Groot, I.J. & de Haan, R.J. 2002. "Beyond disability: perceived participation in people with a chronic disabling condition", Clinical rehabilitation, vol. 16, no. 1, pp. 27-35.

Cardol, M., De Jong, B.A. & Ward, C.D. 2002. "On autonomy and participation in rehabilitation", Disability and rehabilitation, vol. 24, no. 18, pp. 970-4; discussion 975-1004.

Cecchi, F., Molino-Lova, R., Di Iorio, A., Conti, A.A., Mannoni, A., Lauretani, F., Benvenuti, E., Bandinelli, S., Macchi, C. & Ferrucci, L. 2009. "Measures of physical performance capture the excess disability associated with hip pain or knee pain in older persons", Journals of Gerontology Series A: Biological Sciences & Medical Sciences, vol. 64A, no. 12, pp. 1316-1324.

Census data tables 2001. Available: <http://www.ons.gov.uk/ons/guide-method/census/2011/census-data/2001-census-data/2001-census-data-tables/index.html> [2012, 9/9/2012].

Çetin, N., Öztop, P., Bayramoglu, M., Cosar, S. & Özçürümez, G. 2009. "Relation between pain, disability and depression in patients with knee osteoarthritis [Turkish]", Turkish Journal of Rheumatology (Aves Yayincilik Ltd.STI), vol. 24, no. 4, pp. 196-201.

Chapireau, F. & Colvez, A. 1998. "Social disadvantage in the international classification of impairments, disabilities, and handicap", Social science & medicine (1982), vol. 47, no. 1, pp. 59-66.

Charles Christiansen, Bass-Haugen, J., Baum, C.M. & Julie Bass Haugen 2005. Occupational therapy: performance, participation, and well-being Slack, Thorofare, NJ.

Chase, H.W., Camille, N., Michael, A., Bullmore, E.T., Robbins, T.W. & Sahakian, B.J. 2010. "Regret and the negative evaluation of decision outcomes in major

depression ", Cognitive, affective & behavioral neuroscience, vol. 10, no. 3, pp. 406-413.

Chase, H.W., Camille, N., Michael, A., Bullmore, E.T., Robbins, T.W. & Sahakian, B.J. 2010. "Regret and the negative evaluation of decision outcomes in major depression", Cognitive, Affective & Behavioral Neuroscience, vol. 10, no. 3, pp. 406-413.

Cherubino, P., Sarzi-Puttini, P., Zuccaro, S.M. & Labianca, R. 2012. "The Management of Chronic Pain in Important Patient Subgroups", Clinical Drug Investigation, vol. 32, pp. 35-44.

Chew-Graham, C., Kovandžić, M., Gask, L., Burroughs, H., Clarke, P., Sanderson, H. & Dowrick, C. 2012. "Why may older people with depression not present to primary care? Messages from secondary analysis of qualitative data", Health & Social Care In The Community, vol. 20, no. 1, pp. 52-60.

Chubon, R.A. 1998. "Enabling America: Assessing the Role of Rehabilitation Science and Engineering", Journal of rehabilitation, vol. 64, no. 4, pp. 54-55.

Cieza, A., Bostan, C., Oberhauser, C. & Bickenbach, J.E. 2010. "Explaining functioning outcomes across musculoskeletal conditions: a multilevel modelling approach", Disability & Rehabilitation, vol. 32, pp. 85-S93.

Cigolle, C.T., Langa, K.M., Kabeto, M.U., Tian, Z. & Blaum, C.S. 2007. "Geriatric conditions and disability: the Health and Retirement Study", Annals of Internal Medicine, vol. 147, no. 3, pp. 156-164.

Clarke, A.K. 1987, "Community-based rehabilitation in England." International disability studies, vol. 9, no. 1, pp. 18-21.

Clarke, C. 2003. "Clinical application of the Canadian Model of Occupational Performance in a forensic rehabilitation hostel", British Journal of Occupational Therapy, vol. 66, no. 4, pp. 171-174.

Clarke, S.P. & Cossette, S. 2000. "Secondary analysis: theoretical, methodological, and practical considerations", Canadian Journal of Nursing Research, vol. 32, no. 3, pp. 109-129.

Cleves, M.A. 2002. "From the help desk: Comparing areas under receiver operating characteristic curves from two or more probit or logit models." The stata journal, vol. 2, no. 3, pp. 301-313.

Cohen-Mansfield, J. & Parpura-Gill, A. 2007. "Bathing: a framework for intervention focusing on psychosocial, architectural and human factors considerations", Archives of Gerontology and Geriatrics, vol. 45, no. 2, pp. 121-135.

Cole, J.C., Li, T., Lin, P., MacLean, R. & Wallenstein, G.V. 2008. "Treatment impact on estimated medical expenditure and job loss likelihood in rheumatoid arthritis: re-examining quality of life outcomes from a randomized placebo-controlled clinical trial with abatacept", Rheumatology (Oxford, England), vol. 47, no. 7, pp. 1044-1050.

Cole, M.B. & Tufano, R. 2008. Applied theories in occupational therapy: a practical approach, SLACK Incorporated, Thorofare, New Jersey.

Collin, C. 1990. "More on the Barthel Index", The American Journal Of Occupational Therapy.: Official Publication Of The American Occupational Therapy Association, vol. 44, no. 9, pp. 857-857.

Collin, C., Wade, D.T., Davies, S. & Horne, V. 1988. "The Barthel ADL index: a reliability study", International disability studies, vol. 10, no. 2, pp. 61-63.

Comprehensive Area Assessment - Audit Commission 2009. Available: <http://www.audit-commission.gov.uk/inspection-assessment/caa/Pages/Default.aspx> [2012, 9/17/2012].

Community care: detailed statistics on local authority personal social services for adults 2005. Department of Health, Government Statistical Service, England.

Conaghan, P. & Nelson, A.E. 2012. Osteoarthritis. Health Press Limited, Oxfordshire.

Cook, C., Pietrobon, R. & Hegedus, E. 2007. "Osteoarthritis and the impact on quality of life health indicators", Rheumatology international, vol. 27, no. 4, pp. 315-321.

Cook, N.R. 2008. "Statistical evaluation of prognostic versus diagnostic models: beyond the ROC curve", Clinical chemistry, vol. 54, no. 1, pp. 17-23.

Corti, M.C. & Rigon, C. 2003. "Epidemiology of osteoarthritis: prevalence, risk factors and functional impact", Aging clinical and experimental research, vol. 15, no. 5, pp. 359-363.

Covinsky, K.E., Lindquist, K., Dunlop, D.D., Gill, T.M. & Yelin, E. 2008. "Effect of arthritis in middle age on older-age functioning", Journal of the American Geriatrics Society, vol. 56, no. 1, pp. 23-28.

Covinsky, K.E., Lindquist, K., Dunlop, D.D. & Yelin, E. 2009. "Pain, Functional Limitations, and Aging", Journal of the American Geriatrics Society, vol. 57, no. 9, pp. 1556-1561.

Critical Appraisal Skills Programme. 2006. 10 questions to help you make sense of qualitative research 2006 [Homepage of Public Health Resource Unit] [Online]. Available: [http://www.sph.nhs.uk/sph-files/casp-appraisal tools/Qualitative%20Appraisal%20Tool.pdf](http://www.sph.nhs.uk/sph-files/casp-appraisal%20tools/Qualitative%20Appraisal%20Tool.pdf) [2012, September 4th.].

Croft, P., Jordan, K. & Jinks, C. 2005. "Pain elsewhere" and the impact of knee pain in older people", Arthritis & Rheumatism, vol. 52, no. 8, pp. 2350-2354.

Croft, P., Porcheret, M. & Peat, G. 2011. "Managing osteoarthritis in primary care: the GP as public health physician and surgical gatekeeper", The British Journal Of General Practice: The Journal Of The Royal College Of General Practitioners, vol. 61, no. 589, pp. 485-486.

College of Occupational Therapists 2002. [Online]. From Interface to Integration: A strategy for modernising Occupational Therapy services in local health and social

care communities Available:

<http://www.cot.co.uk/sites/default/files/publications/public/from-interface-to-integration.pdf> [2012, 9/9/2012].

College of Occupational Therapists 2004. Guidance-on-international-classification-functiong.pdf (application/pdf Object) [Online]. Available:

[http://www.cot.co.uk/sites/default/files/corporate\\_documents/public/guidance-on-international-classification-functiong.pdf](http://www.cot.co.uk/sites/default/files/corporate_documents/public/guidance-on-international-classification-functiong.pdf) [2012, 9/9/2012].

Couch A, Keniston K. Yeasayers and naysayers: agreeing response set as a personality variable. J Abnorm Soc Psychol. 1960 Mar;60:151-74.

Cugnet, P. Confidence Interval Estimation for Distribution Systems Power Consumption by using the Bootstrap Method. 1997 [Homepage of Digital Library and Archives. Virginia Tech], [Online]. Available:

<http://scholar.lib.vt.edu/theses/available/etd-61697-14555/> [2012, September 11].

Czaja, S.J., Weber, R.A. & Nair, S.N. 1993. "A human factors analysis of ADL activities: a capability-demand approach", Journal of gerontology, vol. 48 Spec No, pp. 44-48.

Dahaghin, S., Bierma-Zeinstra, S., Reijman, M., Pols, H.A.P. & Koes, B.W. 2005. "Prevalence and determinants of one month hand pain and hand related disability in the elderly (Rotterdam study). ", Annals of the Rheumatic Diseases, vol. 64, pp. 99.

Dahaghin, S., Bierma-Zeinstra, S., Ginai, A.Z., Pols, H.A.P., Hazes, J.M.W. & Koes, B.W. 2005. "Prevalence and pattern of radiographic hand osteoarthritis and association with pain and disability (the Rotterdam study)", Annals of the Rheumatic Diseases, vol. 64, no. 5, pp. 682-687.

Davies HTO, Crombie IK. What is a systematic review 2001. Available: <http://www.medicine.ox.ac.uk/bandolier/painres/download/whatis/syst-review.pdf> [2012, 9/13/2012].

Davis, A.M., Palaganas, M.P., Badley, E.M., Gladman, D.D., Inman, R.D. & Gignac, M.A. 2011. "Measuring participation in people with spondyloarthritis using

the social role participation questionnaire", *Annals of the Rheumatic Diseases*, vol. 70, no. 10, pp. 1765-1769.

Davis, M.A., Ettinger, W.H., Neuhaus, J.M. & Mallon, K.P. 1991. "Knee osteoarthritis and physical functioning: evidence from the NHANES I Epidemiologic Followup Study", *The Journal of rheumatology*, vol. 18, no. 4, pp. 591-598.

Davis, S.P. 2006. *Rehabilitation: the use of theories and models in practice* Churchill Livingstone, London.

Dawson, J., Linsell, L., Zondervan, K., Rose, P., Randall, T., Carr, A. & Fitzpatrick, R. 2004. "Epidemiology of hip and knee pain and its impact on overall health status in older adults", *Rheumatology (Oxford, England)*, vol. 43, no. 4, pp. 497-504.

Deal, M. 2003. Definition of impairment and disability. Available: [www.enham.org.uk/data/files/05\\_chapter\\_2.doc](http://www.enham.org.uk/data/files/05_chapter_2.doc) [2012, September 3rd].

Deeg, D.J.H. 2005. "Longitudinal characterization of course types of functional limitations", *Disability & Rehabilitation*, vol. 27, no. 5, pp. 253-261.

Denton, F.T. & Spencer, B.G. 2010. "Chronic health conditions: changing prevalence in an aging population and some implications for the delivery of health care services ", *Canadian journal on aging = La revue canadienne du vieillissement*, vol. 29, no. 1, pp. 11-21.

Department of Health 2010. *Health Profile: Stoke on Trent.*, Crown Copyright, London.

Department for Work and Pensions 2012. Available: <http://statistics.dwp.gov.uk/asd/index.php?page=hbai> [2012, 9/16/2012].

Department of Health (DoH) - Tackling Health Inequalities 2003. Available: [http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_4008268](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4008268) [2012, 9/17/2012].



- Dequeker, J. & Luyten, F.P. 2008. "The history of osteoarthritis-osteoarthrosis", *Annals of the Rheumatic Diseases*, vol. 67, no. 1, pp. 5-10.
- Desai, M.M., Lentzner, H.R. & Weeks, J.D. 2001. "Unmet need for personal assistance with activities of daily living among older adults", *Gerontologist*, vol. 41, no. 1, pp. 82-88.
- Dieppe, P. 1999. "Osteoarthritis: time to shift the paradigm. This includes distinguishing between severe disease and common minor disability", *BMJ (Clinical research ed.)*, vol. 318, no. 7194, pp. 1299-1300.
- Doblhammer, G. & Hoffmann, R. 2010. "Gender Differences in Trajectories of Health Limitations and Subsequent Mortality. A Study Based on the German Socioeconomic Panel 1995-2001 With a Mortality Follow-up 2002-2005", *Journals of Gerontology Series B: Psychological Sciences & Social Sciences*, vol. 65, no. 4, pp. 482-491.
- Dodge, H.H., Du, Y., Saxton, J.A. & Ganguli, M. 2006. "Cognitive domains and trajectories of functional independence in nondemented elderly persons", *Journals of Gerontology: Series A: Biological Sciences and Medical Sciences*, vol. 61A, no. 12, pp. 1330-1337.
- Doherty, M. 2001. "Risk factors for progression of knee osteoarthritis", *Lancet*, vol. 358, no. 9284, pp. 775-776.
- Donald, I.P. 1997. "Development of a modified Winchester disability scale--the elderly at risk rating scale", *Journal of epidemiology and community health*, vol. 51, no. 5, pp. 558-563.
- Donald, I.P. & Foy, C. 2004. "A longitudinal study of joint pain in older people", *Rheumatology*, vol. 43, no. 10, pp. 1256.
- Donati, A., Ruzzi, M., Adrario, E., Pelaia, P., Coluzzi, F., Gabbanelli, V. & Pietropaoli, P. 2004. "A new and feasible model for predicting operative risk", *British journal of anaesthesia*, vol. 93, no. 3, pp. 393-399.

Dong, X., Simon, M.A. & Evans, D.A. 2012. "Prevalence of self-neglect across gender, race, and socioeconomic status: findings from the Chicago Health and Aging Project", *Gerontology*, vol. 58, no. 3, pp. 258-268.

Dost, A. 2007. The evidence pack summary of work in progress., Department of Health, London.

Duncan, R., Peat, G., Thomas, E., Hay, E., McCall, I. & Croft, P. 2007. "Symptoms and radiographic osteoarthritis: not as discordant as they are made out to be?", *Annals of the Rheumatic Diseases*, vol. 66, no. 1, pp. 86-91.

Duncan, R., Peat, G., Thomas, E., Hay, E., McCall, I. & Croft, P. 2007. "Symptoms and radiographic osteoarthritis: not as discordant as they are made out to be? ", *Annals of the Rheumatic Diseases*, vol. 66, no. 1, pp. 86-91.

Dunlop, D.D., Hughes, S.L., Edelman, P., Singer, R.M. & Chang, R.W. 1998. "Impact of joint impairment on disability-specific domains at four years", *Journal of clinical epidemiology*, vol. 51, no. 12, pp. 1253-1261.

Dunlop, D.D., Hughes, S.L. & Manheim, L.M. 1997. "Disability in activities of daily living: patterns of change and a hierarchy of disability", *American Journal of Public Health*, vol. 87, no. 3, pp. 378-383.

Dunlop, D.D., Manheim, L.M., Song, J. & Chang, R.W. 2001. "Arthritis prevalence and activity limitations in older adults", *Arthritis & Rheumatism*, vol. 44, no. 1, pp. 212-221.

Dunnell, K. 2008. "Ageing and mortality in the UK--national statistician's annual article on the population", *Population trends*, no. 134, pp. 6-23.

Dziedzic, K., Thomas, E., Hill, S., Wilkie, R., Peat, G. & Croft, P.R. 2007 (a). "The impact of musculoskeletal hand problems in older adults: findings from the North Staffordshire Osteoarthritis Project (NorStOP)", *Rheumatology (Oxford, England)*, vol. 46, no. 6, pp. 963-967.

Dziedzic, K.S., Thomas, E., Myers, H., Hill, S. & Hay, E.M. 2007 (b). "The Australian/Canadian osteoarthritis hand index in a community-dwelling population

of older adults: reliability and validity", *Arthritis & Rheumatism: Arthritis Care & Research*, vol. 57, no. 3, pp. 423-428.

Ebrahim, S., Papacosta, O., Wannamethee, G., Adamson, J. & British Regional Heart Study 2004. "Social inequalities and disability in older men: prospective findings from the British regional heart study", *Social science & medicine* (1982), vol. 59, no. 10, pp. 2109-2120.

Efron, B. & Tibshirani, R. *An introduction to Bootstrap*, Chapman and Hall, London.

Emerson, E., Graham, H. & Hatton, C. 2006. "The Measurement of Poverty and Socioeconomic Position in Research Involving People with Intellectual Disability" in *International Review of Research in Mental Retardation*, ed. Laraine Masters Glidden, Academic Press, pp. 77-108.

Eng PM, Rimm EB, Fitzmaurice G, Kawachi I. Social ties and change in social ties in relation to subsequent total and cause-specific mortality and coronary heart disease incidence in men. *Am J Epidemiol*. 2002 Apr 15;155(8):700-9.

Engel, G.L. 1981. "The clinical application of the biopsychosocial model", *The Journal of medicine and philosophy*, vol. 6, no. 2, pp. 101-123.

Engel, G.L. 1980. "The clinical application of the biopsychosocial model", *The American Journal of Psychiatry*, vol. 137, no. 5, pp. 535-544.

Engel, G.L. 1992. "The need for a new medical model: A challenge for biomedicine", *Family Systems Medicine*, vol. 10, no. 3, pp. 317-331.

Engels, J.M. & Diehr, P. 2003. "Imputation of missing longitudinal data: a comparison of methods", *Journal of clinical epidemiology*, vol. 56, no. 10, pp. 968-976.

Escalante, A., Haas, R.W. & del Rincón, I. 2005. "A model of impairment and functional limitation in rheumatoid arthritis", *BMC Musculoskeletal Disorders*, vol. 6, pp. 16-16.

Estraviz, B., Quintana, J.M., Valdivieso, A., Bilbao, A., Padierna, A., de Urbina, J.O. & Sarabia, S. 2007. "Factors influencing change in health-related quality of life after liver transplantation", *Clinical transplantation*, vol. 21, no. 4, pp. 481-499.

ERPHO : View Resource : Data : Promoting Mental Health and Well-Being in Later Life . Available: <http://www.erpho.org.uk/viewResource.aspx?id=16172> [2012, 9/11/2012].

Etropolski, M. 2012. "Long term treatment of osteoarthritis pain: achieving a balance between efficacy and tolerability for a successful chronic therapy." in *Osteoarthritis - Diagnosis, treatment and surgery*. ed. Q. Chien, pp. 1.

EUMUSC 2011. Musculoskeletal health in Europe, Executive agency for health and consumers., Europe.

Experian report: Stoke-on-Trent is eighth poorest area in country | This is Staffordshire 2012. Available: <http://www.thisisstaffordshire.co.uk/Stoke-Trent-eighth-poorest-area-country/story-15421602-detail/story.html> [2012, 9/16/2012].

Fair access to care services - guidance on eligibility criteria for adult social care : Department of Health - Publications – 2002. [Homepage of Department of Health], [Online]. Available: [http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_4009653](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4009653) [2012, 9/11/2012].

Fearing, V.G., Law, M. & Clark, J. 1997. "An occupational performance process model: fostering client and therapist alliances", *Canadian Journal of Occupational Therapy*, vol. 64, no. 1, pp. 7-15.

Felson, D.T., Lawrence, R.C., Dieppe, P.A., Hirsch, R., Helmick, C.G., Jordan, J.M., Kington, R.S., Lane, N.E., Nevitt, M.C., Zhang, Y., Sowers, M., McAlindon, T., Spector, T.D., Poole, A.R., Yanovski, S.Z., Ateshian, G., Sharma, L., Buckwalter, J.A., Brandt, K.D. & Fries, J.F. 2000. "Osteoarthritis: new insights. Part 1: the disease and its risk factors", *Annals of Internal Medicine*, vol. 133, no. 8, pp. 635-646.

Felson, D.T., Lawrence, R.C., Hochberg, M.C., McAlindon, T., Dieppe, P.A., Minor, M.A., Blair, S.N., Berman, B.M., Fries, J.F., Weinberger, M., Lorig, K.R., Jacobs, J.J. & Goldberg, V. 2000. "Osteoarthritis: new insights. Part 2: treatment approaches", *Annals of Internal Medicine*, vol. 133, no. 9, pp. 726-737.

Felson, D.T., Niu, J., Clancy, M., Aliabadi, P., Sack, B., Guermazi, A., Hunter, D.J., Amin, S., Rogers, G. & Booth, S.L. 2007. "Low levels of vitamin D and worsening of knee osteoarthritis: results of two longitudinal studies", *Arthritis and Rheumatism*, vol. 56, no. 1, pp. 129-136.

Felson, D.T. & Chaisson, C.E. 1997. "2 Understanding the relationship between body weight and osteoarthritis ", *Baillière's Clinical Rheumatology*, vol. 11, no. 4, pp. 671 -681.

Fernandez-Lopez, J., Laffon, A., Blanco, F.J. & Carmona, L. 2008. "Prevalence, risk factors, and impact of knee pain suggesting osteoarthritis in Spain", *Clinical and experimental rheumatology*, vol. 26, no. 2, pp. 324-332.

Ferreira, V.M. & Sherman, A.M. 2007. "Relationship of optimism, pain and social support to well-being in older adults with osteoarthritis", *Aging and Mental Health*, vol. 11, no. 1, pp. 89-98.

Field, A. 2005. *Discovering statistics using SPSS (2nd ed.)* edn, Sage, London.

Field, A.P. 2009. *Discovering statistics using SPSS: (and sex and drugs and rock 'n' roll)* SAGE, Los Angeles.

Fishbain, D.A., Cutler, R., Rosomoff, H.L. & Rosomoff, R.S. 1997. "Chronic pain-associated depression: antecedent or consequence of chronic pain? A review", *The Clinical journal of pain*, vol. 13, no. 2, pp. 116-137.

Fishbain, D.A., Lewis, J., Cole, B., Cutler, B., Smets, E., Rosomoff, H. & Rosomoff, R.S. 2005. "Multidisciplinary pain facility treatment outcome for pain-associated fatigue", *Pain medicine (Malden, Mass.)*, vol. 6, no. 4, pp. 299-304.

Fitzpatrick, R., Davey, C., Buxton, M.J., Jones, D.R. Evaluating patient-based outcome measures for use in clinical trials. *Health Technology Assessment* 1998; Vol.2: No.14.

Forsberg, K. & Nilsson, B.E. 1992. "Coxarthrosis on the island of Gotland. Increased prevalence in a rural population", *Acta Orthopaedica Scandinavica*, vol. 63, no. 1, pp. 1-3.

Fougeyrollas, P. & Beauregard, L. 2001. The interactive person-environment in disability in creation, G. L. Albrecht, K. D. Seelman, & M. Bury (Eds.), *Handbook of disability studies*. (pp. 171-194) edn, Sage., Thousand Oaks, CA.

Francescutti, C., Frattura, L., Troiano, R., Gongolo, F., Martinuzzi, A., Sala, M., Meucci, P., Raggi, A., Russo, E., Buffoni, M., Gorini, G., Conclave, M., Petrangeli, A., Solipaca, A. & Leonardi, M. 2009. "Towards a common disability assessment framework: theoretical and methodological issues for providing public services and benefits using ICF", *Disability and rehabilitation*, vol. 31 Suppl 1, pp. S8-15.

Fransen, M., McConnell, S. & Bell, M. 2002. "Therapeutic exercise for people with osteoarthritis of the hip or knee. A systematic review", *The Journal of rheumatology*, vol. 29, no. 8, pp. 1737-1745.

Frazier, P.A., Tix, A.P. & Barron, K.E. 2004. "Testing Moderator and Mediator Effects in Counseling Psychology Research", *Journal of Counseling Psychology*, vol. 51, no. 1, pp. 115-134.

Fried, L.P., Ettinger, W.H., Lind, B., Newman, A.B. & Gardin, J. 1994. "Physical disability in older adults: A physiological approach", *Journal of clinical epidemiology*, vol. 47, no. 7, pp. 747-760.

Fries, J.F., Spitz, P., Kraines, R.G. & Holman, H.R. 1980. "Measurement of patient outcome in arthritis ", *Arthritis and Rheumatism*, vol. 23, no. 2, pp. 137-145.

Gabriel, S.E., Crowson, C.S. & O'Fallon, W.M. 1999. "Comorbidity in arthritis", *The Journal of rheumatology*, vol. 26, no. 11, pp. 2475-2479.

Gaidhane, A.M., Zahiruddin, Q.S., Waghmare, L., Zodpey, S., Goyal, R.C. & Johrapurkar, S.R. 2008. "Assessing self-care component of activities and participation domain of the international classification of functioning, disability and health (ICF) among people living with HIV/AIDS", *AIDS Care*, vol. 20, no. 9, pp. 1098-1104.

Gannon, M., Northern, J. & Carroll, S., 1971. "Characteristics of non-respondents among workers", *Journal of Applied Psychology* 55, 586- 588.

Garmon Bibb, S.,C. 2007. "Issues associated with secondary analysis of population health data", *Applied Nursing Research: ANR*, vol. 20, no. 2, pp. 94-99.

Garrow AP, Papageorgiou AC, Silman AJ, Thomas E, Jayson MIV, Macfarlane GJ: Development and validation of a questionnaire to assess disabling foot pain. *Pain* 2000, 85:107-113.

Garrow, A., P., Silman, A., J. & Macfarlane, G., J. 2004. "The Cheshire Foot Pain and Disability Survey: a population survey assessing prevalence and associations", *Pain* (03043959), vol. 110, no. 1-2, pp. 378-384.

Giacomini, M. 2004. "Interdisciplinarity in health services research: dreams and nightmares, maladies and remedies ", *Journal of health services research & policy*, vol. 9, no. 3, pp. 177-183.

Gignac, M.A.M., Cott, C. & Badley, E.M. 2000. "Adaptation to chronic illness and disability and its relationship to perceptions of independence and dependence", *Journals of Gerontology: Series B: Psychological Sciences and Social Sciences*, vol. 55B, no. 6, pp. P362-P372.

Gignac, M.A. 2000. "An evaluation of a psychotherapeutic group intervention for persons having difficulty coping with musculoskeletal disorders", *Social work in health care*, vol. 32, no. 1, pp. 57-75.

Gignac, M., Davis, A.M., Hawker, G., Wright, J.G., Mahomed, N., Fortin, P.R. & Badley, E.M. 2006. "'What do you expect? You're just getting older': a comparison of perceived osteoarthritis-related and aging-related health experiences in middle-

and older-age adults", *Arthritis & Rheumatism: Arthritis Care & Research*, vol. 55, no. 6, pp. 905-912.

Gill, T.M., Richardson, E.D. & Tinetti, M.E. 1995 (a). "Evaluating the risk of dependence in activities of daily living among community-living older adults with mild to moderate cognitive impairment", *J GERONTOL A BIOL SCI MED SCI*, vol. 50, no. 5, pp. M235.

Gill, T.M., Williams, C.S. & Tinetti, M.E. 1995 (b). "Assessing risk for the onset of functional dependence among older adults: the role of physical performance... presented in abstract form at the Annual Meeting of the American Geriatrics Society, Los Angeles, California, May 20, 1994", *Journal of the American Geriatrics Society*, vol. 43, no. 6, pp. 603-609.

Gill, T.M., Robison, J.T. & Tinetti, M.E. 1998. "Difficulty and dependence: two components of the disability continuum among community-living older persons", *Annals of Internal Medicine*, vol. 128, no. 2, pp. 96-101.

Gill, T.M., Hardy, S.E. & Williams, C.S. 2002 (a). "Underestimation of disability in community-living older persons", *Journal of the American Geriatrics Society*, vol. 50, no. 9, pp. 1492-1497.

Gill, T.M., Baker, D.I., Gottschalk, M., Peduzzi, P.N., Allore, H. & Byers, A. 2002 (b). "A program to prevent functional decline in physically frail, elderly persons who live at home", *The New England journal of medicine*, vol. 347, no. 14, pp. 1068-1074.

Gill, T.M., Allore, H. & Guo, Z. 2003. "Restricted activity and functional decline among community-living older persons", *Archives of Internal Medicine*, vol. 163, no. 11, pp. 1317-1322.

Gill, T.M., Allore, H.G., Holford, T.R. & Guo, Z. 2004 (a). "Hospitalization, restricted activity, and the development of disability among older persons", *JAMA: Journal of the American Medical Association*, vol. 292, no. 17, pp. 2115-2124.



- Gill, T.M., Allore, H., Holford, T.R. & Guo, Z. 2004 (b). "The development of insidious disability in activities of daily living among community-living older persons", *The American Journal of Medicine*, vol. 117, no. 7, pp. 484-491.
- Gill, T.M., Guo, Z. & Allore, H.G. 2006 (a). "The epidemiology of bathing disability in older persons", *Journal of the American Geriatrics Society*, vol. 54, no. 10, pp. 1524-1530.
- Gill, T.M., Allore, H.G. & Han, L. 2006 (b). "Bathing disability and the risk of long-term admission to a nursing home", *The Journals Of Gerontology.Series A, Biological Sciences And Medical Sciences*, vol. 61, no. 8, pp. 821-825.
- Gill, T.M., Han, L. & Allore, H.G. 2007 (a). "Predisposing factors and precipitants for bathing disability in older persons", *Journal of the American Geriatrics Society*, vol. 55, no. 4, pp. 534-540.
- Gill, T.M., Han, L. & Allore, H.G. 2007 (b). "Bath aids and the subsequent development of bathing disability in community-living older persons", *Journal of the American Geriatrics Society*, vol. 55, no. 11, pp. 1757-1763.
- Gill, T.M. & Gahbauer, E.A. 2008. "Evaluating disability over discrete periods of time", *The Journals Of Gerontology.Series A, Biological Sciences And Medical Sciences*, vol. 63, no. 6, pp. 588-594.
- Gill, T.M., Murphy, T.E., Barry, L.C. & Allore, H.G. 2009. "Risk factors for disability subtypes in older persons", *Journal of the American Geriatrics Society*, vol. 57, no. 10, pp. 1850-1855.
- Gill, T.M., Allore, H.G., Gahbauer, E.A. & Murphy, T.E. 2010. "Change in disability after hospitalization or restricted activity in older persons", *JAMA: The Journal Of The American Medical Association*, vol. 304, no. 17, pp. 1919-1928.
- Gill, T.M. 2010. "Assessment of function and disability in longitudinal studies", *Journal of the American Geriatrics Society*, vol. 58 Suppl 2, pp. S308-12.

Graham, J.W., Olchowski, A.E. & Gilreath, T.D. 2007. "How many imputations are really needed? Some practical clarifications of multiple imputation theory", *Prevention Science*, vol. 8, no. 3, pp. 206-213.

Greenhill, Y., Wilkie, R., Beech, R. & Peat, G. 2011. Self-care needs in community-dwelling older adults with joint pain: A descriptive cross-sectional study, *Ann Rheum Dis* 2011;70(Suppl3):698.

Grimby, G. & Smedby, B. 2001. "ICF approved as the successor of ICIDH", *Journal of Rehabilitation Medicine* (Taylor & Francis Ltd), vol. 33, no. 5, pp. 193-194.

Grotle, M., Hagen, K.B., Natvig, B., Dahl, F.A. & Kvien, T.K. 2008. "Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological study in the general population with 10 years follow-up", *BMC Musculoskeletal Disorders*, vol. 9, pp. 132-132.

Grundy, E. 2000. "Socio-demographic differences in the onset and progression of disability in early old age: a longitudinal study ", *Age and Ageing*, vol. 29, no. 2, pp. 149 - 157.

Guo, Z., Gill, T.M. & Allore, H.G. 2008. "Modeling repeated time-to-event health conditions with discontinuous risk intervals. An example of a longitudinal study of functional disability among older persons", *Methods of information in medicine*, vol. 47, no. 2, pp. 107-116.

Guralnik, J.M. 1995. "Lower extremity function in persons over the age of 70 years as a predictor of subsequent disability", *New England Journal of Medicine*, vol. 332, no. 9, pp. 556-561.

Gureje, O., Simon, G, E & Von Korff, M 2001. "A cross-national study of the course of persistent pain in primary care", *Pain* (03043959), vol. 92, no. 1-2, pp. 195-200.

Gureje, O., Ademola, A. & Olley, B.O. 2008. "Depression and disability: comparisons with common physical conditions in the Ibadan Study of Aging", *Journal of the American Geriatrics Society*, vol. 56, no. 11, pp. 2033-2038.

- Haglund, L. & Kjellberg, A. 1999. "A critical analysis of the model of human occupation", *Canadian Journal of Occupational Therapy*, vol. 66, no. 2, pp. 102-108.
- Handley, T.E., Inder, K.J., Kelly, B.J., Attia, J.R., Lewin, T.J., Fitzgerald, M.N. & Kay-Lambkin, F. 2012. "You've got to have friends: the predictive value of social integration and support in suicidal ideation among rural communities", *Social psychiatry and psychiatric epidemiology*, vol. 47, no. 8, pp. 1281-1290.
- Hardy, S.E. & Gill, T.M. 2004. "Recovery from disability among community-dwelling older persons", *JAMA: Journal of the American Medical Association*, vol. 291, no. 13, pp. 1596-1602.
- Hardy, S.E., Dubin, J.A., Holford, T.R. & Gill, T.M. 2005. "Transitions between states of disability and independence among older persons", *American Journal of Epidemiology*, vol. 161, no. 6, pp. 575-584.
- Hardy, S.E. & Gill, T.M. 2005. "Factors associated with recovery of independence among newly disabled older persons", *Archives of Internal Medicine*, vol. 165, no. 1, pp. 106-112.
- Harrell, F.E.J. 2001. Model uncertainty, penalization and parsimony [Homepage of University of Virginia], [Online]. Available: <http://biostat.mc.vanderbilt.edu/wiki/pub/Main/FHHandouts/iscb98.pdf> [2012, September 3rd].
- Harrell, F.E., Jr, Lee, K.L. & Mark, D.B. 1996. "Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors ", *Statistics in medicine*, vol. 15, no. 4, pp. 361-387.
- Harwood,, R, H, Jitapunkul, , S, Dickinson, , E & Ebrahim, , S 1994. "Measuring handicap: motives, methods, and a model", *Quality in Health Care*, vol. 3, no. 1, pp. 53-57.
- Haslam, D. & Cook, D. 2011. "Obesity in the elderly patient." *Midlife and beyond*, , pp. 27.

Hawker, G.A., Gignac, M.A.M., Badley, E., Davis, A.M., French, M.R., Li, Y., Perruccio, A.V., Power, J.D., Sale, J. & Lou, W. 2011. "A longitudinal study to explain the pain-depression link in older adults with osteoarthritis", *Arthritis Care & Research*, vol. 63, no. 10, pp. 1382-1390.

Hayden, J.A., Côté, P., Steenstra, I.A. & Bombardier, C. 2008. "Identifying phases of investigation helps planning, appraising, and applying the results of explanatory prognosis studies", *Journal of clinical epidemiology*, vol. 61, no. 6, pp. 552-560.

Hayden, J.A. 2007. *Methodological issues in systematic reviews of prognosis and prognostic factors: low back pain*, University of Toronto: Canada.

Hayden, J.A., Côté, P. & Bombardier, C. 2006. "Academia and clinic. Evaluation of the quality of prognosis studies in systematic reviews", *Annals of Internal Medicine*, vol. 144, no. 6, pp. 427-437.

Health survey for England 2005. *Chronic diseases: the health of older people.*, The Information Centre, Leeds.

Heerkens, Y.F., Brandsma, J.W., Lakerveld-Heyl, K. & van Ravensberg, C. 1994. "Impairments and disabilities -- the difference: proposal for adjustment of the International Classification of Impairments, Disabilities, and Handicaps", *Physical Therapy*, vol. 74, no. 5, pp. 430-442.

Hemingway, H., Stafford, M. , Stansfeld, S., Shipley, M. , and Marmot, M. Is the SF-36 a valid measure of change in population health? Results from the Whitehall II Study. *BMJ*. 1997 November 15; 315(7118): 1273–1279.

Hemmingsson, H. & Jonsson, H. 2005. "An occupational perspective on the concept of participation in the International Classification of Functioning, Disability and Health--some critical remarks", *AMERICAN JOURNAL OF OCCUPATIONAL THERAPY*, vol. 59, no. 5, pp. 569.

Hennekens, C. & Buring, J.E. 1987. "Need for large sample sizes in randomized trials", *Pediatrics*, vol. 79, no. 4, pp. 569-571.

Hennekens, C.H., Buring, J.E. & Hebert, P.R. 1987. "Implications of overviews of randomized trials", *Statistics in medicine*, vol. 6, no. 3, pp. 397-409.

Herrmann, C. 1997. "International experiences with the Hospital Anxiety and Depression Scale--a review of validation data and clinical results", *Journal of psychosomatic research*, vol. 42, no. 1, pp. 17-41.

Hochberg, M.C., Kasper, J., Williamson, J., Skinner, A. & Fried, L.P. 1995. "The contribution of osteoarthritis to disability: preliminary data from the Women's Health and Aging Study", *The Journal Of Rheumatology. Supplement*, vol. 43, pp. 16-18.

Hochberg, M.C., Vignon, E. & Maheu, E. 2000. Session 2: Clinical assessment of hand OA. *Osteoarthritis Cartilage*, vol. 8, pp. 38.

Households Below Average Income (HBAI) 2010/2011. Available: <http://statistics.dwp.gov.uk/asd/index.php?page=hbai> [2012, 9/16/2012].

Hosie, G. & Dickson, J. 2000. *Managing osteoarthritis in primary care*. Blackwell Publishing., Edinburgh.

Hosmer, D. & Lemeshow, S. 1989. *Applied Logistic Regression*. John Wiley & Sons, New York.

Hosmer, D.W. & Lemeshow, S. 1995. "Confidence interval estimates of an index of quality performance based on logistic regression models", *Statistics in medicine*, vol. 14, no. 19, pp. 2161-2172.

Howse, K. 2006. *Increasing life expectancy and the compression of morbidity: A critical review of the debate.*, Oxford Institute of ageing working papers., Oxford.

Hughes, S.L. & Dunlop, D. 1995. "The prevalence and impact of arthritis in older persons", *Arthritis Care And Research: The Official Journal Of The Arthritis Health Professions Association*, vol. 8, no. 4, pp. 257-264.

Hung, W.W., Ross, J.S., Boockvar, K.S. & Siu, A.L. 2012. "Association of chronic diseases and impairments with disability in older adults: a decade of change?" *Medical care*, vol. 50, no. 6, pp. 501-507.

Huwiler-Muntener, K., Juni, P., Junker, C. & Egger, M. 2002. "Quality of reporting of randomized trials as a measure of methodologic quality ", JAMA : the journal of the American Medical Association, vol. 287, no. 21, pp. 2801-2804.

Hybels, C.F., Pieper, C.F., Blazer, D.G., Fillenbaum, G.G. & Steffens, D.C. 2010. "Trajectories of mobility and IADL function in older patients diagnosed with major depression", International journal of geriatric psychiatry, vol. 25, no. 1, pp. 74-81.

Imrie, R. 2004. "Demystifying disability: a review of the International Classification of Functioning, Disability and Health ", Sociology of health & illness, vol. 26, no. 3, pp. 287-305.

Indices of deprivation 2010. Communities and neighbourhoods. Department for Communities and Local Government . Available:

<http://www.communities.gov.uk/communities/research/indicesdeprivation/deprivation10/> [2012, 9/17/2012].

Institute of Medicine, National Research Council, Wunderlich, G.S., Rice, D.P. & Nicole L. Amado 2002. The dynamics of disability measuring and monitoring disability for social security programs National Academy Press, Washington, D.C.

J Fang , Activity limitation and personal care participation restriction of older inpatients in an acute care setting. 2010. [Homepage of La Trobe University], [Online]. Available:

[http://www.hkiot.org/2012otc/docs/pdf/3S1.1\\_Ms%20NEO%20Cui%20Fang%20Josephine.pdf](http://www.hkiot.org/2012otc/docs/pdf/3S1.1_Ms%20NEO%20Cui%20Fang%20Josephine.pdf) [2012, September/Monday].

Jacobsen, S., Sonne-Holm, S., Soballe, K., Gebuhr, P. & Lund, B. 2004. "Radiographic case definitions and prevalence of osteoarthritis of the hip: a survey of 4 151 subjects in the Osteoarthritis Substudy of the Copenhagen City Heart Study ", Acta Orthopaedica Scandinavica, vol. 75, no. 6, pp. 713-720.

Jagger, C., Spiers, N. & Arthur, A. 2005. "The role of sensory and cognitive function in the onset of activity restriction in older people", Disability and rehabilitation, vol. 27, no. 5, pp. 277-283.

Jagger, C., Arthur, A.J., Spiers, N.A. & Clarke, M. 2001. "Patterns of onset of disability in activities of daily living with age", *Journal of the American Geriatrics Society*, vol. 49, no. 4, pp. 404-409.

Jenkinson, C. & Layte, R. 1997. "Development and testing of the UK SF-12 (short form health survey)", *Journal of health services research & policy*, vol. 2, no. 1, pp. 14-18.

Jenkinson, C., Layte, R., Coulter, A. & Wright, L. 1996. "Evidence for the sensitivity of the SF-36 health status measure to inequalities in health: results from the Oxford healthy lifestyles survey", *Journal of epidemiology and community health*, vol. 50, no. 3, pp. 377-380.

Jette, A.M. 1994. "How measurement techniques influence estimates of disability in older populations", *Social science & medicine*, vol. 38, no. 7, pp. 937-942.

Jette, A.M. 2006. "Toward a Common Language for Function, Disability, and Health", *Physical Therapy*, vol. 86, no. 5, pp. 726-734.

Jette, A.M., Haley, S.M. & Kooyoomjian, J.T. 2003. "Are the Icf Activity and Participation Dimensions Distinct?", *Journal of Rehabilitation Medicine* (Taylor & Francis Ltd), vol. 35, no. 3, pp. 145.

Jewell, P. N. & Hubbard, A. 2006. *Analysis of longitudinal studies in epidemiology*, University of California, University of California.

Jinks, C., Jordan, K., Ong, B.N. & Croft, P. 2004. "A brief screening tool for knee pain in primary care (KNEST). 2. Results from a survey in the general population aged 50 and over", *Rheumatology* (Oxford, England), vol. 43, no. 1, pp. 55-61.

Jinks, C., Lewis, M., Ong, B.N. & Croft, P. 2001. "A brief screening tool for knee pain in primary care. 1. Validity and reliability", *Rheumatology*, vol. 40, no. 5, pp. 528.

Jinks, C., Jordan, K. & Croft, P. 2003. "Evaluation of a computer-assisted data entry procedure (including Teleform) for large-scale mailed surveys", *Computers in biology and medicine*, vol. 33, no. 5, pp. 425-437.

Jinks, C., Jordan, K. & Croft, P. 2002. "Measuring the population impact of knee pain and disability with the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)", *Pain* (03043959), vol. 100, no. 1-2, pp. 55-64.

Jordan, K., Ong, B.N. & Croft, P.R. 1998. *Mastering Statistics - A Guide for Health Service Professionals and Researchers*: Amazon.co.uk: Nelson Thornes.

Jordan, K.M., Sawyer, S., Coakley, P., Smith, H.E., Cooper, C. & Arden, N.K. 2004. "The use of conventional and complementary treatments for knee osteoarthritis in the community", *Rheumatology* (Oxford, England), vol. 43, no. 3, pp. 381-384.

Jordan, K.P., Wilkie, R., Muller, S., Myers, H. & Nicholls, E. 2009. "Measurement of change in function and disability in osteoarthritis: current approaches and future challenges", *Current opinion in rheumatology*, vol. 21, no. 5, pp. 525-530.

Jordan, K., P., Thomas, E., Peat, G., Wilkie, R. & Croft, P. 2008. "Social risks for disabling pain in older people: a prospective study of individual and area characteristics", *Pain* (03043959), vol. 137, no. 3, pp. 652-661.

Joss, M. 2002. "Occupational therapy and rehabilitation for work", *British Journal of Occupational Therapy*, vol. 65, no. 3, pp. 141-148.

Kaplan, M.S., Huguet, N., Newsom, J.T. & McFarland, B.H. 2003. "Characteristics of physically inactive older adults with arthritis: results of a population-based study", *Preventive medicine*, vol. 37, no. 1, pp. 61-67.

Katz, S. & Akpom, C.A. 1976. "Index of ADL", *Medical care*, vol. 14, no. 5, pp. 116-118.

Katz, S., Downs, T.D., Cash, H.R. & Grotz, R.C. 1970. "Progress in development of the index of ADL", *The Gerontologist*, vol. 10, no. 1, pp. 20-30.

Kelley, K., Clark, B., Brown, V. & Sitzia, J. 2003. "Good practice in the conduct and reporting of survey research", *International Journal for Quality in Health Care*, vol. 15, no. 3, pp. 261-266.



Kempen, G.I. & Suurmeijer, T.P. 1990. "The development of a hierarchical polychotomous ADL-IADL scale for noninstitutionalized elders" *The Gerontologist*, vol. 30, no. 4, pp. 497-502.

Kempen, G.I.J.M., Miedema, I., Ormel, J. & Molenaar, W. 1996, "The assessment of disability with the Groningen Activity Restriction Scale. Conceptual framework and psychometric properties ", *Social science & medicine*, vol. 43, no. 11, pp. 1601 - 1610.

Kempen, G. & Suurmeijer, P. 1991. "Factors influencing professional home care utilization among the elderly", *Social science & medicine*, vol. 32, no. 1, pp. 77-81.

Kennedy, J., LaPlante, M.P. & Kaye, H.S. 1997. Need for assistance in the activities of daily living, United States Department of Education. National Institute on Disability and Rehabilitation Research, Washington, DC.

Kennedy, J. 2001. "Unmet and undermet need for activities of daily living and instrumental activities of daily living assistance among adults with disabilities", *Medical care*, vol. 39, no. 12, pp. 1305-1312.

Kielhofner, G. & Kielhofner, G. 2002. *A Model of Human Occupation: theory and application* Lippincott Williams & Wilkins, Baltimore, MD.

Kielhofner, G. 2008. *A Model of Human Occupation: theory and application* Lippincott Williams & Wilkins, Baltimore, MD.

Kingston, A., Collerton, J., Davies, K., Bond, J., Robinson, L. & Jagger, C. 2012. "Losing the ability in activities of daily living in the oldest old: a hierarchic disability scale from the Newcastle 85+ study", *Plos One*, vol. 7, no. 2, pp. e31665-e31665.

Kjeken, I., Dagfinrud, H., Slatkowsky-Christensen, B., Mowinckel, P., Uhlig, T., Kvien, T.K. & Finset, A. 2005. "Activity limitations and participation restrictions in women with hand osteoarthritis: patients' descriptions and associations between dimensions of functioning", *Annals of the Rheumatic Diseases*, vol. 64, no. 11, pp. 1633-1638.

- Klebanoff, M.A. & Cole, S.R. 2008. "Use of multiple imputation in the epidemiologic literature", *American Journal of Epidemiology*, vol. 168, no. 4, pp. 355-357.
- Klenk, J., Rapp, K., Buchele, G., Keil, U. & Weiland, S.K. 2007. "Increasing life expectancy in Germany: quantitative contributions from changes in age- and disease-specific mortality", *European journal of public health*, vol. 17, no. 6, pp. 587-592.
- Klijs, B., Nusselder, W.J., Looman, C.W. & Mackenbach, J.P. 2011. "Contribution of chronic disease to the burden of disability", *Plos One*, vol. 6, no. 9.
- Kloppenburg M, Stamm T, Watt I, Kainberger F, Cawston TE, Birrell FN, Petersson IF, Saxne T, Kvien TK, Slatkowsky-Christensen B, Dougados M, Gossec L, Breedveld FC, Smolen JS. Research in hand osteoarthritis: time for reappraisal and demand for new strategies. An opinion paper. *Ann Rheum Dis*. 2007;66:1157–1161. doi: 10.1136/ard.2007.070813.
- Knottnerus, J.A., van Weel, C. & Muris, J.W.M. 2002. "Evaluation of diagnostic procedures", *BMJ (Clinical research ed.)*, vol. 324, no. 7335, pp. 477-480.
- Kostanjsek, N. & Üstün, T.B. 2010. "The International Classification of Functioning, Disability and Health (ICF): Structure, concept, use and linkages with the International Classification of Disease (ICD)", *Die Psychiatrie: Grundlagen & Perspektiven*, vol. 7, no. 3, pp. 170-178.
- Lacey RJ, Lewis M, Jordan K, Jinks C, Sim J. Interrater reliability of scoring of pain drawings in a self-report health survey. *Spine (Phila Pa 1976)*. 2005 Aug 15;30(16):E455-8.
- Lane, N.E. & Wallace, D.J. 2002. *All About Osteoarthritis: The Definitive Resource for Arthritis Patients and Their Families*, Oxford University Press, Inc., New York 10016.
- Lang, T.A. & Secic, M. 1997. *How to report statistics in medicine*, American College of Physicians, Philadelphia.

Laub, C. & Kuhl, T. 2005. How bad is good? : A Critical Look at the Fitting of Reflectivity Models using the Reduced Chi-Square Statistic. Available: [http://neutrons.ornl.gov/workshops/sns\\_hfir\\_users/posters/Laub\\_Chi-Square\\_Data\\_Fitting.pdf](http://neutrons.ornl.gov/workshops/sns_hfir_users/posters/Laub_Chi-Square_Data_Fitting.pdf) [2012, 9/17/2012].

Laukkanen, P., Era, P., Heikkinen, R.L., Suutama, T., Kauppinen, M. & Heikkinen, E. 1994. "Factors related to carrying out everyday activities among elderly people aged 80", Aging (Milan, Italy), vol. 6, no. 6, pp. 433-443.

Law, M., Baum, C.M. & Dunn, W. 2001. Measuring occupational performance: Supporting best practice in occupational therapy, Thorofare, Slack Inc.

Lawton, M.P. 1990. "Aging and performance of home tasks", Human factors, vol. 32, no. 5, pp. 527-536.

Lawton, M.P. & Brody, E.M. 1969. "Assessment of older people: Self-maintaining and instrumental activities of daily living", Gerontologist, vol. 9, pp. 179.

Leon, D.A. 2007. "Commentary: Preston and mortality trends since the mid-1970s " International journal of epidemiology, vol. 36, no. 3, pp. 500 - 501.

Leroux, I., Dionne, C.E., Bourbonnais, R. & Brisson, C. 2005. "Prevalence of musculoskeletal activity limitation and associated factors among adults in the general population in the 1998 Quebec Health Survey", The Journal of rheumatology, vol. 32, no. 9, pp. 1794-1804.

Leung, Y.Y. & Thumboo, J. 2011. "An overview of OA research in two urban APLAR populations", International Journal Of Rheumatic Diseases, vol. 14, no. 2, pp. 130-135.

Leveille, S.G., Ling, S., Hochberg, M.C., Resnick, H.E., Bandeen-Roche, K.J., Won, A. & Guralnik, J.M. 2001. "Widespread musculoskeletal pain and the progression of disability in older disabled women", Annals of Internal Medicine, vol. 135, no. 12, pp. 1038-1046.

- Lewis, M., Jordan, K., Lacey, R.J., Jinks, C. & Sim, J. 2002. "Inter-rater reliability assessment of the scoring of the body pain manikin", *Rheumatology*, vol. 2002;41:52.
- Lin, E., Katon, W., Von Korff, M., Tang, L., Williams, J., J., Kroenke, K., Hunkeler, E., Harpole, L., Hegel, M., Arian, P., Hoffing, M., Della Penna, R., Langston, C. & Unützer, J. 2003. "Effect of improving depression care on pain and functional outcomes among older adults with arthritis: a randomized controlled trial", *JAMA: Journal of the American Medical Association*, vol. 290, no. 18, pp. 2428-2434.
- Lisspers, J., Nygren, A. & Söderman, E. 1997. "Hospital Anxiety and Depression Scale (HAD): some psychometric data for a Swedish sample", *Acta Psychiatrica Scandinavica*, vol. 96, no. 4, pp. 281-286.
- Litwin, H. & Sapir, E.V. 2009. "Perceived income adequacy among older adults in 12 countries: findings from the survey of health, ageing, and retirement in Europe", *Gerontologist*, vol. 49, no. 3, pp. 397-406.
- Lloyd, C. & King, R. 2001. "Work-related stress and occupational therapy", *Occupational Therapy International*, vol. 8, no. 4, pp. 227-243.
- Lohr, K.N., Aaronson, N.K., Alonso, J., Burnam, M.A., Patrick, D.L., Perrin, E.B. & Roberts, J.S. 1996. "Evaluating quality-of-life and health status instruments: development of scientific review criteria", *Clinical therapeutics*, vol. 18, no. 5, pp. 979-992.
- Machado, G.P., Gignac, M.A. & Badley, E.M. 2008. "Participation restrictions among older adults with osteoarthritis: A mediated model of physical symptoms, activity limitations, and depression", *Arthritis & Rheumatism: Arthritis Care & Research*, vol. 59, no. 1, pp. 129-135.
- Madden, R., Madden, R., Choi, C., Tallis, K. & Xingyan. 2005. Use of ICF in health information systems and surveys. [Homepage of WHO-FIC], [Online]. Available: [http://apps.who.int/classifications/apps/icd/meetings/tokyomeeting/B\\_5-6%20Use%20of%20ICF%20in%20health%20information%20systems%20and%20surveys.pdf](http://apps.who.int/classifications/apps/icd/meetings/tokyomeeting/B_5-6%20Use%20of%20ICF%20in%20health%20information%20systems%20and%20surveys.pdf) [2012, September 3rd].

Magasi, S. & Post, M.W. 2010. "A Comparative Review of Contemporary Participation Measures' Psychometric Properties and Content Coverage", Archives of Physical Medicine & Rehabilitation, vol. 91, no. 9, pp. 17-28.

Magni, G., Caldieron, C., Rigatti-Luchini, S. & Merskey, H. 1990, "Chronic musculoskeletal pain and depressive symptoms in the general population. An analysis of the 1st National Health and Nutrition Examination Survey data", Pain, vol. 43, no. 3, pp. 299-307.

Magni, G., Marchetti, M., Moreschi, C., Merskey, H. & Luchini, S.R. 1993."Chronic musculoskeletal pain and depressive symptoms in the National Health and Nutrition Examination. I. Epidemiologic follow-up study", Pain, vol. 53, no. 2, pp. 163-168.

Mallen, C.D., Peat, G., Thomas, E., Dunn, K.M. & Croft, P.R. 2007. "Prognostic factors for musculoskeletal pain in primary care: a systematic review", British Journal of General Practice, vol. 57, no. 541, pp. 655-661.

Mallen, C.D., Peat, G., Thomas, E., Wathall, S., Whitehurst, T., Clements, C., Bailey, J., Gray, J. & Croft, P.R. 2006. "The assessment of the prognosis of musculoskeletal conditions in older adults presenting to general practice: a research protocol ", BMC musculoskeletal disorders, vol. 7, pp. 84.

Mallen, C., Peat, G., Thomas, E. & Croft, P. 2005. "Severely disabling chronic pain in young adults: prevalence from a population-based postal survey in North Staffordshire", BMC Musculoskeletal Disorders, vol. 6, pp. 42-42.

Mallinson, S 1998. "The Short-Form 36 and older people: some problems encountered when using postal administration", Journal of Epidemiology & Community Health, vol. 52, no. 5, pp. 324-328.

Mallinson S. Listening to respondents: a qualitative assessment of the Short-Form 36 Health Status Questionnaire. Soc Sci Med. 2002 Jan;54(1):11-21.

Malta, M., Cardoso, L.O., Bastos, F.I., Magnanini, M.M.F. & Silva, Cosme Marcelo Furtado Passos da 2010. "STROBE initiative: guidelines on reporting observational studies", Revista De Saúde Pública, vol. 44, no. 3, pp. 559-565.

- Mangione, T.W. 1995. Mail surveys: improving the quality Sage Publications, Thousand Oaks, Calif.
- Mann, C.J. 2003. "Observational research methods. Research design II: cohort, cross sectional, and case-control studies", *Emergency Medicine Journal*, vol. 20, no. 1, pp. 54-60.
- Mannoni, A., Briganti, M.P., Di Bari, M., Ferrucci, L., Costanzo, S., Serni, U., Masotti, G. & Marchionni, N. 2003. "Epidemiological profile of symptomatic osteoarthritis in older adults: a population based study in Dicomano, Italy", *Annals of the Rheumatic Diseases*, vol. 62, no. 6, pp. 576-578.
- Mannoni, A., Briganti, M.P., Di Bari, M., Ferrucci, L., Serni, U., Masotti, G. & Marchionni, N. 2000. "Prevalence of symptomatic hand osteoarthritis in community-dwelling older persons: the ICARe Dicomano study. Insufficienza Cardiaca negli Anziani Residenti a Dicomano", *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society*, vol. 8 Suppl A, pp. S11-S13.
- Manthorpe, J. & Liffé, S. 2010. "Suicide in later life: public health and practitioner perspectives", *International journal of geriatric psychiatry*, vol. 25, no. 12, pp. 1230-1238.
- Margolis RB, Tait RC, Krause SJ. A rating system for use with patient pain drawings. *Pain*. 1986 Jan;24(1):57-65.
- Martin, L.G., Freedman, V.A., Schoeni, R.F. & Andreski, P.M. 2010. "Trends In Disability And Related Chronic Conditions Among People Ages Fifty To Sixty-Four", *Health affairs*, vol. 29, no. 4, pp. 725-731.
- Masala, C. & Petretto, D.R. 2008. "From disablement to enablement: conceptual models of disability in the 20th century", *Disability & Rehabilitation*, vol. 30, no. 17, pp. 1233-1244.
- May, S. 2010. "Self-management of chronic low back pain and osteoarthritis", *Nature reviews.Rheumatology*, vol. 6, no. 4, pp. 199-209.

McConnell S, Kolopack P, Davis AM. The Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC): a review of its utility and measurement properties. *Arthritis Rheum.* 2001; 45(5):453-461.

McDermott, S., Moran, R., Platt, T. & Dasari, S. 2006. "Variation in health conditions among groups of adults with disabilities in primary care", *Journal of community health*, vol. 31, no. 3, pp. 147-159.

McDonald-Miszczak, L. & Wister, A.V. 2005. "Predicting self-care behaviors among older adults coping with arthritis: a cross-sectional and 1-year longitudinal comparative analysis", *Journal of Aging & Health*, vol. 17, no. 6, pp. 836-857.

McDonough, C.M. & Jette, A.M. 2010. "The contribution of osteoarthritis to functional limitations and disability ", *Clinics in geriatric medicine*, vol. 26, no. 3, pp. 387-399.

McGauran, N., Wieseler, B., Kreis, J., Schuler, Y., Kolsch, H. & Kaiser, T. 2010. "Reporting bias in medical research - a narrative review", *Trials*, vol. 11, no. 1, pp. 37.

McKeough, DM. 2009. Available:

<http://www.google.co.uk/url?sa=t&rct=j&q=&esrc=s&frm=1&source=web&cd=3&ved=0CDsQFjAC&url=http%3A%2F%2Fwww.csus.edu%2Fdiv%2Fm%2Fmckeough%2FPT204%2FConceptMapsShow%2FDisablConcept%2520Map.pps&ei=HqRZUYqMBKKM0AXTrYC4BA&usq=AFQjCNGnZak4WgBxivHRn9PuXJyhsjrMxQ&sig2=hqbsutXs3Qv2qbPizaOotg> [2012, 12/15/2012].

Medical Research Council - A strategy for collaborative ageing research in the UK 2010. Available:

<http://www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC007289> [2012, 9/8/2012].

Melzer, D., Gardener, E. & Guralnik, J.M. 2005. "Mobility disability in the middle-aged: cross-sectional associations in the English Longitudinal Study of Ageing", *Age & Ageing*, vol. 34, no. 6, pp. 594-602.

Merskey and Bogduk - Pain: Anatomy 1994. Available:

<http://jubilation.uwaterloo.ca/~ranney/painanat.html> [2012, 9/16/2012].

Mertler, C.A. Designing scoring rubrics for your classroom. 2001. Available:

<http://pareonline.net/getvn.asp?v=7&n=25> [2012, 9/13/2012].

Michael, Y.L., Berkman, L.F., Colditz, G.A. & Kawachi, I. 2001. "Living arrangements, social integration, and change in functional health status", American Journal of Epidemiology, vol. 153, no. 2, pp. 123-131.

Michael, Y.L., Berkman, L.F., Colditz, G.A. & Kawachi, I. 2001. "Living arrangements, social integration, and change in functional health status", American Journal of Epidemiology, vol. 153, no. 2, pp. 123-131.

Miller, R.R., Zhang, Y., Silliman, R.A., Hayes, M.K., Leveille, S.G., Murabito, J.M., Kiel, D., O'Connor, G.T. & Felson, D.T. 2004. "Effect of medical conditions on improvements in self-reported and observed functional performance of elders", Journal of the American Geriatrics Society, vol. 52, no. 2, pp. 217-223.

Minaire, P. 1992. "Disease, illness and health: theoretical models of the disablement process", Bulletin of the World Health Organization, vol. 70, no. 3, pp. 373-379.

Moher, D., Cook, D.J., Eastwood, S., Olkin, I., Rennie, D. & Stroup, D.F. 1999. "Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of Reporting of Meta-analyses", Lancet, vol. 354, no. 9193, pp. 1896-1900.

Moher, D., Tetzlaff, J., Tricco, A.C., Sampson, M. & Altman, D.G. 2007. "Epidemiology and reporting characteristics of systematic reviews", Plos Medicine, vol. 4, no. 3, pp. 78-e78.

Mont, D. 2007. Measuring Disability Prevalence, Social Protection The World Bank.Washington.



- Montazeri, A., Vahdaninia, M., Ebrahimi, M. & Jarvandi, S. 2003. "The Hospital Anxiety and Depression Scale (HADS): translation and validation study of the Iranian version", *Health And Quality Of Life Outcomes*, vol. 1, pp. 14-14.
- Moore, D. & Tarnai, J. (eds) 2002. *Evaluating Nonresponse in Mail Surveys*, Groves, R. M., Dillman, D. A., Eltinge, J. L., & Little, R. J. A. (Eds.) *Survey Nonresponse* edn, Wiley-Interscience, New York.
- Moore, E.G., Rosenberg, M.W. & Fitzgibbon, S.H. 1999. "Activity limitation and chronic conditions in Canada's elderly, 1986-2011", *Disability and rehabilitation*, vol. 21, no. 5-6, pp. 196-210.
- Moore, M., Hofer, S., McGee, H. & Ring, L. 2005. "Can the concepts of depression and quality of life be integrated using a time perspective?", *Health and quality of life outcomes*, vol. 3, pp. 1.
- Moore, P. 2011. *The pain and self-care toolkit*. [Homepage of Pain toolkit], [Online]. Available: <http://www.pain toolkit.org/assets/downloads/Self-Care-Toolkit-Booklet-Nov-2011.pdf> [2012, September 4th].
- Mosby Medical Dictionary. 2009. *Osteoarthritis*. [Homepage of Elsevier.], [Online]. Available: <http://medical-dictionary.thefreedictionary.com/osteoarthritis> [2012, September 4th].
- Moskowitz, R.W. 2009. "The burden of osteoarthritis: clinical and quality-of-life issues", *The American Journal of Managed Care*, vol. 15, no. 8, pp. 223-S229.
- Muller, S. 2010. *The measurement of locomotor disability in epidemiological studies*, Keele University.
- Muller, S. & Roddy, E. 2009. "A rasch analysis of the Manchester foot pain and disability index", *Journal Of Foot And Ankle Research*, vol. 2, pp. 29-29.
- Mullick, A. 2005. *Bathing for older people with disabilities*. Available: <http://www.ap.buffalo.edu/idea/Publications/Bathing%20for%20Older%20People.htm> [2012, September, 9].

Murray, C. 2002. Summary measures of population health: concepts, ethics, measurement and applications World Health Organization, Geneva.

Myers, H., Nicholls, E., Handy, J., Peat, G., Thomas, E., Duncan, R., Wood, L., Marshall, M., Tyson, C., Hay, E. & Dziedzic, K. 2007. "The Clinical Assessment Study of the Hand (CAS-HA): a prospective study of musculoskeletal hand problems in the general population", BMC Musculoskeletal Disorders, vol. 8, pp. 85-85.

Naeim, A., Keeler, E.B. & Reuben, D. 2007. "Perceived causes of disability added prognostic value beyond medical conditions and functional status", Journal of clinical epidemiology, vol. 60, no. 1, pp. 79-85.

Nagi, S. 1965. Some conceptual issues in disability and rehabilitation, American Sociological Association, Washington, DC.

Nagi SZ. 1991. Disability concepts revised: implications for prevention. In: Pope AM, Tarlov AR, editors. Disability in America: Toward a national agenda for prevention. Washington (DC): National Academy Press.

Nagurney, J.T., Brown, D.F., Sane, S., Weiner, J.B., Wang, A.C. & Chang, Y. 2005. "The accuracy and completeness of data collected by prospective and retrospective methods ", Academic emergency medicine : official journal of the Society for Academic Emergency Medicine, vol. 12, no. 9, pp. 884-895.

Naik, A.D., Concato, J. & Gill, T.M. 2004. "Bathing disability in community-living older persons: common, consequential and complex", Journal of the American Geriatrics Society, vol. 52, no. 11, pp. 1805-1810.

Naik, A.D. & Gill, T.M. 2005. "Underutilization of environmental adaptations for bathing in community-living older persons", Journal of the American Geriatrics Society, vol. 53, no. 9, pp. 1497-1503.

Naik, A.D., Lai, J.M., Kunik, M.E. & Dyer, C.B. 2008. "Assessing capacity in suspected cases of self-neglect", Geriatrics, vol. 63, no. 2, pp. 24.

- Nakash, R.A., Hutton, J.L., Jørstad-Stein, E.,C., Gates, S. & Lamb, S.E. 2006. "Maximising response to postal questionnaires--a systematic review of randomised trials in health research", *BMC Medical Research Methodology*, vol. 6, pp. 5-5.
- Nante, N., Groth, N., Guerrini, M., Galeazzi, M., Kodraliu, G. & Apolone, G. 1999. "Using the SF-36 in a rural population of elderly in Italy; a pilot study.", *Journal of preventative medicine and hygiene.*, vol. 40, pp. 1.
- National Institute for Clinical Excellence. 2009. The guideline development group. National Institute for Clinical Excellence., London.
- Nelson, M.E., Rejeski, W.J., Blair, S.N., Duncan, P.W., Judge, J.O., King, A.C., Macera, C.A. & Castaneda-Sceppa, C. 2007. "Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association", *Medicine & Science in Sports & Exercise*, vol. 39, no. 8, pp. 1435-1445.
- Nicholls, E., E., A.,W.M., Jordan, J., L., Dziedzic, K., S. & Thomas, E. 2012. "Factors Associated with the Severity and Progression of Self-Reported Hand Pain and Functional Difficulty in Community-Dwelling Older Adults: A Systematic Review", *Musculoskeletal Care*, vol. 10, no. 1, pp. 51-62.
- NICE Osteoarthritis 2008. [Homepage of National Institute for Health and Clinical Excellence], [Online]. Available: <http://www.nice.org.uk/CG59> [2012, 9/13/2012].
- NIH 1993. Research Plan for the National Center for Medical Rehabilitation Research, NIH edn, National Institutes of Health, Department of Health and Human Services, Washington, DC.
- Noah, N. 2008. "The STROBE initiative: STrengthening the Reporting of OBservational studies in Epidemiology (STROBE)", *Epidemiology and infection*, vol. 136, no. 7, pp. 865-865.
- Nolte, E., Knai, C. & Mckee, M. 2008. Managing chronic conditions: experience in eight countries. World Health Organisation, Denmark.

Norburn, J.E., Bernard, S.L., Konrad, T.R., Woomert, A., DeFries, G.H., Kalsbeek, W.D., Koch, G.G. & Ory, M.G. 1995. "Self-care and assistance from others in coping with functional status limitations among a national sample of older adults", *The Journals Of Gerontology. Series B, Psychological Sciences And Social Sciences*, vol. 50, no. 2, pp. S101-S109.

Nordenfelt, L. 2003. *Action Theory, Disability And ICF*, *Disability and Rehabilitation* 2003;25:1075-79.

Nordenskiöld, U., Grimby, G., Hedberg, M., Wright, B. & Linacre, J.M. 1996. "The structure of an instrument for assessing the effects of assistive devices and altered working methods in women with rheumatoid arthritis ", *Arthritis Care and Research The Official Journal of the Arthritis Health Professions Association*, vol. 9, no. 5, pp. 358-367.

Norman, G.R. & Streiner, D.L. 2010. "Binary dependent variables: logistic regression", *Community Oncology*, vol. 7, no. 8, pp. 367-368.

Nusselder, W.J., Looman, C. & Mackenbach, J.P. 2006. "The level and time course of disability: trajectories of disability in adults and young elderly", *Disability & Rehabilitation*, vol. 28, no. 16, pp. 1015-1026.

Palacios-Ceña, D., Jiménez-García, R., Hernández-Barrera, V., Alonso-Blanco, C., Carrasco-Garrido, P. & Fernández-de-las-Peñas, C. 2012. "Has the Prevalence of Disability Increased Over the Past Decade (2000–2007) in Elderly People? A Spanish Population-based Survey", *Journal of the American Medical Directors Association*, vol. 13, no. 2, pp. 136-142.

Pallant, J.F. & Bailey, C.M. 2005. "Assessment of the structure of the Hospital Anxiety and Depression Scale in musculoskeletal patients", *Health And Quality Of Life Outcomes*, vol. 3, pp. 82-82.

Pan, A.W., Fan, C.W., Chung, L.I., Chen, T.J., Kielhofner, G., Wu, M.Y. & Chen, Y.L. 2011. "Examining the validity of the Model of Human Occupation Screening Tool: Using classical test theory and item response theory", *BRITISH JOURNAL OF OCCUPATIONAL THERAPY*, vol. 74, no. 1, pp. 34.

Papaioannou, E.C., R  ih  , I. & Kivel  , S. 2012."Self-neglect of the elderly. An overview", *The European Journal Of General Practice*, vol. 18, no. 3, pp. 187-190.

Paterson, D.H. & Warburton, D.E. 2010."Physical activity and functional limitations in older adults: a systematic review related to Canada's Physical Activity Guidelines", *The International Journal Of Behavioral Nutrition And Physical Activity*, vol. 7, pp. 38-38.

Peat, G., McCarney, R. & Croft, P. 2001. "Knee pain and osteoarthritis in older adults: a review of community burden and current use of primary health care ", *Annals of the Rheumatic Diseases*, vol. 60, no. 2, pp. 91-97.

Peat, G., Thomas, E., Handy, J., Wood, L., Dziedzic, K., Myers, H., Wilkie, R., Duncan, R., Hay, E., Hill, J. & Croft, P. 2004. "The Knee Clinical Assessment Study--CAS(K). A prospective study of knee pain and knee osteoarthritis in the general population", *BMC Musculoskeletal Disorders*, vol. 5, pp. 4-4.

Peat G, Thomas E, Handy J, Croft P. Social networks and pain interference with daily activities in middle and old age. *Pain*. 2004 Dec;112(3):397-405.

Peat, G., Greig, J., Wood, L., Wilkie, R., Thomas, E. & Croft, P. 2005. "Diagnostic discordance: we cannot agree when to call knee pain 'osteoarthritis'", *Family practice*, vol. 22, no. 1, pp. 96-102.

Peat, G., Thomas, E., Handy, J., Wood, L., Dziedzic, K., Myers, H., Wilkie, R., Duncan, R., Hay, E., Hill, J., Lacey, R. & Croft, P. 2006."The Knee Clinical Assessment Study-CAS(K). A prospective study of knee pain and knee osteoarthritis in the general population: baseline recruitment and retention at 18 months", *BMC Musculoskeletal Disorders*, vol. 7, pp. 30-30.

Peat, G., Thomas, E., Duncan, R., Wood, L., Wilkie, R., Hill, J., Hay, E.M. & Croft, P. 2007. "Estimating the probability of radiographic osteoarthritis in the older patient with knee pain", *Arthritis & Rheumatism: Arthritis Care & Research*, vol. 57, no. 5, pp. 794-802.

Peat, G. & Thomas, E. 2009. "When knee pain becomes severe: a nested case-control analysis in community-dwelling older adults", *Journal of Pain*, vol. 10, no. 8, pp. 798-808.

Peeters, A., Barendregt, J.J., Willekens, F., Mackenbach, J.P., Al Mamun, A., Bonneux, L. & NEDCOM the Netherlands Epidemiology and Demography Compression of Morbidity Research Group 2003. "Obesity in adulthood and its consequences for life expectancy: a life-table analysis ", *Annals of Internal Medicine*, vol. 138, no. 1, pp. 24-32.

Pepe, M.S. 2003. *The statistical evaluation of medical tests for classification and prediction*. New York, NY: Oxford.

Perenboom, R.J. & Chorus, A.M. 2003. "Measuring participation according to the International Classification of Functioning, Disability and Health (ICF) " *Disability and rehabilitation*, vol. 25, no. 11-12, pp. 577-587.

Period and cohort life expectancy tables, 2010. Available:  
<http://www.ons.gov.uk/ons/rel/lifetables/period-and-cohort-life-expectancy-tables/2010-based/index.html> [2012, 9/8/2012].

Peters, T.J., Sanders, C., Dieppe, P. & Donovan, J. 2005. "Factors associated with change in pain and disability over time: a community-based prospective observational study of hip and knee osteoarthritis", *British Journal of General Practice*, vol. 55, no. 512, pp. 205-211.

Pettersson, I., Pettersson, V. & Frisk, M. 2011. "ICF from an occupational therapy perspective in adult care: an integrative literature review", *Scandinavian Journal of Occupational Therapy*, vol. 19, no. 3, pp. 260-273.

Pham, B., Klassen, T.P., Lawson, M.L. & Moher, D. 2005. "Language of publication restrictions in systematic reviews gave different results depending on whether the intervention was conventional or complementary", *Journal of clinical epidemiology*, vol. 58, no. 8, pp. 769-776.

Picavet, H.S. & Schouten, J.S. 2003. "Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study", *Pain*, vol. 102, no. 1-2, pp. 167.

Pickles, A. 2011. Missing data, problems and solutions. [Homepage of University of Manchester], [Online]. Available: [www.uvm.edu/.../Missing\\_Data/missing%20dataPickles.doc](http://www.uvm.edu/.../Missing_Data/missing%20dataPickles.doc) [2012, September 4th].

Pincus, T. & Morley, S. 2001. "Cognitive-processing bias in chronic pain: a review and integration", *Psychological bulletin*, vol. 127, no. 5, pp. 599-617.

Pincus, T., Woodcock, A. & Vogel, S. 2010. "Returning back pain patients to work: how private musculoskeletal practitioners outside the national health service perceive their role (an interview study).", *Journal of Occupational Rehabilitation*, vol. 20, no. 3, pp. 322-330.

Pollard, B., Johnston, M. & Dieppe, P. 2006. "What do osteoarthritis health outcome instruments measure? Impairment, activity limitation, or participation restriction? ", *The Journal of rheumatology*, vol. 33, no. 4, pp. 757-763.

Pollard, B., Dixon, D., Dieppe, P. & Johnston, M. 2009. "Measuring the ICF components of impairment, activity limitation and participation restriction: an item analysis using classical test theory and item response theory", *Health And Quality Of Life Outcomes*, vol. 7, pp. 41-41.

Pollard, B., Johnston, M. & Dieppe, P. 2011. "Exploring the relationships between International Classification of Functioning, Disability and Health (ICF) constructs of Impairment, Activity Limitation and Participation Restriction in people with osteoarthritis prior to joint replacement", *BMC Musculoskeletal Disorders*, vol. 12, pp. 97-97.

Policy Digest: Common Core Principles To Support Self-Care, 2008. Independent Nurse, pp. 58-58.

Põlluste, K., Kallikorm, R., Mättik, E. & Lember, M. 2012. "Assistive devices, home adjustments and external help in rheumatoid arthritis", *Disability & Rehabilitation*, vol. 34, no. 10, pp. 839-845.

Pope, A.M. & Tarlov, A.R. 1991. *Disability in America: Toward a national agenda for prevention*. National Academy Press, Washington.

Porta, M. (ed) 2008. *Dictionary of Epidemiology*, 5th edition edn, Oxford University Press Inc., New York.

Pua Y-H, Cowan SM, Wrigley TV, Bennell KL. Discriminant validity of the Western Ontario and McMaster Universities Osteoarthritis Index physical functioning subscale in community samples with hip osteoarthritis. *Arch Phys Med Rehabil*. 2009; 90:1772-1777.

Public Health Resource Unit: Critical Appraisal Skills Programme 2008. Available: <http://www.phru.nhs.uk/casp/casp.htm> [2012, 9/13/2012].

Prior, J.A., Jordan, K.P. & Kadam, U.T. 2012. "Influence of chronic diseases on long-term change in physical health: a consultation-survey linkage cohort study in general practice", *Quality of Life Research*, vol. 21, no. 4, pp. 581-591.

Quintana, J.M., Arostegui, I., Escobar, A., Azkarate, J., Goenaga, J.I. & Lafuente, I. 2008 (a). "Prevalence of knee and hip osteoarthritis and the appropriateness of joint replacement in an older population", *Archives of Internal Medicine*, vol. 168, no. 14, pp. 1576-1584.

Quintana, J.M., Escobar, A., Arostegui, I., Bilbao, A., Armendariz, P., Lafuente, I. & Agirre, U. 2008 (b), "Prevalence of symptoms of knee or hip joints in older adults from the general population", *Aging Clinical and Experimental Research*, vol. 20, no. 4, pp. 329-336.

Rabins, P. 1996. "Barriers to diagnosis and treatment of depression in elderly patients", *American Journal of Geriatric Psychiatry*, vol. 4, pp. 79-84.



Radloff, L.S. 1977. Center for epidemiological studies depression scale. Available: [www.iprc.unc.edu/.../Center%20for%20Epidemiologic%20Studies%20Depression%20Scale.pdf](http://www.iprc.unc.edu/.../Center%20for%20Epidemiologic%20Studies%20Depression%20Scale.pdf) [2012, September 4th].

Rait, G., Fletcher, A., Smeeth, L., Brayne, C., Stirling, S., Nunes, M., Breeze, E., Ng, E.S., Bulpitt, C.J. & Jones, D. 2005. "Prevalence of cognitive impairment: results from the MRC trial of assessment and management of older people in the community", *Age and Ageing*, vol. 34, no. 3, pp. 242-248.

Read, J. 1991. "The Read Clinical Classification (Read codes): general description", *Br Homoeopath J*, vol. 80, no. 1, pp. 14.

Reed, G.M., Lux, J.B., Bufka, L.F., Trask, C., Peterson, D.B., Stark, S., Threats, T.T., Jacobson, J.W. & Hawley, J.A. 2005. "Operationalizing the International Classification of Functioning, Disability and Health in Clinical Settings. ", *Rehabilitation Psychology*, vol. 50, no. 2, pp. 122 - 131.

Rehab General Neuro-Musculo Best Practice Team 2004. Areas of function: Instrumental activities of daily living. Vancouver Health Authority. Vancouver.

Reid, M.C., Williams, C.S., Concato, J., Tinetti, M.E. & Gill, T.M. 2003 (a). "Depressive symptoms as a risk factor for disabling back pain in community-dwelling older persons", *Journal of the American Geriatrics Society*, vol. 51, no. 12, pp. 1710-1717.

Reid, M.C., Williams, C.S. & Gill, T.M. 2003 (b). "The relationship between psychological factors and disabling musculoskeletal pain in community-dwelling older persons", *Journal of the American Geriatrics Society*, vol. 51, no. 8, pp. 1092-1098.

Reid, M.C., Williams, C.S. & Gill, T.M. 2005. "Back pain and decline in lower extremity physical function among community-dwelling older persons", *The Journals Of Gerontology. Series A, Biological Sciences And Medical Sciences*, vol. 60, no. 6, pp. 793-797.

Reilly, M. 1974. *Play as exploratory learning: studies of curiosity behavior*, Sage, Beverly Hills.

Research for healthy ageing 2010. A strategy for collaborative ageing research in the UK, Medical Research Council, London.

Reynolds, S.L. & Silverstein, M. 2003. "Observing the onset of disability in older adults", *Social Science and Medicine*, vol. 57, no. 10, pp. 1875-1889.

Rodda, J., Walker, Z. & Carter, J. 2011. Musculoskeletal problems and work in the UK - Time for a new approach?", *Rheumatology*, vol. 48, no. 7, pp. 709-710  
"Depression in older adults", *BMJ (Clinical research ed.)*, vol. 343, pp. 5219-d5219.

Roddy, E., Muller, S. & Thomas, E. 2009. "Defining disabling foot pain in older adults: Further examination of the Manchester Foot Pain and Disability Index", *Rheumatology*, vol. 48, no. 8, pp. 992.

Rothman K.J., Greenland S., Lash T.L., eds. *Modern Epidemiology*. 3<sup>rd</sup> ed. Philadelphia: Lippincott-Raven, 2008.

Royston, P. 2009. Multiple imputation of missing values: Further update of ice, with an emphasis on categorical variables. *Stata Journal* 9(3): 466-477.

Royston, P. 2007. Multiple imputation of missing values: further update of ice, with an emphasis on interval censoring. *Stata Journal* (7)4: 445-464.

Royston, P. 2005. Multiple imputation of missing values: Update of ice., *Stata Journal* 5(4): 527-536.

Rubin, D.B. 1976. "Inference and missing data." *Biometrika*, vol. 63, no. 3, pp. 581.

Rubin, D.B. 1996. "Multiple imputation after 18+ years", *Journal of the American Statistical Association*, vol. 91: 434, no. 9/17/2012, pp. 473-489.

Rubin, D.B. 1987. *Multiple Imputation for Nonresponse in Surveys*. Wiley & Sons, New York.

Rüdiger, W. 2010. "Assessing nonresponse bias in activist surveys", *Quality & Quantity: International Journal of Methodology*, vol. 44, no. 1, pp. 173-180.

Saarni, S.I., Suvisaari, J., Sintonen, H., Koskinen, S., Harkanen, T. & Lonnqvist, J. 2007. "The health-related quality-of-life impact of chronic conditions varied with age in general population ", *Journal of clinical epidemiology*, vol. 60, no. 12, pp. 1288-1297.

Sackley, C. & Lett, K. 2007. "How fair is Fair Access to Care Services? A survey of occupational therapy teams", *International Journal of Therapy & Rehabilitation*, vol. 14, no. 8, pp. 340-340.

Sackley, C.M., van den Berg, M.,E., Lett, K., Patel, S., Hollands, K., Wright, C.C. & Hoppitt, T.J. 2009. "Effects of a physiotherapy and occupational therapy intervention on mobility and activity in care home residents: a cluster randomised controlled trial", *BMJ (Clinical research ed.)*, vol. 339, pp. 3123-b3123.

Salaffi, F., De Angelis, R. & Grassi, W. 2005. "Prevalence of musculoskeletal conditions in an Italian population sample: results of a regional community-based study. The MAPPING study", *Clinical and experimental rheumatology*, vol. 23, no. 6, pp. 819-828.

Sanderson, S., Tatt, I.D. & Higgins, J.P.T. 2007. "Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: a systematic review and annotated bibliography", *International journal of epidemiology*, vol. 36, no. 3, pp. 666-676.

Schäfer, I., Hansen, H., Schön, G., Höfels, S., Altiner, A., Dahlhaus, A., Gensichen, J., Riedel-Heller, S., Weyerer, S., Blank, W.A., König, H., von, d.K., Wegscheider, K., Scherer, M., van, d.B. & Wiese, B. 2012. "The influence of age, gender and socio-economic status on multimorbidity patterns in primary care. First results from the multicare cohort study", *BMC Health Services Research*, vol. 12, pp. 89-89.

Schafer, J.L. 1997. *Analysis of Incomplete Multivariate Data*. Chapman and Hall, London.

Schlesselman, J.J. 1982. *Case-Control Studies. Design, Conduct, Analysis*, Oxford University Press, New York.

Schouten, J.S., van den Ouweland, F.A. & Valkenburg, H.A. 1992. "A 12 year follow up study in the general population on prognostic factors of cartilage loss in osteoarthritis of the knee ", *Annals of the Rheumatic Diseases*, vol. 51, no. 8, pp. 932-937.

Schuntermann, M.F. 2005. "The implementation of the International Classification of Functioning, Disability and Health in Germany: experiences and problems", *International Journal of Rehabilitation Research*, vol. 28, no. 2, pp. 93-102.

Schröder A, Oernboel E, Licht RW, Sharpe M, Fink P. Outcome measurement in functional somatic syndromes: SF-36 summary scores and some scales were not valid. *J Clin Epidemiol*. 2012 Jan;65(1):30-41. doi: 10.1016/j.jclinepi.2011.03.012. Epub 2011 Jul 20.

Seshamani, M. 2004. The impact of ageing on healthcare expenditures: Impending crisis or misguided concern? Office for health economics. London.

Sharma, L., Cahue, S., Song, J., Hayes, K., Pai, Y. & Dunlop, D. 2003. "Physical functioning over three years in knee osteoarthritis: role of psychosocial, local mechanical, and neuromuscular factors", *Arthritis & Rheumatism*, vol. 48, no. 12, pp. 3359-3370.

Sharma, L., Song, J., Felson, D.T., Cahue, S., Shamiyeh, E. & Dunlop, D.D. 2001. "The role of knee alignment in disease progression and functional decline in knee osteoarthritis", *JAMA : the journal of the American Medical Association*, vol. 286, no. 2, pp. 188-195.

Shelton, R., C., McNeill, L., H., Puleo, E., Wolin, K., Y., Emmons, K., M. & Bennett, G., G. 2011. "The Association Between Social Factors and Physical Activity Among Low-Income Adults Living in Public Housing", *American Journal of Public Health*, vol. 101, no. 11, pp. 2102-2110.

Sherbourne, C.D., Asch, S.M., Shugarman, L.R., Goebel, J.R., Lanto, A.B., Rubenstein, L.V., Wen, L., Zubkoff, L. & Lorenz, K.A. 2009. "Early identification of co-occurring pain, depression and anxiety", *Journal of general internal medicine*, vol. 24, no. 5, pp. 620-625.

Silman, A., J. & Macfarlane, G.J. 2002. *Epidemiological Studies: A practical guide*, Cambridge University Press, Cambridge, United Kingdom.

Sim, J. & Wright, C.C. 2005. "The Kappa Statistic in Reliability Studies: Use, Interpretation, and Sample Size Requirements", *Physical Therapy*, vol. 85, no. 3, pp. 257-268.

Simeonsson, R.J., Leonardi, M., Bjorck-Akesson, E., Hollenweger, J., & Lollar, D. 2003. "Applying the International Classification of Functioning Disability and Health to measure childhood disability." *Disability & Rehabilitation* 25(11–12):602–610.

Simeonsson, R.J., Lollar, D., Hollowell, J. & Adams, M. 2000. "Revision of the International Classification of Impairments, Disabilities, and Handicaps: developmental issues", *Journal of clinical epidemiology*, vol. 53, no. 2, pp. 113-124.

Simon, H. 2009. [Homepage of University of Maryland Medical Center (UMMC) and A.D.A.M., Inc.], [Online]. Available: [http://www.umm.edu/patiented/articles/what\\_osteoarthritis\\_000035\\_1.htm](http://www.umm.edu/patiented/articles/what_osteoarthritis_000035_1.htm) [2012, 07/2012].

Sitzia, J. & Wood, N. 1998. "Response rate in patient satisfaction research: an analysis of 210 published studies ", *International Journal for Quality in Health Care*, vol. 10, no. 4, pp. 311-317.

Slater, M., Perruccio, A., V. & Badley, E., M. 2011. "Musculoskeletal comorbidities in cardiovascular disease, diabetes and respiratory disease: the impact on activity limitations; a representative population-based study", *BMC Public Health*, vol. 11, no. 1, pp. 77-77.

Song, J., Chang, R.W. & Dunlop, D.D. 2006. "Population impact of arthritis on disability in older adults", *Arthritis and Rheumatism*, vol. 55, no. 2, pp. 248-255.

South Staffordshire PCT and NHS North Staffordshire 2010. *Health and social profile of Staffordshire, North Staffordshire and South Staffordshire County Council*, Staffordshire.

Spector, W.D. & Fleishman, J.A. 1998. "Combining activities of daily living with instrumental activities of daily living to measure functional disability", *The Journals Of Gerontology. Series B, Psychological Sciences And Social Sciences*, vol. 53, no. 1, pp. 46-S57.

Spector, W.D., Katz, S., Murphy, J.B. & Fulton, J.P. 1987. "The hierarchical relationship between activities of daily living and instrumental activities of daily living", *Journal of chronic diseases*, vol. 40, no. 6, pp. 481-489.

Spratt, M., Carpenter, J., Sterne, J., Carlin, J.B., Heron, J., Henderson, J. & Tilling, K. 2010. "Strategies for multiple imputation in longitudinal studies", *American Journal of Epidemiology*, vol. 172, no. 4, pp. 478-487.

Staffordshire observatory. 2010. Indices of deprivation summary of findings for Staffordshire county., Staffordshire observatory, Staffordshire.

Stemler, Steven E. 2004. A comparison of consensus, consistency, and measurement approaches to estimating interrater reliability. Available: <http://www.pareonline.net/getvn.asp?v=9&n=4> [2012, 9/13/2012].

Sterne, J.A.C., White, I.R., Carlin, J.B., Spratt, M., Royston, P., Kenward, M.G., Wood, A.M. & Carpenter, J.R. 2009. "Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls", *BMJ (Clinical research ed.)*, vol. 338, pp. b2393-b2393.

Steven, M.M. 1992. "Prevalence of chronic arthritis in four geographical areas of the Scottish Highlands", *Annals of the Rheumatic Diseases*, vol. 51, no. 2, pp. 186-194.

Stewart, A.L., Hays, R.D. & Ware, J.E., J. 1988. "The MOS short-form general health survey. Reliability and validity in a patient population", *Medical care*, vol. 26, no. 7, pp. 724-735.

Steyerberg, E.W., Harrell, F.E., J., Borsboom, G.J., Eijkemans, M.J., Vergouwe, Y. & Habbema, J.D. 2001. "Internal validation of predictive models: efficiency of some procedures for logistic regression analysis", *Journal of clinical epidemiology*, vol. 54, no. 8, pp. 774-781.

Steyerberg, E.W., Eijkemans, M.J., Harrell, F.E., Jr & Habbema, J.D. 2000. "Prognostic modelling with logistic regression analysis: a comparison of selection and estimation methods in small data sets", *Statistics in medicine*, vol. 19, no. 8, pp. 1059-1079.

Steyerberg, E.W., Vickers, A.J., Cook, N.R., Gerds, T., Gonen, M., Obuchowski, N., Pencina, M.J. & Kattan, M.W. 2010. "Assessing the performance of prediction models: a framework for traditional and novel measures ", *Epidemiology (Cambridge, Mass.)*, vol. 21, no. 1, pp. 128-138.

Steyerberg, E.W., Bleeker, S.E., Moll, H.A., Grobbee, D.E. & Moons, K.G.M. 2003. "Internal and external validation of predictive models: a simulation study of bias and precision in small samples", *Journal of clinical epidemiology*, vol. 56, no. 5, pp. 441-447.

Stineman, M.G., Henry-Sanchez, J.T., Kurichi, J.E., Pan, Q., Xie, D., Saliba, D., Zhang, Z. & Streim, J.E. 2012. "Staging activity limitation and participation restriction in elderly community-dwelling persons according to difficulties in self-care and domestic life functioning", *Am J Phys Med Rehabil*, vol. 91, no. 2, pp. 126.

Stuck, A.E., Walthert, J.M., Nikolaus, T., Bula, C.J., Hohmann, C. & Beck, J.C. 1999. "Risk factors for functional status decline in community-living elderly people: a systematic literature review", *Social Science And Medicine*, vol. 48, no. 4, pp. 445.

Stucki, G., Ewert, T. & Cieza, A. 2002. "Value and application of the ICF in rehabilitation medicine ", *Disability and rehabilitation*, vol. 24, no. 17, pp. 932-938.

Suurmeijer, T.P., Doeglas, D.M., Moum, T., Briançon, S., Krol, B., Sanderman, R., Guillemin, F., Bjelle, A. & van den Heuvel, W.J. 1994. "The Groningen Activity Restriction Scale for measuring disability: its utility in international comparisons", *American Journal of Public Health*, vol. 84, no. 8, pp. 1270-1273.

Stratford PW, Kennedy DM, Woodhouse LJ, Spadoni GF. Measurement properties of the WOMAC LK 3.1 pain scale. *Osteoarthritis Cartilage*. 2006; 15:266-272.

STROBE statement--checklist of items that should be included in reports of observational studies (STROBE initiative) 2008. International Journal Of Public Health, vol. 53, 1, pp. 3-4.

Tackling inequalities in life expectancy in areas with the worst health and deprivation - National Audit Office 2010. [Homepage of Department of Health], [Online]. Available: [http://www.nao.org.uk/publications/1011/health\\_inequalities.aspx](http://www.nao.org.uk/publications/1011/health_inequalities.aspx) [2012, 9/17/2012].

Tamm, A., Lintrop, M., Veske, K., Hansen, U. & Tamm, A. 2008. "Prevalence of patello- and tibiofemoral osteoarthritis in Elva, Southern Estonia", The Journal of rheumatology, vol. 35, no. 3, pp. 543-544.

Tanna, S. 2004. Osteoarthritis: opportunities to address pharmaceutical gaps., World Health Organisation., London.

The British pain society & The Royal College of General Practitioners. 2004. A practical guide to the provision of chronic pain services for adults in primary care., The British Pain Society & The Royal College of General Practitioners., London.

The Health Survey of England 2005. [Homepage of The information centre for health and social care], [Online]. Available: <http://www.ic.nhs.uk/pubs/hse05olderpeople> [2012, September 9].

The National Collaborating Centre for Chronic Conditions 2008. Osteoarthritis national clinical guideline for care and management in adults., Royal college of physicians, London.

The National Statistics Socio-economic Classification (NS-SEC rebased on the SOC2010). Available: <http://www.ons.gov.uk/ons/guide-method/classifications/current-standard-classifications/soc2010/soc2010-volume-3-ns-sec--rebased-on-soc2010--user-manual/index.html> [2012, 9/9/2012].

The NHS Information Centre 2005. [Homepage of The NHS information centre for health and social care], [Online]. Available: <http://www.ic.nhs.uk/statistics-and-data-collections/social-care/adult-social-care-information> [2012, September 9].



The STROBE Statement: Home 2009. Available: <http://www.strobe-statement.org/> [2012, 9/13/2012].

The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies, 2007. Bulletin of the World Health Organization, vol. 85, no. 11, pp. 867-872.

The original bootstrap method. Available: <http://scholar.lib.vt.edu/theses/available/etd-61697-14555/unrestricted/Ch4.pdf> [2012, September 3rd].

Thomas, R. Income- commentary 1999. Available: <http://survey.net.ac.uk/sqb/topics/commentaries.asp> [2012, 9/16/2012].

Thomas, E., Peat, G., Harris, L., Wilkie, R. & Croft, P.R. 2004 (a). "The prevalence of pain and pain interference in a general population of older adults: cross-sectional findings from the North Staffordshire Osteoarthritis Project (NorStOP)", Pain, vol. 110, no. 1-2, pp. 361-368.

Thomas, E., Wilkie, R., Peat, G., Hill, S., Dziedzic, K. & Croft, P. 2004 (b). "The North Staffordshire Osteoarthritis Project--NorStOP: prospective, 3 years study of the epidemiology and management of clinical osteoarthritis in a general population of older adults", BMC Musculoskeletal Disorders, vol. 5, pp. 2-2.

Thomas, E., Peat, G., Mallen, C., Wood, L., Lacey, R., Duncan, R. & Croft, P. 2008. "Predicting the course of functional limitation among older adults with knee pain: do local signs, symptoms and radiographs add anything to general indicators?", Annals of the Rheumatic Diseases, vol. 67, no. 10, pp. 1390-1398.

Thomas, E., Mottram, S., Peat, G., Wilkie, R. & Croft, P. 2007. "The effect of age on the onset of pain interference in a general population of older adults: prospective findings from the North Staffordshire Osteoarthritis Project (NorStOP)", Pain, vol. 129, no. 1-2, pp. 21-27.

Tierney, R. & Simon, M. What's still wrong with rubrics: focusing on the consistency of performance criteria across scale levels. 2004. Available: <http://pareonline.net/getvn.asp?v=9&n=2> [2012, 9/13/2012].

- Tranmer, M. & Elliot, M. 2008. Binary logistic regression, <http://www.ccsr.ac.uk/publications/teaching/blr.pdf> edn, The Cathie Marsh centre for census and survey research.
- Trigg, R. & Wood, V.A. 2003, "The validation of the subjective index of physical and social outcome", *Clinical rehabilitation*, vol. 17, no. 3, pp. 238.
- Tripepi, G., Jager, K.J., Dekker, F.W. & Zoccali, C. 2009. "Statistical methods for the assessment of prognostic biomarkers(part II): calibration and re-classification ", *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*.
- Tsang, A., Von Korff, M., Lee, S., Alonso, J., Karam, E., Angermeyer, M.C., Borges, G.L., Bromet, E.J., de Girolamo, G., de Graaf, R., Gureje, O., Lepine, J., Haro, J.M., Levinson, D., Oakley Browne, M., Posada-Villa, J., Seedat, S., Watanabe, M. & de Graaf, R. 2008. "Common chronic pain conditions in developed and developing countries: gender and age differences and comorbidity with depression-anxiety disorders", *Journal of Pain*, vol. 9, no. 10, pp. 883-891.
- Turley, M., Tobias, M. & Paul, S. 2006. "Non-fatal disease burden associated with excess body mass index and waist circumference in New Zealand adults", *Australian & New Zealand Journal of Public Health*, vol. 30, no. 3, pp. 231-237.
- Turner, A., Foster, M. & Johnson, S. 2002. *Occupational Therapy and Physical Dysfunction, Principles, Skills and Practice 5th Edition* edn, Churchill Livingstone, Edinburgh.
- Twisk, J. & de Vente, W. 2002, "Attrition in longitudinal studies. How to deal with missing data", *Journal of clinical epidemiology*, vol. 55, no. 4, pp. 329-337.
- Occupational Therapy Practice Framework: domain and process, 2002. *The American Journal Of Occupational Therapy: Official Publication Of The American Occupational Therapy Association*, vol. 56, no. 6, pp. 609-639.
- Office of National Statistics (ONS): *Standard Occupational Classification 2000*. Available: <http://www.ons.gov.uk/ons/guide-method/classifications/archived->

[standard-classifications/standard-occupational-classification-2000/index.html](http://www.ons.gov.uk/ons/guide-method/classifications/archived-standard-classifications/standard-occupational-classification-2000/index.html)  
[2012, 9/16/2012].

Office of National Statistics (ONS): Standard Occupational Classification 2001.  
Available: <http://www.ons.gov.uk/ons/guide-method/classifications/archived-standard-classifications/standard-occupational-classification-2000/index.html>  
[2012, 9/16/2012].

Office of National Statistics (ONS): Deaths in older adults in England 2010.  
National End of Life Care Intelligence Network. ISBN: 978-0-9569225-4-0

Ueda, S. & Okawa, Y. 2003. "The subjective dimension of functioning and disability: what is it and what is it for?" *Disability & Rehabilitation*, vol. 25, no. 11, pp. 596.

UK National Statistics 2006. Demographic statistics: current UK population.  
Available: [www.statistics.gov.uk/](http://www.statistics.gov.uk/) [2012, September 3rd].

University of California 2004. Statistical computing seminars: Multiple imputation in stata part 1. [Homepage of University of California.], [Online]. Available: [http://www.ats.ucla.edu/stat/stata/seminars/missing\\_data/mi\\_in\\_stata\\_pt1.htm](http://www.ats.ucla.edu/stat/stata/seminars/missing_data/mi_in_stata_pt1.htm)  
[2012, September 4th].

University Of York 2009. Systematic reviews: CRD's guidance for undertaking reviews in health care. [Homepage of University of York], [Online]. Available: [www.york.ac.uk/inst/crd/SysRev/!SSL!/WebHelp/SysRev3.htm](http://www.york.ac.uk/inst/crd/SysRev/!SSL!/WebHelp/SysRev3.htm) [2012, September 4th].  
Definition of impairment and disability 2003. Available: [www.enham.org.uk/data/files/05\\_chapter\\_2.doc](http://www.enham.org.uk/data/files/05_chapter_2.doc) [2012, September/Monday].

Urwin, M., Symmons, D., Allison, T., Brammah, T., Busby, H., Roxby, M., Simmons, A. & Williams, G. 1998. "Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation", *Annals of the Rheumatic Diseases*, vol. 57, no. 11, pp. 649-655.

Üstün, T., Chatterji, S., Bickenbach, J., Kostanjsek, N. & Schneider, M. 2003. "The International Classification of Functioning, Disability and Health: a new tool for

understanding disability and health", *Disability & Rehabilitation*, vol. 25, no. 11, pp. 565-571.

Valanis, B. 1999. *Epidemiology in health care* Appleton & Lange, Stamford, Conn.

Van der Windt, D.A., Thomas, E., Pope, D.P., de Winter, A.F., Macfarlane, G.J., Bouter, L.M. & Silman, A.J. 2000, "Occupational risk factors for shoulder pain: a systematic review", *Occupational and environmental medicine*, vol. 57, no. 7, pp. 433-442.

Van Dijk, G.,M., Veenhof, C., Schellevis, F., Hulsmans, H., Bakker, J.P., Arwert, H., Dekker, J.H., Lankhorst, G.J. & Dekker, J. 2008. "Comorbidity, limitations in activities and pain in patients with osteoarthritis of the hip or knee", *BMC Musculoskeletal Disorders*, vol. 9, pp. 95-95.

Van Dijk, G., Veenhof, C., Lankhorst, G.J. & Dekker, J. 2009."Limitations in activities in patients with osteoarthritis of the hip or knee: the relationship with body functions, comorbidity and cognitive functioning", *Disability & Rehabilitation*, vol. 31, no. 20, pp. 1685-1691.

Van Houwelingen, J.C. & Le Cessie, S. 1990. "Predictive value of statistical models ", *Statistics in medicine*, vol. 9, no. 11, pp. 1303-1325.

Vandenbroucke, J.P., von Elm, E., Altman, D.G., Gotzsche, P.C., Mulrow, C.D., Pocock, S.J., Poole, C., Schlesselman, J.J., Egger, M. & STROBE Initiative 2007. "Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration ", *PLoS medicine*, vol. 4, no. 10, pp. e297.

Vargas-Chanes, D., Schroeder, D.R. & Offord, K.P. 2003. *An introduction to multiple imputation methods: handling missing data with SAS v8.2*, Mayo Foundation, USA.

Verbrugge, L.M. & Jette, A.M. 1994, "The disablement process", *Social science & medicine* (1982), vol. 38, no. 1, pp. 1-14.

- Verbrugge, L.M., Reoma, J.M. & Gruber-Baldini, A.L. 1994, "Short-term dynamics of disability and well-being", *Journal of health and social behavior*, vol. 35, no. 2, pp. 97-117.
- Verbrugge, L.M. & Juarez, L. 2001. "Profile of arthritis disability", *Public health reports*, vol. 116, pp. 157-179.
- Verzijl, N., Bank, R.A., TeKoppele, J.M. & DeGroot, J. 2003. "AGEing and osteoarthritis: a different perspective", *Current opinion in rheumatology*, vol. 15, no. 5, pp. 616-622.
- Victor, C.R., Triggs, E., Ross, F., Lord, J. & Axford, J.S. 2005. "Lack of benefit of a primary care-based nurse-led education programme for people with osteoarthritis of the knee", *Clinical rheumatology*, vol. 24, no. 4, pp. 358-364.
- Victor, C.R. 2005. *Social context of ageing*, Routledge, London, England.
- Von Elm, E., Altman, D.G., Egger, M., Pocock, S.J., Gøtzsche, P.C. & Vandenbroucke, J.P. 2007(a). "The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies", *Preventive Medicine: An International Journal Devoted to Practice and Theory*, vol. 45, no. 4, pp. 247-251.
- Von Elm, E. 2007 (b). "Commentary: Assessing the quality of observational studies--or a lesson from Mars", *International journal of epidemiology*, vol. 36, no. 3, pp. 677-678.
- Von Hippel, T.P. 2007. "Regression with missing Y's: An improved strategy for analyzing multiply imputed data." *Sociological methodology*, vol. 37, pp. 1-54.
- Von Korff, M., Ormel, J., Keefe, F.J. & Dworkin, S.F. 1992. "Grading the severity of chronic pain", *Pain*, vol. 50, no. 2, pp. 133-149.
- Wade, D.T. & Collin, C. 1988. "The Barthel ADL index: a standard measure of physical disability?" *International disability studies*, vol. 10, no. 2, pp. 64-67.

- Walters, S.J., Munro, J.F. & Brazier, J.E. 2001. "Using the SF-36 with older adults: a cross-sectional community-based survey", *Age and Ageing*, vol. 30, no. 4, pp. 337-343.
- Ward, G.E., Jagger, C. & Harper, W.M. 1996. "The Canadian occupational performance measure: what do users consider important?", *BRITISH JOURNAL OF THERAPY AND REHABILITATION*, vol. 3, no. 8, pp. 448.
- Ware, J.E., Jr & Sherbourne, C.D. 1992. "The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection", *Medical care*, vol. 30, no. 6, pp. 473-483.
- Ware, J.E. 1993. *SF-36 Health Survey: Manual and Interpretation Guide*, The Health Institute, New England Medical Center. Boston.
- Ware, J.E. (ed) 1995. *How to Score the SF-12 Physical and Mental Health Summary Scales*, 2nd edn, The Health Institute, New England Medical Center, Boston: MA.
- Ware, J.E. 2000. "SF-36 health survey update", *Spine*, vol. 25, no. 24, pp. 3130-3139.
- Ware, J.E. & Kosinski, M. 2001. "Interpreting SF-36 summary health measures: a response", *QUALITY OF LIFE RESEARCH*, vol. 10, no. 5, pp. 405.
- Watson, D., Pennebaker, J. W., & Folger, R. (1987). Beyond Negative Affectivity: Measuring stress and satisfaction in the workplace. *Journal of Organizational Behavior Management*, 8, 141-157.
- Weigl, M. 2003. "Linking osteoarthritis-specific health-status measures to the International Classification of Functioning, Disability, and Health (ICF) ", *Osteoarthritis and Cartilage*, vol. 11, no. 7, pp. 519 -523.
- Wells, G. 2010. Standardization (of rates): *Encyclopedia of Public Health 2002*. Available: <http://www.enotes.com/public-health-encyclopedia/standardization-rates> [2012].

Wen, X. 2004. "Trends in the prevalence of disability and chronic conditions among the older population: Implications for survey design and measurement of disability", *Australasian Journal on Ageing*, vol. 23, no. 1, pp. 3-6.

Wenham, C.Y.J. & Conaghan, P.G. 2009. "Imaging the painful osteoarthritic knee joint: what have we learned?", *Nature Clinical Practice.Rheumatology*, vol. 5, no. 3, pp. 149-158.

Westoby, C.J., Mallen, C.D. & Thomas, E. 2009. "Cognitive complaints in a general population of older adults: prevalence, association with pain and the influence of concurrent affective disorders", *European Journal of Pain*, vol. 13, no. 9, pp. 970-976.

White, A.M., Philogene, G.S., Fine, L. & Sinha, S. 2009. "Social Support and Self-Reported Health Status of Older Adults in the United States", *American Journal of Public Health*, vol. 99, no. 10, pp. 1872-1878.

White, I.R., Royston, P. & Wood, A.M. 2011. "Multiple imputation using chained equations: Issues and guidance for practice", *Statistics in medicine*, vol. 30, no. 4, pp. 377-399.

Whiteneck, G. 2006. *Conceptual models of disability AND past, present, and future*. National Academies Press, Washington, DC.

White paper - briefings for members 2010.

<http://www.nhsconfed.org/Networks/SDONet/Pages/WhitePaperBriefingForMembers.aspx>

WHO Scientific Group on the Burden of Musculoskeletal Conditions at the Start of the New Millennium 2003. "The burden of musculoskeletal conditions at the start of the new millennium ", *World Health Organization technical report series*, vol. 919, pp. i-x, 1-218, back cover.

WHO & ESCAP. 2008. *Training manual on disability statistics.*, World Health Organisation., Denmark.

Wiener, J.M., Hanley, R.J., Clark, R. & Van Nostrand, J.F. 1990. "Measuring the activities of daily living: comparisons across national surveys", *Journal of gerontology*, vol. 45, no. 6, pp. 229-S237.

Wilkie, R., Peat, G., Thomas, E. & Croft, P.R. 2004. "Measuring the consequences of osteoarthritis and joint pain in population-based studies: can existing health measurement instruments capture levels of participation?" *Arthritis and Rheumatism*, vol. 51, no. 5, pp. 755-762.

Wilkie, R. 2005. A study of participation restriction and joint pain in community-dwelling older adults, Keele University.

Wilkie, R., Peat, G., Thomas, E., Hooper, H. & Croft, P.R. 2005. "The Keele Assessment of Participation: A new instrument to measure participation restriction in population studies. Combined qualitative and quantitative examination of its psychometric properties", *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation*, vol. 14, no. 8, pp. 1889-1899.

Wilkie, R., Peat, G., Thomas, E. & Croft, P. 2006. "The prevalence of person-perceived participation restriction in community-dwelling older adults", *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation*, vol. 15, no. 9, pp. 1471-1479.

Wilkie, R., Peat, G., Thomas, E. & Croft, P. 2007 (a). "Factors associated with participation restriction in community-dwelling adults aged 50 years and over", *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*, vol. 16, no. 7, pp. 1147-1156.

Wilkie, R., Peat, G., Thomas, E. & Croft, P. 2007 (b). "Factors associated with restricted mobility outside the home in community-dwelling adults ages fifty years and older with knee pain: an example of use of the International Classification of Functioning to investigate participation restriction", *Arthritis and Rheumatism*, vol. 57, no. 8, pp. 1381-1389.



Wilkie, R. & Peat, G. 2008. "Social participation and osteoarthritis: the foundations have been laid where next?" *The Journal of rheumatology*, vol. 35, no. 8, pp. 1484-1486.

Wilkie, R., Thomas, E., Mottram, S., Peat, G. & Croft, P. 2008. "Onset and persistence of person-perceived participation restriction in older adults: a 3 years follow-up study in the general population", *Health And Quality Of Life Outcomes*, vol. 6, pp. 92-92.

Wilkie, R., Jordan, J.L., Muller, S., Nicholls, E., Healey, E.L. & Van der Windt, D.,A. 2011. "Measures of social function and participation in musculoskeletal populations: Impact on Participation and Autonomy (IPA), Keele Assessment of Participation (KAP), Participation Measure for Post-Acute Care (PM-PAC), Participation Objective, Participation Subjective (POPS), Rating of Perceived Participation (ROPP), and The Participation Scale", *Arthritis Care & Research*, vol. 63 Suppl 11, pp. 325-S336.

Wolfe, F. & Kong, S.X. 1999. "Rasch analysis of the Western Ontario MacMaster Questionnaire (WOMAC) in 2205 patients with osteoarthritis, rheumatoid arthritis, and fibromyalgia", *Annals of the Rheumatic Diseases*, vol. 58, no. 9, pp. 563.

Wood, L.R., Peat, G., Thomas, E. & Duncan, R. 2007. "Knee osteoarthritis in community-dwelling older adults: are there characteristic patterns of pain location?" *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society*, vol. 15, no. 6, pp. 615-623.

Wood, L., Peat, G., Thomas, E., Hay, E.M. & Sim, J. 2008. "Associations Between Physical Examination and Self-Reported Physical Function in Older Community-Dwelling Adults With Knee Pain", *Physical Therapy*, vol. 88, no. 1, pp. 33-42.

Wood-Dauphinee, S., Opzoomer, M.A., Williams, J.I., Marchand, B. & Spitzer, W.O. 1988. "Assessment of global function: the Reintegration to Normal Living Index", *Archives of Physical Medicine & Rehabilitation*, vol. 69, no. 8, pp. 583-590.

Woods, D., Kirk, D., Agarwal, S., Annandale, E., Arthur, T., Harvey, J., Hsu, R., Katbamna, S., Olsen, R., Smith, L., Riley, R. & Sutton, A. Vulnerable groups and

access to health care: a critical interpretive review 2005. Available: [http://www.netscc.ac.uk/hsdr/files/project/SDO\\_FR\\_08-1210-025\\_V01.pdf](http://www.netscc.ac.uk/hsdr/files/project/SDO_FR_08-1210-025_V01.pdf) [2012, 9/17/2012].

Woolf, A.D. & Pfleger, B. 2003, "Burden of major musculoskeletal conditions", Bulletin of the World Health Organization, vol. 81, no. 9, pp. 646-656.

World Health Organisation. 1980. International classifications disabilities and handicaps., World Health Organisation., Geneva.

World Health Organization 2001. International Classification of Functioning, Disability and Health, Geneva: WHO.

World Health Organization: Employment Conditions and Health Inequalities 2007. [Homepage of World Health Organisation], [Online]. Available: [http://www.who.int/social\\_determinants/resources/articles/emconet\\_who\\_report.pdf](http://www.who.int/social_determinants/resources/articles/emconet_who_report.pdf) [2012, 9/17/2012].

World Federation of Occupational Therapy 2011. Available: <http://www.wfot.org/> [2012, 9/8/2012].

World Health Organization 2003. Available: [http://whqlibdoc.who.int/trs/WHO\\_TRS\\_919.pdf](http://whqlibdoc.who.int/trs/WHO_TRS_919.pdf) [2012, 9/13/2012].

Wu, S.C., Leu, S. & Li, C. 1999. "Incidence of and predictors for chronic disability in activities of daily living among older people in Taiwan", Journal of the American Geriatrics Society, vol. 47, no. 9, pp. 1082-1086.

Wynne-Jones, G., Mallen, C.D., Main, C.J. & Dunn, K.M. 2010. "Sickness certification and the GP: what really happens in practice?", Family practice, vol. 27, no. 3, pp. 344-350.

Xiang, H., Leff, M. & Stallones, I. 2005. "Non-fatal injuries among adults with activity limitations and participation restrictions", Injury Prevention, vol. 11, no. 3, pp. 157-162.

Yang, Y. & George, L.K. 2005. "Functional disability, disability transitions, and depressive symptoms in late life", *Journal of aging and health*, vol. 17, no. 3, pp. 263-292.

Yelin, E.H. & Katz, P.P. 1990. "Transitions in health status among community-dwelling elderly people with arthritis. A national, longitudinal study", *Arthritis and Rheumatism*, vol. 33, no. 8, pp. 1205-1215.

Yelin, E., Lubeck, D., Holman, H. & Epstein, W. 1987. "The impact of rheumatoid arthritis and osteoarthritis: the activities of patients with rheumatoid arthritis and osteoarthritis compared to controls", *The Journal of rheumatology*, vol. 14, no. 4, pp. 710-717.

Young, A.F., Powers, J.R. & Bell, S.L. 2006. "Attrition in longitudinal studies: who do you lose?", *Australian & New Zealand Journal of Public Health*, vol. 30, no. 4, pp. 353-361.

Zelman, D., C., Hoffman, D., L., Seifeldin, R. & Dukes, E., M. 2003. "Development of a metric for a day of manageable pain control: derivation of pain severity cut-points for low back pain and osteoarthritis", *Pain (03043959)*, vol. 106, no. 1-2, pp. 35-42.

Zigmond, A.S. & Snaith, R.P. 1983. "The hospital anxiety and depression scale", *Acta Psychiatrica Scandinavica*, vol. 67, no. 6, pp. 361-370.

## **Appendix 1: Baseline Health Questionnaire**

# Health Questionnaire

Would you be willing to be contacted again about the possibility of taking part in this study? Giving us your permission to contact you again does not mean that you must take part. You are just agreeing to be contacted again.

Yes, I am happy to be contacted again.....☐

No, I do not want to be contacted again.....☐

It is important for us to find out what types of treatments and tests people need. we can do this by reviewing medical records.

Would you be willing to give your permission for this? When we review the medical records, your name will not be used so that you will not be identified personally. We can assure you that any information will be held in the strictest confidence.

Yes, I give my permission for my medical records to be reviewed...☐

No, I do not want my medical records to be reviewed.....☐

Signed ..... Date .....

Please print your name and address - .....

.....  
.....  
.....

**Thank you for taking the time to fill in this  
questionnaire.**

## **SECTION C - CONTINUING TO HELP WITH THIS STUDY**

Thank you very much for completing this questionnaire.

There will be more stages to this study, and we hope that people who have taken part will be able to help us again.


Please complete the two questions on the opposite page with regards to further help.

**Even if you have decided not to participate further in this study or would prefer us not to review your records, the answers in this questionnaire will still be very useful to us**

**Please check that you have answered all the questions.**

**Please return this questionnaire in the envelope provided (no stamp needed).**

**If you have any questions, telephone Ross Wilkie on 01782 583904.**



## **Instructions for this questionnaire**

The aim of this questionnaire is to find out about the general health of local people and about how many people suffer from joint pain. The answers you give in the questionnaire will be treated in the **strictest confidence**.

**Please answer even if you have no problems with your health**

Please answer all of the questions unless the instructions ask you to do something else.

Please write in **BLOCK CAPITALS** where appropriate.

Most of the questions can be answered by putting a cross in a box like this: ☒

When you have finished please check that you have answered all of the questions and then return the questionnaire in the envelope enclosed. You do not need a stamp. Please return the questionnaire in the next two weeks.

More details about this project are available in the information sheet enclosed. If you would like further information about this project please contact Ross Wilkie, survey co-ordinator, on 01782 583904.

**Thank you for your help with this important research project**

**Section A**

This section is made up of questions about your health, the activities you do, and some of the ways in which people do things in everyday life.

Please answer each set of questions as the instructions tell you to.

**Thank you for completing Section B**  
**Please turn to Section C**



Part 4

The following items are about activities you might do during a typical day. Does **your health now limit you** in these activities? If so, how much?  
*(Please put a cross in one box on each line)*

	Yes, limited a lot	Yes, limited a little	No,not limited at all
a. <b>Vigorous activities</b> , such as running, lifting heavy objects, participating in strenuous sports.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. <b>Moderate activities</b> , such as moving a table, pushing a vacuum cleaner, bowling or playing golf.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Lifting or carrying groceries.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Climbing <b>several</b> flights of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Climbing <b>one</b> flight of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Bending, kneeling, or stooping.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Walking <b>more than a mile</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Walking <b>half a mile</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Walking <b>100 yards</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Bathing or dressing yourself.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 1 - Your health

We are interested in your general health.

**Please answer every question.** Some questions may look similar to others but each one is different. We would like you to answer each one. Please take the time to read and answer each question carefully by placing a cross in the box of your choice. Please cross one box only on each line.

1. In general would you say your health is:  
*(Please put a cross in one box only)*

Excellent	Very good	Good	Fair	Poor
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. The following questions are about activities you might do during a typical day. Does your health now limit you in these activities? If so, how much?  
*(Please put a cross in one box on each line)*

	Yes, limited a lot	Yes, limited a little	No, not limited at all
a. <b>Moderate activities</b> , such as moving a table, pushing a vacuum cleaner, bowling or playing golf.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Climbing <b>several</b> flights of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. During the **past 4 weeks**, have you had any of the following problems with your work or other regular daily activities as a result of your physical health?  
*(Please put a cross in one box on each line)*

	Yes	No
a. <b>Accomplished less</b> than you would like.....	<input type="checkbox"/>	<input type="checkbox"/>
b. Were limited in the <b>kind</b> of work or other activities .....	<input type="checkbox"/>	<input type="checkbox"/>

4. During the **past 4 weeks**, have you had any of the following problems with your work or other regular daily activities as a result of any emotional problems (such as feeling depressed or anxious)?

*(Please put a cross in one box on each line)*

Yes No

a) **Accomplished less** than you would like..... ☐ ☐

b) Didn't do work or other activities as **carefully** as usual..... ☐ ☐

5. During the **past 4 weeks**, how much did **pain** interfere with your normal work (including both work outside the home and housework)?

*(Please put a cross in one box only)*

Not at all      A little bit      Moderately      Quite a bit      Extremely

☐      ☐      ☐      ☐      ☐

6. These questions are about how you feel and how things have been with you during the **past 4 weeks**. For each question, please give the one answer that comes closest to the way you have been feeling. How much of the time during the **past 4 weeks**.....

*(Please put a cross in one box on each line)*

	All of the time	Most of the time	A good bit of the time	Some of the time	A little of the time	None of the time
a) Have you felt calm and peaceful?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b) Did you have a lot of energy?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c) Have you felt downhearted and depressed?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. During the **past 4 weeks**, how much of the time has your physical health or emotional problems interfered with your social activities (like visiting friends, relatives, etc.)? *(Please put a cross in one box only)*

All of the time      Most of the time      Some of the time      A little of the time      None of the time

☐      ☐      ☐      ☐      ☐

We are also interested in **any problems** you may have had with your **legs**. For these questions, please think about problems with your **hip, knee or foot**; for example **pain, stiffness, giving way or locking**.

3. Have you had any problems with your **KNEES**, over the **last year**?

*(Please put a cross in one box only)*

Yes.... ☐      No.... ☐

4. Have you had pain in the last year in and **around the KNEE**?

*(Please put a cross in one box only)*

Yes.... ☐      No.... ☐

5. Have you had any problems with your **HIPS**, over the **last year**?

*(Please put a cross in one box only)*

Yes.... ☐      No.... ☐

6. Have you had pain in the last year in and **around the HIP**?

*(Please put a cross in one box only)*

Yes.... ☐      No.... ☐

7. Have you had any problems with your **FEET**, over the **last year**?

*(Please put a cross in one box only)*

Yes.... ☐      No.... ☐

8. Have you had pain in the last year in and **around the FOOT**?

*(Please put a cross in one box only)*

Yes.... ☐      No.... ☐

Part 2

It would be helpful if you could tell us about any tablets, pills or creams you may have used to reduce pain, in the **past 4 weeks**.

1. In the **past 4 weeks**, how often have you taken any medicines for your recent pain? **(Please put a cross in one box on each line)**

	All days	Most days	Some days	Few days	No days
a. Painkillers, Anti-inflammatories (e.g paracetamol, aspirin, Diclofenac,Ibuprofen).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Creams, gels or rub-ons (e.g. Ibuleve, Ibuprofen, Ralgex).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Natural medicine(e.g. herbal remedies, cod liver oil).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Glucosamine/Chondroitin Sulphate.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 3

We are interested to know how many people suffer from specific joint pains and problems. Please answer each of the following questions about your hands, hips, knees, and feet, even if you do not suffer from any problems.

We would like you to tell us if you have had **any problems** with your **hands**. **'Hand problems'** relate to **any hand symptoms** you may have experienced; for example, **pain or stiffness**, or any **knobbly swellings** on your hands, including your fingers and thumbs.

1. Have you had any problems with your **HANDS**, including your fingers and thumbs, over the **last year**?

**(Please put a cross in one box only)**

Yes....☐ No....☐

2. Have you had any pain in your **HANDS**, including in your fingers and thumbs, over the **last year**?

**(Please put a cross in one box only)**

Yes....☐ No....☐

Part 2 - Taking Part

We are interested in some of the things that are necessary for you to live your life in the way you choose. We are particularly interested in how often these things are achieved in the way you would like.

When answering the questions, please think about the **past 4 weeks**. It does not matter if you require the help of other people or from gadgets and machines. We would simply like to know if the activity **IS** achieved to the extent that you want it to be.

Please read each statement below and put a cross in the box which comes closest to how much you agree with the statement. Please put a cross in one box only for each line.

1. During the past 4 weeks, I have moved around in my home, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. During the past 4 weeks, I have moved around outside my home, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. During the past 4 weeks, my self-care needs (examples are washing, toileting, dressing, feeding, maintaining health) have been met, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. During the past 4 weeks, my home has been looked after, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. During the past 4 weeks, my things (belongings) have been looked after, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. Do you have any relatives, or other people, who depend on you?

Yes.... ☐                      No.... ☐

If yes, during the past 4 weeks, were these people looked after, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. During the past 4 weeks, I have met and spoken to other people, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. During the past 4 weeks, I, or someone else on my behalf, have managed my money, **as I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 1 - Body Chart

This question is about any **recent pain** you may have had in **any part of your body**. By pain we also mean ache, discomfort or stiffness. Please **do not** include pain due to a feverish illness such as flu. If you are a woman please **do not** include pain related to your monthly period.


1. In the **past 4 weeks**, have you had pain that has lasted for **one day or longer** in **any part of your body**?

Yes ..... ☐

Please shade in the diagram below **any pain** that has lasted for **one day or longer in the past 4 weeks**

No ..... ☐

Please turn the page over and continue with **Part 2**.



Please turn over to complete Part 2

**Section B**

This section is about any pains you may have or any problems with your joints.

Please fill in all the parts even if you do not suffer from any pains or joint problems.

Please follow the instructions for each part.

9. Do you choose to take part in paid or voluntary work?

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in paid or voluntary work, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Do you choose to take part in education or training courses?

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in education or training, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

11. Do you choose to take part in social activities?  
(Examples of social activities are community and religious activities, meeting up with friends, going to clubs)

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in social activities, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 3 - How often

We would like to know about how often you do things during your normal daily routine.

Look at the following list. Thinking about the past 4 weeks, please consider how often you did each thing, and put a cross in one box on each line.

	All days	Most days	Some days	Few days	No days
a. Go out for a walk.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Go out of the house and go somewhere.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Go out to work.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Spend most or all of the day in bed or in a chair.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Go shopping.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Take a bus or drive a car.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Go in a car as a passenger.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Go to a club, church or social event.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Play a sport.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Go on an education or training course.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Do a hobby.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
l. Take a bath/shower.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
m. Do home maintenance activities.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Thinking over the **past 4 weeks**, did you?

(Please put a cross in one box on each line)

	Not at all	On some nights	On most nights
a. Have trouble falling asleep.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Wake up several times per night.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Have trouble staying asleep.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Wake up after your usual amount of sleep feeling tired and worn out.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Thank you for completing Section A

Please turn to Section B

3. Please put a cross in the box to show whether you agree (yes box) or disagree (no box) with each of the following statements.....

(Please put a cross in one box on each line)

a. I am confused and start to do more than one thing at a time.....

Yes....☐

No....☐

b. I have more minor accidents than usual (e.g.I drop things, I trip and fall, or bump into things).....

Yes....☐

No....☐

c. I react slowly to things that are said or done.....

Yes....☐

No....☐

d. I do not finish things that I start.....

Yes....☐

No....☐

e. I have difficulty reasoning and solving problems (e.g. making plans, making decisions, or learning new things).....

Yes....☐

No....☐

f. I sometimes get confused (e.g. I do not know where I am, who is around, or what day it is).....

Yes....☐

No....☐

g. I forget a lot (e.g. things that happened recently, where I put things, or to keep appointments).....

Yes....☐

No....☐

h. I do not keep my attention on any activity for long.....

Yes....☐

No....☐

i. I make mistakes more than usual.....

Yes....☐

No....☐

j. I have difficulty doing things which involve thought and concentration.....

Yes....☐

No....☐

	All days	Most days	Some days	Few days	No days
n. Wash clothes.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
o. Cook and clean.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
p. Look after others.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
q. Heavy housework (e.g. spring cleaning, moving furniture, scrubbing floors by hand).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
r. Heavy gardening (e.g.digging, tree felling).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
s. Heavy DIY work at home (e.g. decorating, plastering).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
t. Walks of at least a quarter of a mile (5-10 minutes continuous walking).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
u. Walks of two miles or more (at least 40 minutes continuous walking).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

art 4 - How you do things

We would like to know if you have changed the way you do your normal daily activities and if you require the help of other people or special devices.

Please answer each question and put a cross in one box for each one.

1. Thinking back over the **past 4 weeks**, have you had to reduce the amount of time or change how you have done most activities because of your health?

All days	Most days	Some days	Few days	No days
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. During the **past 4 weeks** have you required the assistance of others or aids (examples are a walking stick or a wheelchair) to move around your home?

Yes... ☐No... ☐

3. During the **past 4 weeks**, have you required the assistance of others or aids (examples are a walking stick or a wheelchair) to go to places outside of your home?

Yes... ☐No... ☐

4. During the **past 4 weeks** have you required the assistance of others to look after yourself?

Yes... ☐No... ☐

5. During the **past 4 weeks**, have you required the assistance of others to look after your home?

Yes... ☐No... ☐

6. Compared to **12 months ago**, have you reduced the time or changed how you have done any of your activities?

Yes, a lot

☐

Yes, a little

☐

No, not at all

☐

### **Part 5 - Friends and Family**

We are interested in the contact you may have with your friends and family.

**Please answer each question and put a cross in one box for each line.**

1. How often do you go to religious meetings or services?

More than  
once a  
week☐Once a  
week☐1 to 3  
times per  
month☐Less than  
once per  
month☐Never or  
almost  
never☐

### **Part 10 - About your health**

We would like to know if you have any other health problems. Please put a cross in the box if you suffer from any of the listed problems.

1. Do you suffer from any of the following?

**(Please place a cross in the box for any which apply to you)**

a. Chest problems..... ☐b. Heart problems..... ☐c. Deafness..... ☐d. Problems with eyesight  
(excluding the need for glasses)..... ☐e. Raised blood pressure..... ☐f. Diabetes..... ☐

2. Thinking back over **the past 3 months**, have you suffered from any of the following?

**(Please place a cross in the box for any which apply to you)**

a. A fall or falls..... ☐b. Difficulty remembering things..... ☐c. Cough with spit..... ☐d. Breathless when walking..... ☐e. Dizziness or unsteadiness..... ☐f. Weakness in an arm or leg..... ☐



11. On average, how often do you drink alcohol?  
*(Please put a cross in one box only)*

Daily or most days	Once or twice a week	Once or twice a month	Once or twice a year	Never
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. How old were you when you left school?  years old

13. Did you go from school to full-time education or university?

Yes.....☐

If yes, what age did you finish full-time education?

years old

No.....☐

14. Have you gained qualifications through study as an adult?

Yes.....☐

No.....☐

15. Thinking about the cost of living as it affects you, which of these descriptions best describes your situation:  
*(Please put a cross in one box only)*

Find it a strain to get by from week to week.....	<input type="checkbox"/>
Have to be careful with money.....	<input type="checkbox"/>
Able to manage without much difficulty.....	<input type="checkbox"/>
Quite comfortably off.....	<input type="checkbox"/>

16. Is your ethnic origin? *(Please put a cross in one box only)*

White UK/European.....	<input type="checkbox"/>	Asian.....	<input type="checkbox"/>
Afro Caribbean.....	<input type="checkbox"/>	African.....	<input type="checkbox"/>
Chinese.....	<input type="checkbox"/>	Other.....	<input type="checkbox"/>

2. How many hours **each week** do you participate in any groups such as social or work group, church-connected group, self-help group, charity, public service or community group?

None	1 to 2 hours	3 to 5 hours	6 to 10 hours	11 to 15 hours	16 or more hours
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. How many living children do you have?

None	1 to 2	3 to 5	6 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. How many of your children do you see at least **once a month**?

None	1 to 2	3 to 5	6 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Apart from your children, how many relatives do you have with whom you feel close?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. How many close relatives do you see at least **once a month**?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. How many close friends do you have?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. How many of these friends do you see at least **once a month**?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. Is there any one special person you know that you feel very close to; someone you feel you can share confidences and feelings with?

Yes... ☐

No... ☐

If yes, how often do you see or talk with this person?

Daily

☐

Weekly

☐

Monthly

☐

Several times  
a year

☐

Once  
a year  
or less

☐

### **Part 6 - Access**

We would like you to tell us how easy it is to get hold of or gain access to things that you need in daily life.

***(Please put a cross in one box for each question)***

1. Do you have access to a car when you personally need it?

Yes... ☐

No... ☐

2. Do you have access to public transport?

Yes... ☐

No... ☐

3. Do you have access to a telephone?

Yes... ☐

No... ☐

4. Do you have good access to your doctor (GP), as and when you need it?

Yes... ☐

No... ☐

5. Do you have access to a chemist?

Yes... ☐

No... ☐

7. Do you have a spouse or partner who is currently living with you?

Yes... ☐

No... ☐

If they are working, what is their job title (examples - factory worker, welder, office worker, shop assistant, lawyer)?

.....

If they are not working or are retired, what was their last job title?

.....

If you are a widow, or widower, what was your spouse's last job title?

.....

8. What is your weight?  stones  lbs or  kgs

9. What is your height?  feet  inches or  cms

10. What is your current smoking status?

***(Please put a cross in one box only)***

Never smoked..... ☐

Previously smoked..... ☐

Currently smoking..... ☐

2. Are you:

Female.....☐

Male.....☐

3. What is your current marital status? *(Please put a cross in one box only)*

Married.....☐

Separated.....☐

Divorced.....☐

Widowed.....☐

Cohabiting.....☐

Single.....☐

4. Do you live alone?

Yes.....☐

No.....☐

5. What is your current employment status?  
*(Please put a cross in one box only)*

Employed.....☐

Not working due to ill health.....☐

Retired.....☐

Unemployed/seeking work.....☐

Housewife.....☐

Other.....☐

6. If working, what is your job title (examples - factory worker, welder, office worker, shop assistant, lawyer) ?

.....

If you are not working, or are retired, what was your last job title?

.....

6. Do you have access to a bank?

Yes...☐

No...☐

7. Do you have access to advice or help with your income (for example, relatives or the benefits system)?

Yes...☐

No...☐

8. If you wanted to take part in an education course, is there the opportunity?

Yes...☐

No...☐

9. If you wanted to do paid or voluntary work, is there the opportunity?

Yes...☐

No...☐

**Part 7 - How you feel**

The next set of questions are about how you feel at the moment. Please read each item and put a cross next to the reply that comes closest to how you have been feeling **in the past week**. Don't take too long over your replies; your immediate reaction to each item will usually be more accurate than a long thought out response.

1. I feel tense or 'wound up':

Most of the time

A lot of the time

From time to time, occasionally

Not at all

☐

☐

☐

☐

2. I still enjoy the things I used to enjoy:

Definitely as much

Not quite as much

Only a little

Hardly at all

☐

☐

☐

☐

3. I get a sort of frightened feeling as if something awful is about to happen:

Very definitely and quite badly      Yes, but not too badly      A little, but it doesn't worry me      Not at all

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------

4. I can laugh and see the funny side of things:

As much as I always could      Not quite so much now      Definitely not so much now      Not at all

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------

5. Worrying thoughts go through my mind:

A great deal of the time      A lot of the time      Not too often      Very little

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------

6. I feel cheerful:

Never      Not often      Sometimes      Most of the time

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------

7. I can sit at ease and feel relaxed:

Definitely      Usually      Not often      Not at all

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------

8. I feel as if I am slowed down:

Nearly all the time      Very often      Sometimes      Not at all

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------

Strongly disagree      Disagree      Neither agree nor disagree      Agree      Strongly agree

i. Problems with your joints are a normal part of growing old.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

j. Doctors can do a lot to help people with joint problems.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

k. Joint problems always get worse over time.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

Treatments are effective in controlling pain.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

m. If a pain lasts for a week or more, you have a serious illness.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

n. I do not expect doctors to be able to do much about pain.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

o. The thought of pain makes me afraid.....

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
--------------------------	--------------------------	--------------------------	--------------------------	--------------------------

### **Part 9 - About you**

Here are some general questions about yourself.

Please follow the instructions and answer ALL of the following questions.

1. What is your date of birth?  /  /

(For example - if you were born on the 5th June 1936, this would be entered as 05/06/36)

3. When you are ill, do you usually go straight away to the doctor, or do you wait to see if you get better? **(Please put a cross in one box only)**

- a. Go straight to the doctor..... ☐
- b. Wait a day or two to see if it gets better..... ☐
- c. Wait several days to see if it gets better..... ☐
- d. Put it off as long as possible..... ☐

### Section 3

We are interested in your views of health and life. Please indicate how much you agree or disagree with the following statements about health issues by putting a cross in one box for each statement.

**(Please put a cross in one box on each line)**

- |   | Strongly disagree        | Disagree                 | Neither agree nor disagree | Agree                    | Strongly agree           |
|---|--------------------------|--------------------------|----------------------------|--------------------------|--------------------------|
| a. There is a lot which I can do to control my health.....      | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| b. I do will affect whether my health gets better or worse..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| c. Treatments are effective in controlling illness.....         | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| d. My health is very unpredictable.....                         | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| e. Illness makes me feel afraid.....                            | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| f. The course of my life depends on me.....                     | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| g. I have the power to influence what happens in my life.....   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |
| h. Osteoarthritis is a serious condition.....                   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/> | <input type="checkbox"/> |

9. I get a sort of frightened feeling like 'butterflies' in my stomach:

- |                          |                          |                          |                          |
|--------------------------|--------------------------|--------------------------|--------------------------|
| Not at all               | Occasionally             | Quite often              | Very often               |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

10. I have lost interest in my appearance:

- |                          |                                       |                                   |                                  |
|--------------------------|---------------------------------------|-----------------------------------|----------------------------------|
| Definitely               | I don't take as much care as I should | I may not take quite as much care | I take just as much care as ever |
| <input type="checkbox"/> | <input type="checkbox"/>              | <input type="checkbox"/>          | <input type="checkbox"/>         |

11. I feel restless as if I have to be on the move:

- |                          |                          |                          |                          |
|--------------------------|--------------------------|--------------------------|--------------------------|
| Very much indeed         | Quite a lot              | Not very much            | Not at all               |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

12. I look forward with enjoyment to things:

- |                          |                            |                                |                          |
|--------------------------|----------------------------|--------------------------------|--------------------------|
| As much as I ever did    | Rather less than I used to | Definitely less than I used to | Hardly at all            |
| <input type="checkbox"/> | <input type="checkbox"/>   | <input type="checkbox"/>       | <input type="checkbox"/> |

13. I get sudden feelings of panic:

- |                          |                          |                          |                          |
|--------------------------|--------------------------|--------------------------|--------------------------|
| Very often indeed        | Quite often              | Not very often           | Not at all               |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

14. I can enjoy a good book or radio or television programme:

- |                          |                          |                          |                          |
|--------------------------|--------------------------|--------------------------|--------------------------|
| Often                    | Sometimes                | Not often                | Very seldom              |
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

## Part 8- Your views

### Section 1

We would like to know what your views are on **what causes illness**.  
Please indicate your agreement or disagreement with each of the following as  
a possible cause of illness.

**(Please put a cross in one box on each line)**

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
a. Stress or worry.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Hereditary - it runs in the family.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. A germ or virus.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Diet or eating habit.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Chance or bad luck.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Poor medical care in the past.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Pollution in the environment.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Someone's own behaviour.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Someone's mental attitude (e.g. thinking about life negatively).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Family problems or worries.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Overwork.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
l. Someone's emotional state (e.g. feeling down, lonely, anxious, empty).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
m. Ageing.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
n. Alcohol.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
o. Smoking.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
p. Accident or injury.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
q. Someone's personality.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
r. Altered immunity (the body is unable to fight illness).....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

### Section 2

We are also interested in how you think illness can be controlled and how  
you may use your doctor (GP).

1. In your opinion, is it a matter of luck whether you are well or ill, or is  
it something which can be controlled?

**(Please put a cross in one box only)**

All luck	Mostly luck	Bit of both	Mostly under control	Almost all under control
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. How often do you visit the doctor (GP) for yourself?


**(Please put a cross in one box only)**

Very often	Often	Occasionally	Seldom	Hardly ever
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## **Appendix 2: Health Questionnaire 3 years Follow-up**

# Health Questionnaire

**3 Year Follow-up**



LREC Number: 05/Q2604/20

Version 2, Date: 07/02/2005





## SECTION D CONTINUING TO HELP WITH THIS STUDY

Thank you very much for completing this questionnaire.

There will be further stages to this study, and we hope that people who have taken part so far will be able to help us again.

We may want to contact you again as part of this study and we are asking your permission to do this. Giving us permission to contact you again does not mean that you must take part.

Please put a cross in one of the boxes below to tell us if you are happy to be contacted again, and then sign and date where shown.

Yes, I am happy to be contacted again..... ☐

No, I do not want to be contacted again..... ☐

Please return this questionnaire whether or not you want to be contacted again.

Signature..... Date.....

Please print your name and address - .....

.....

.....


.....

**Please check that you have answered all the questions.**

**Please return this questionnaire in the envelope provided (no stamp needed).**

**If you have any questions, telephone Rosie Lacey on 01782 583921.**

**Thank you for your help with this research project**



5024

## Instructions for this questionnaire

The aim of this questionnaire is to find out if your health, including any pains or other symptoms, has changed in the last three years. The answers you give in the questionnaire will be treated in the **strictest confidence**.

**Please answer even if you have no problems with your health**

Please answer all of the questions unless the instructions ask you to do something else.

Please write in **BLOCK CAPITALS** where appropriate.

Most of the questions can be answered by putting a cross in a box like this:


**X**

When you have finished please check that you have answered all of the questions and then return the questionnaire in the envelope enclosed. You do not need a stamp. Please return the questionnaire in the next two weeks.

More details about this project are available in the information sheet enclosed.

If you would like further information about this project please contact Rosie Lacey, survey co-ordinator, on 01782 583921.

**Thank you for your help with this research project**



## **SECTION A YOUR GENERAL HEALTH**

This section is made up of questions about your health, the activities you do, and some of the ways in which people do things in everyday life.

Please answer each set of questions as the instructions tell you to.

## **SECTION D CONTINUING TO HELP WITH THIS STUDY**

# **For everyone to complete**

Please turn over the page to complete  
the questions in Section D



**Part 1 - Your health**

**Please answer every question.** Some questions may look similar to others but each one is different. Please take the time to read and answer each question carefully by placing a cross in the box of your choice.

1. In **general** would you say your health is:  
**(Please put a cross in one box only)**

Excellent	Very good	Good	Fair	Poor
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. **Compared to one year ago**, how would you rate your health in general now?  
**(Please put a cross in one box only)**

Much better than a year ago	Somewhat better than a year ago	About the same	Somewhat worse now than a year ago	Much worse now than a year ago
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. How TRUE or FALSE is each of the following statements for you?  
**(Please put a cross in one box on each line)**

	Definitely true	Mostly true	Don't know	Mostly false	Definitely false
a. I seem to get ill more easily than other people.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. I am as healthy as anybody I know.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. I expect my health to get worse...	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. My health is excellent.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



4. The following questions are about activities you might do during a typical day. Does your health now limit you in these activities? If so, how much?  
*(Please put a cross in one box on each line)*

	Yes, limited a lot	Yes, limited a little	No, not limited at all
a. <b>Vigorous activities</b> , such as running, lifting heavy objects, participating in strenuous sports.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. <b>Moderate activities</b> , such as moving a table, pushing a vacuum cleaner, bowling or playing golf.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Lifting or carrying groceries.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Climbing <b>several</b> flights of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Climbing <b>one</b> flight of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Bending, kneeling or stooping.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Walking <b>more than a mile</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Walking <b>half a mile</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Walking <b>one hundred yards</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Bathing and dressing yourself.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. During the **past 4 weeks**, have you had any of the following problems with your work or other regular daily activities as a result of your physical health?

*(Please put a cross in one box on each line)*

	Yes	No
a. Cut down on the <b>amount of time</b> you spent on work or other activities.....	<input type="checkbox"/>	<input type="checkbox"/>
b. <b>Accomplished less</b> than you would like.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Were limited in the <b>kind</b> of work or other activities .....	<input type="checkbox"/>	<input type="checkbox"/>
d. Had <b>difficulty</b> performing the work or other activities (for example, it took more effort).....	<input type="checkbox"/>	<input type="checkbox"/>

7. Is there any one special person you know that you feel very close to; someone you feel you can share confidences and feelings with?

Yes.... ☐

No.... ☐

If yes, how often do you see or talk with this person?

Daily	Weekly	Monthly	Several times a year	Once a year or less
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. How often do you go to religious meetings or services?

More than once a week	Once a week	1 to 3 times per month	Less than once per month	Never or almost never
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. How many hours **each week** do you participate in any groups such as social or work group, church-connected group, self-help group, charity, public service or community group?

None	1 to 2 hours	3 to 5 hours	6 to 10 hours	11 to 15 hours	16 or more hours
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Thank you for completing Section C**

**Please continue with Section D  
on page 29**

Part 2 - Friends and family

We are interested in the contact you may have with your friends and family.  
(Please answer each question and put a cross in one box on each line.)

1. How many living children do you have?

None                      1 to 2                      3 to 5                      6 or more

☐                      ☐                      ☐                      ☐

2. How many of your children do you see at least once a month?

None                      1 to 2                      3 to 5                      6 or more

☐                      ☐                      ☐                      ☐

3. Apart from your children, how many relatives do you have with whom you feel close?

None                      1 to 2                      3 to 5                      6 to 9                      10 or more

☐                      ☐                      ☐                      ☐                      ☐

4. How many close relatives do you see at least once a month?

None                      1 to 2                      3 to 5                      6 to 9                      10 or more

☐                      ☐                      ☐                      ☐                      ☐

5. How many close friends do you have?

None                      1 to 2                      3 to 5                      6 to 9                      10 or more

☐                      ☐                      ☐                      ☐                      ☐

6. How many of these friends do you see at least once a month?

None                      1 to 2                      3 to 5                      6 to 9                      10 or more

☐                      ☐                      ☐                      ☐                      ☐

6. During the **past 4 weeks**, have you had any of the following problems with your work or other regular daily activities as a result of any emotional problems (such as feeling depressed or anxious)?  
(Please put a cross in one box on each line)

	Yes	No
a) Cut down on the <b>amount of time</b> you spent on work or other activities.....	<input type="checkbox"/>	<input type="checkbox"/>
b) <b>Accomplished less</b> than you would like.....	<input type="checkbox"/>	<input type="checkbox"/>
c) Didn't do work or other activities as <b>carefully</b> as usual.....	<input type="checkbox"/>	<input type="checkbox"/>

7. During the **past 4 weeks**, to what extent have your physical health or emotional problems interfered with your normal social activities with family, friends, neighbours, or groups?  
(Please put a cross in one box only)

Not at all                      Slightly                      Moderately                      Quite a bit                      Extremely

☐                      ☐                      ☐                      ☐                      ☐

8. How much bodily pain have you had during the **past 4 weeks**?  
(Please put a cross in one box only)


None                      Very mild                      Mild                      Moderate                      Severe                      Very severe

☐                      ☐                      ☐                      ☐                      ☐                      ☐

9. During the **past 4 weeks**, how much did pain interfere with your normal work (including both work outside the home and housework)?  
(Please put a cross in one box only)

Not at all                      A little bit                      Moderately                      Quite a bit                      Extremely

☐                      ☐                      ☐                      ☐                      ☐



10. These questions are about how you feel and how things have been with you during the **past 4 weeks**. For each question, please give the one answer that comes closest to the way you have been feeling.

*(Please put a cross in one box on each line)*

How much of the time during the past 4 weeks.....	All of the time	Most of the time	A good bit of time	Some of the time	A little of the time	None of the time
a. Did you feel full of life?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Have you been a very nervous person?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Have felt so down in the dumps that nothing could cheer you up?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Have you felt calm and peaceful?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Did you have a lot of energy?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Have you felt downhearted and low?...	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Did you feel worn out?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Have you been a happy person?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Did you feel tired?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

11. During the **past 4 weeks**, how much of the time has your physical health or emotional problems interfered with your social activities (like visiting with friends, relatives, etc.)?

*(Please put a cross in one box only)*

All of the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. If working, what is your job title (examples - factory worker, welder, office worker, shop assistant, lawyer) ?

.....

If you are not working, or are retired, what was your last job title?

.....

7. Do you have a spouse or partner who is currently living with you?

Yes.... ☐ No.... ☐

If they are working, what is their job title (examples - factory worker, welder, office worker, shop assistant, lawyer)?

.....

If they are not working or are retired, what was their last job title?

.....

If you are a widow, or widower, what was your spouse's last job title?

.....

8. Thinking about the cost of living as it affects you, which of these descriptions best describes your situation:

*(Please put a cross in one box only)*

Find it a strain to get by from week to week.....	<input type="checkbox"/>
Have to be careful with money.....	<input type="checkbox"/>
Able to manage without much difficulty.....	<input type="checkbox"/>
Quite comfortably off.....	<input type="checkbox"/>

Part 1 - About you

1. What is your date of birth?      /   /

(For example - if you were born on the 5th of June 1936, this would be entered as 05/06/36)

2. Are you :

Female..... ☐

Male..... ☐

3. What is your current marital status?  
(Please put a cross in one box only)

Married..... ☐

Widowed..... ☐

Separated..... ☐

Cohabiting..... ☐

Divorced..... ☐

Single..... ☐

4. Do you live alone?

Yes..... ☐

No..... ☐

5. What is your current employment status?  
(Please put a cross in one box only)

Employed..... ☐

Not working due to ill health..... ☐

Retired..... ☐

Unemployed/seeking work..... ☐

Housewife..... ☐

Other..... ☐

Part 2 - Taking part

We are interested in some of the ways that are necessary for you to live your life in the way you choose. We are particularly interested in how often these things are achieved in the way you would like.

When answering the questions, please think about the **past 4 weeks**. It does not matter if you require the help of other people or from gadgets and machines. We would simply like to know if the activity **IS** achieved to the extent that you want it to be.

Please read each statement below and put a cross in the box which comes closest to how much you agree with the statement. **Please put a cross in one box only for each line.**

1. During the past 4 weeks, I have moved around in my home, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. During the past 4 weeks, I have moved around outside my home, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. During the past 4 weeks, my self-care needs (examples are washing, toileting, dressing, feeding, maintaining health) have been met, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>





4. During the past 4 weeks, my home has been looked after, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. During the past 4 weeks, my things (belongings) have been looked after, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. Do you have any relatives, or other people, who depend on you?

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, were these people looked after, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. During the past 4 weeks, I have met and spoken to other people, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. During the past 4 weeks, I, or someone else on my behalf, have managed my money, **as I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## SECTION C ABOUT YOU

This section contains general questions about yourself and your circumstances.

Please follow the instructions and answer ALL of the following questions.

## Part 4 - Coping with pain

This question is about how you **cope** with **pain**.

Below is a list of things that people have reported doing when they feel pain. Please cross a box for each statement to indicate **how often** you use that way of **cop**ing **when you are in pain** (where 0 is you "never do that", and 6 is you "always do that" when you are experiencing pain).

*(Please cross a box on each line)*

When I have pain.....	I never do that 0	1	2	3	4	5	I always do that 6
a. I think of things I enjoy.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. I just think of it as some other sensation, such as numbness...	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. It is terrible and I feel it is never going to get any better.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. I don't pay any attention to it.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. I pray for the pain to stop.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. I tell myself I can't let the pain stand in the way of what I have to do.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. I do something active, like household chores or projects....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Other (please state).....							

**Thank you for completing Section B**

**Please continue with Section C**

9. Do you choose to take part in paid or voluntary work?

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in paid or voluntary work, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Do you choose to take part in education or training courses?

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in education or training, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

11. Do you choose to take part in social activities?

(Examples of social activities are community or religious activities, meeting up with friends, going to clubs)

Yes.... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in social activities, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



### Part 3 - How you have felt in the past week

The next questions are about how you feel at the moment. Please read each item and put a cross next to the reply that comes closest to how you have been feeling **in the past week**. Don't take too long over your replies; your immediate reaction to each item will usually be more accurate than a long thought out response.

1. I feel tense or 'wound up':

Most of the time	A lot of the time	From time to time, occasionally	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. I still enjoy the things I used to enjoy:

Definitely as much	Not quite as much	Only a little	Hardly at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. I get a sort of frightened feeling as if something awful is about to happen:

Very definitely and quite badly	Yes, but not too badly	A little, but it doesn't worry me	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. I can laugh and see the funny side of things:

As much as I always could	Not quite so much now	Definitely not so much now	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Worrying thoughts go through my mind:

A great deal of the time	A lot of the time	Not too often	Very little
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. I feel cheerful:

Never	Not often	Sometimes	Most of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

### Part 3 - Body Chart

This question is about **recent pain** you may have had in **any part of your body**. By pain we also mean ache, discomfort or stiffness. Please **do not** include pain due to a feverish illness such as flu. If you are a woman please **do not** include pain related to your monthly period.

1. In the **past 4 weeks**, have you had pain that has lasted for **one day or longer** in **any part of your body**?


(Please put a cross in one box only)

Yes .... ☐

Please shade in the diagram below **any pain** that has lasted for **one day or longer in the past 4 weeks**

No .... ☐

Please continue with **Part 4**.



On this page we want you to think about **HOW INTENSE (how severe)** the pain in different parts of your body has been in the **last 6 months**.

2. In the **past 6 months, on average, how intense were each of these pains** rated on a 0-10 scale where 0 is "no pain" and 10 is "pain as bad as could be"? (That is, your usual pain at times you were experiencing pain.)

**For each pain, please put a cross in one box. For pains that do not apply to you please put a cross in box 0 to indicate No Pain.**

	No Pain 0	1	2	3	4	5	6	7	8	Pain as bad as could be 9 10
a. Headache	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Neck pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Shoulder pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Elbow pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Hand pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Chest pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Abdominal pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Back pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Hip pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Knee pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Foot pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. I can sit at ease and feel relaxed:

Definitely	Usually	Not often	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. I feel as if I am slowed down:

Nearly all the time	Very often	Sometimes	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. I get a sort of frightened feeling like 'butterflies' in my stomach:

Not at all	Occasionally	Quite often	Very often
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. I have lost interest in my appearance:

Definitely	I don't take as much care as I should	I may not take quite as much care	I take just as much care as ever
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

11. I feel restless as if I have to be on the move:

Very much indeed	Quite a lot	Not very much	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. I look forward with enjoyment to things:

As much as I ever did	Rather less than I used to	Definitely less than I used to	Hardly at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13. I get sudden feelings of panic:

Very often indeed	Quite often	Not very often	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. I can enjoy a good book or radio or television programme:

Often	Sometimes	Not often	Very seldom
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 4 - Specific health problems

We would like to know about some specific health problems.

1. Have you **ever** fractured your hip or your wrist?

Yes... ☐

No... ☐

2. Do you suffer from any of the following?

*(Please place a cross in the box for any which apply to you)*

a. Chest problems..... ☐

b. Heart problems..... ☐

c. Deafness..... ☐

d. Problems with eyesight  
(excluding the need for glasses)..... ☐

e. Raised blood pressure..... ☐

f. Diabetes..... ☐

g. Stroke..... ☐

h. Cancer..... ☐

i. Liver disease..... ☐

j. Kidney disease..... ☐

k. Circulation problems in the legs..... ☐

3. What is your weight?   stones   lbs or    kgs

4. What is your height?   feet   inches or    cms

## Part 2 - Pain in different parts of the body in the last 6 months

This is about pain in different parts of your body in the last 6 months. Please complete each of the following questions **even if you have not suffered pain in any of these areas** in the last 6 months.

The first question is about **HOW OFTEN** you have had pain in different parts of the body in the **last 6 months**.

1. On how many days in the **last 6 months** have you had.....?

*(For each pain, please put a cross in one box)*

	No Days	1 to 30 Days	31 to 89 Days	90 or more Days
a. Headache	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Neck pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Shoulder pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Elbow pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Hand pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Chest pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Abdominal pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Back pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Hip pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Knee pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Foot pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 1 - Specific joint pains and problems in the last year**

We are interested to know how many people suffer from specific joint pains and problems. Please answer each of the following questions about your hands, hips, knees, and feet, **even if you do not suffer from any problems**.

We would like you to tell us if you have had **any problems** with your **hands**. 'Hand problems' relate to **any hand symptoms** you may have experienced; for example, **pain or stiffness**, or any **knobbly swellings** on your hands, including your fingers and thumbs.

1. Have you had any problems with your **HANDS**, including your fingers and thumbs, over the **last year**?  
*(Please put a cross in one box only)*

Yes.... ☐ No.... ☐

2. Have you had any pain in your **HANDS**, including in your fingers and thumbs, over the **last year**?  
*(Please put a cross in one box only)*

Yes.... ☐ No.... ☐

We are also interested in any pain you may have had in your legs.

3. Have you had pain in the last year in and **around the HIP**?  
*(Please put a cross in one box only)*

Yes.... ☐ No.... ☐

4. Have you had pain in the last year in and **around the KNEE**?  
*(Please put a cross in one box only)*

Yes.... ☐ No.... ☐

5. Have you had pain in the last year in and **around the FOOT**?  
*(Please put a cross in one box only)*

Yes.... ☐ No.... ☐

5. Thinking back over the **past 3 months**, have you suffered from any of the following?  
*(Please place a cross in the box for any which apply to you)*

- a. A fall or falls..... ☐
- b. Difficulty remembering things..... ☐
- c. Cough with spit..... ☐
- d. Breathless when walking..... ☐
- e. Dizziness or unsteadiness..... ☐
- f. Weakness in an arm or leg..... ☐
- g. Shaking in your hands..... ☐
- h. Swelling in your ankles or feet..... ☐

6. Thinking over the **past 4 weeks**, did you?

*(Please put a cross in one box on each line)*

	Not at all	On some nights	On most nights
a. Have trouble falling asleep.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Wake up several times per night.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Have trouble staying asleep.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Wake up after your usual amount of sleep feeling tired and worn out.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



7. Please put a cross in the box to show whether you agree (yes box) or disagree (no box) with each of the following statements.

*(Please put a cross in one box on each line)*

	Yes	No
a. I am confused and start to do more than one thing at a time.....	<input type="checkbox"/>	<input type="checkbox"/>
b. I have more minor accidents than usual (e.g. I drop things, I trip and fall, or bump into things).....	<input type="checkbox"/>	<input type="checkbox"/>
c. I react slowly to things that are said or done.....	<input type="checkbox"/>	<input type="checkbox"/>
d. I do not finish things that I start.....	<input type="checkbox"/>	<input type="checkbox"/>
e. I have difficulty reasoning and solving problems (e.g. making plans, making decisions, or learning new things)..	<input type="checkbox"/>	<input type="checkbox"/>
f. I sometimes get confused (e.g. I do not know where I am, who is around, or what day it is).....	<input type="checkbox"/>	<input type="checkbox"/>
g. I forget a lot (e.g. things that happened recently, where I put things, or to keep appointments).....	<input type="checkbox"/>	<input type="checkbox"/>
h. I do not keep my attention on any activity for long.....	<input type="checkbox"/>	<input type="checkbox"/>
i. I make mistakes more than usual.....	<input type="checkbox"/>	<input type="checkbox"/>
j. I have difficulty doing things which involve thought and concentration.....	<input type="checkbox"/>	<input type="checkbox"/>

**Thank you for completing Section A**

**Please continue with Section B**

## SECTION B PAINS AND JOINT PROBLEMS

This section is about any pains you may have or any problems with your joints.

Please fill in all the parts even if you do not suffer from any pains or joint problems.

Please follow the instructions for each part.

### **Appendix 3: Health Questionnaire 6 years Follow-up**




# Health Questionnaire

**6 Year Follow-up**

REC Number: 05/Q2604/20

Version 3, Date: 06/02/2008



16993

## SECTION D: CONTINUING TO HELP WITH THIS STUDY

Thank you very much for completing this questionnaire.

There will be further stages to this study, and we hope that people who have taken part so far will be able to help us again.

We may want to contact you again as part of this study and we are asking your permission to do this. Giving us permission to contact you again does not mean that you must take part.

Please put a cross in one of the boxes below to tell us if you are happy to be contacted again, and then sign and date where shown.

Yes, I am happy to be contacted again..... ☐

No, I do not want to be contacted again..... ☐

Please return this questionnaire whether or not you want to be contacted again.

Signature..... Date.....

Please print your **name** and **address** - .....

.....

.....

.....

**Please check that you have answered all the questions.**

**Please return this questionnaire in the envelope provided (no stamp needed).**

**If you have any questions, telephone Rosie Lacey on 01782 584721.**

**Thank you for your help with this research project**

## Instructions for this questionnaire

The aim of this questionnaire is to find out if your health, including any pains or other symptoms, has changed in the last three years. The answers you give in the questionnaire will be treated in the **strictest confidence**.

**Please answer even if you have no problems with your health**

Please answer all of the questions unless the instructions ask you to do something else.

Please write in **BLOCK CAPITALS** where appropriate.

Most of the questions can be answered by putting a cross in a box like this:


☐

When you have finished please check that you have answered all of the questions and then return the questionnaire in the envelope enclosed. You do not need a stamp. Please return the questionnaire in the next two weeks.

More details about this project are available in the information sheet enclosed.

If you would like further information about this project please contact Rosie Lacey, survey co-ordinator, on 01782 584721.

**Thank you for your help with this research project**



## SECTION A: YOUR GENERAL HEALTH

This section is made up of questions about your health, the activities you do, and some of the ways in which people do things in everyday life.

Please answer each set of questions as the instructions tell you to.

## SECTION D: CONTINUING TO HELP WITH THIS STUDY

# For everyone to complete

Please turn over the page to complete the questions in Section D



32140

14. How satisfied are you with being retired?  
(Please put a cross in one box only)

Very satisfied	Satisfied	Neither satisfied nor dissatisfied	Not very satisfied	Not at all satisfied
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

15. Did you have the opportunity to work past retirement age?  
(Please put a cross in one box only)

Yes.....☐

No.....☐

**Thank you for completing Section C**

**Please continue with Section D on page 45**

## Part 1 - Your health

**Please answer every question.** Some questions may look similar to others but each one is different. Please take the time to read and answer each question carefully by placing a cross in the box of your choice.

1. In **general** would you say your health is:  
(Please put a cross in one box only)

Excellent	Very good	Good	Fair	Poor
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. **Compared to one year ago**, how would you rate your health in general now?

(Please put a cross in one box only)

Much better than a year ago	Somewhat better than a year ago	About the same	Somewhat worse now than a year ago	Much worse now than a year ago
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. How TRUE or FALSE is each of the following statements for you?  
(Please put a cross in one box on each line)

	Definitely true	Mostly true	Don't know	Mostly false	Definitely false
a. I seem to get ill more easily than other people.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. I am as healthy as anybody I know.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. I expect my health to get worse.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. My health is excellent.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



4. The following questions are about activities you might do during a typical day. Does your health now limit you in these activities? If so, how much?  
*(Please put a cross in one box on each line)*

	Yes, limited a lot	Yes, limited a little	No, not limited at all
a. <b>Vigorous activities</b> , such as running, lifting heavy objects, participating in strenuous sports.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. <b>Moderate activities</b> , such as moving a table, pushing a vacuum cleaner, bowling or playing golf.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Lifting or carrying groceries.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Climbing <b>several</b> flights of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Climbing <b>one</b> flight of stairs.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Bending, kneeling or stooping.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Walking <b>more than a mile</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Walking <b>half a mile</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Walking <b>one hundred yards</b> .....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Bathing and dressing yourself.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. During the **past 4 weeks**, have you had any of the following problems with your work or other regular daily activities as a result of your physical health?

<i>(Please put a cross in one box on each line)</i>	Yes	No
a. Cut down on the <b>amount of time</b> you spent on work or other activities.....	<input type="checkbox"/>	<input type="checkbox"/>
b. <b>Accomplished less</b> than you would like.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Were limited in the <b>kind</b> of work or other activities.....	<input type="checkbox"/>	<input type="checkbox"/>
d. Had <b>difficulty</b> performing the work or other activities (for example, it took more effort).....	<input type="checkbox"/>	<input type="checkbox"/>

12. Which of the following aspects of your last job before you retired would have enabled you to continue working in your last job?  
*(Please put a cross in the box for any that apply to you)*

a. Less physical demand.....	<input type="checkbox"/>
b. Less mental demand.....	<input type="checkbox"/>
c. Flexible working (e.g. flexible hours, job sharing, work at home).....	<input type="checkbox"/>
d. Special equipment or workplace adaptation.....	<input type="checkbox"/>
e. Being allowed to change how I did my job.....	<input type="checkbox"/>
f. Retraining and skill development.....	<input type="checkbox"/>
g. More support from people in my workplace.....	<input type="checkbox"/>
h. Better health facilities in my place of work.....	<input type="checkbox"/>

13. In the final 12 months of your last job before you retired how many days were you off work because of **joint pain**?  
*(Please put a cross in one box only)*

I was not off work at all due to joint pain.....	<input type="checkbox"/>
Less than 7 days.....	<input type="checkbox"/>
1-4 weeks.....	<input type="checkbox"/>
More than 1 month, but less than 3 months.....	<input type="checkbox"/>
More than 3 months.....	<input type="checkbox"/>

Please complete questions 10 to 15 ONLY if you are fully retired.

10. At what age did you retire?  years

11. What were your reasons for retiring?  
(Please put a cross in the box for any that apply to you)

- a. Joint pain..... ☐
- b. Ill health or disability other than joint pain..... ☐
- c. Job was too tiring/stressful..... ☐
- d. Ill health or disability of a relative/friend..... ☐
- e. Made redundant/dismissed/had no choice..... ☐
- f. Offered reasonable financial terms to retire early or took voluntary redundancy..... ☐
- g. Could not find another job..... ☐
- h. Dissatisfied or fed up with job..... ☐
- i. Lack of flexibility in job..... ☐
- j. Lack of support at work for people with health problems..... ☐
- k. To spend more time with/look after partner/family..... ☐
- l. To enjoy life while still fit and young..... ☐
- m. Reached retirement age (65 years for men, 60 years for women)..... ☐
- n. Because I could afford to..... ☐
- o. To retire at the same time as husband/wife/partner..... ☐
- p. To retire at a different time to husband/wife/partner..... ☐

6. During the **past 4 weeks**, have you had any of the following problems with your work or other regular daily activities as a result of any emotional problems (such as feeling depressed or anxious)?  
(Please put a cross in one box on each line)

- |   | Yes                      | No                       |
|---|--------------------------|--------------------------|
| a) Cut down on the <b>amount of time</b> you spent on work or other activities..... | <input type="checkbox"/> | <input type="checkbox"/> |
| b) <b>Accomplished less</b> than you would like.....                                | <input type="checkbox"/> | <input type="checkbox"/> |
| c) Didn't do work or other activities as <b>carefully</b> as usual.....             | <input type="checkbox"/> | <input type="checkbox"/> |

7. During the **past 4 weeks**, to what extent have your physical health or emotional problems interfered with your normal social activities with family, friends, neighbours, or groups?  
(Please put a cross in one box only)

- | Not at all               | Slightly                 | Moderately               | Quite a bit              | Extremely                |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

8. How much bodily pain have you had during the **past 4 weeks**?  
(Please put a cross in one box only)

- | None                     | Very mild                | Mild                     | Moderate                 | Severe                   | Very severe              |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

9. During the **past 4 weeks**, how much did pain interfere with your normal work (including both work outside the home and housework)?  
(Please put a cross in one box only)

- | Not at all               | A little bit             | Moderately               | Quite a bit              | Extremely                |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |



10. These questions are about how you feel and how things have been with you during the **past 4 weeks**. For each question, please give the one answer that comes closest to the way you have been feeling.  
*(Please put a cross in one box on each line)*

How much of the time during the past 4 weeks.....	All of the time	Most of the time	A good bit of time	Some of the time	A little of the time	None of the time
a. Did you feel full of life?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Have you been a very nervous person?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Have you felt so down in the dumps that nothing could cheer you up?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Have you felt calm and peaceful?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Did you have a lot of energy?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Have you felt downhearted and low?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Did you feel worn out?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Have you been a happy person?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Did you feel tired?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

11. During the **past 4 weeks**, how much of the time has your physical health or emotional problems interfered with your social activities (like visiting with friends, relatives, etc.)?  
*(Please put a cross in one box only)*

All of the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Please complete questions 8 and 9 **ONLY** if you have **been off work due to ill health for 6 months or more, OR if you are unemployed**.

8. We are interested to know if joint pain or ill health has played a part in your current employment status.

What are the reasons for you being **off work due to ill health for six months or more, or for being unemployed**?

*(Please put a cross in one box on each line)*

	Yes	No
a. Joint pain.....	<input type="checkbox"/>	<input type="checkbox"/>
b. Ill health or disability other than joint pain.....	<input type="checkbox"/>	<input type="checkbox"/>

9. We are interested in knowing whether certain aspects of work will make it easier for people "who have been off work due to ill health for six months or more or unemployed" to return to work. Which of the following **aspects of a job** would make it easier for you to return to work?  
*(Please put a cross in the box for any that apply to you)*

a. Fewer physical demands.....	<input type="checkbox"/>
b. Fewer mental demands.....	<input type="checkbox"/>
c. Flexible working (e.g. flexible hours, job sharing, work at home).....	<input type="checkbox"/>
d. Special equipment or workplace adaptation.....	<input type="checkbox"/>
e. Being allowed to change how you do the job.....	<input type="checkbox"/>
f. Retraining and skill development.....	<input type="checkbox"/>
g. More support for people with health problems at work.....	<input type="checkbox"/>
h. Better health facilities in the workplace.....	<input type="checkbox"/>

PLEASE NOW GO TO SECTION D, ON PAGE 45



6. How satisfied are you with your **current** job?  
(Please put a cross in one box only)

Very satisfied	Satisfied	No opinion	Dissatisfied	Very dissatisfied
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. Which of the following statements **best describes** the work that you do in your **current** job?  
(Please put a cross in one box only)

**Sedentary occupation** (e.g. secretary - where you spend most of your time sitting)..... ☐

**Standing occupation** (e.g. shop assistant, security guard - where you spend most of your time standing or walking, although the work does not require intense physical effort)..... ☐

**Physical work** (e.g. plumber, nurse - a job that involves some physical effort including handling of heavy objects and use of tools)..... ☐

**Heavy manual work** (e.g. miner or bricklayer - a job that involves very vigorous physical activity including handling very heavy objects)..... ☐

PLEASE NOW GO TO SECTION D, ON PAGE 45

**Part 2 - Taking part**

We are interested in some of the ways that are necessary for you to live your life in the way you choose. We are particularly interested in how often these things are achieved in the way you would like.

When answering the questions, please think about the **past 4 weeks**. It does not matter if you require the help of other people or from gadgets and machines. We would simply like to know if the activity **IS** achieved to the extent that you want it to be.

Please read each statement below and put a cross in the box which comes closest to how much you agree with the statement.  
(Please put a cross in one box on each line)

1. During the past 4 weeks, I have moved around my home, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. During the past 4 weeks, I have moved around outside my home, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. During the past 4 weeks, my self-care needs (examples are washing, toileting, dressing, feeding, maintaining health) have been met, **as and when I have wanted**.

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



4. During the past four weeks, my home has been looked after, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. During the past 4 weeks, my things (belongings) have been looked after, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. Do you have any relatives, or other people, who depend on you?

Yes.....☐ No....☐

If yes, during the past 4 weeks, were these people looked after, **as and when you have wanted?**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. During the past 4 weeks, I have met and spoken to other people, **as and when I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. During the past 4 weeks, I, or someone else on my behalf, have managed my money, **as I have wanted.**

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Please answer questions 4 to 7 **ONLY** if you are in full-time or part-time paid work, OR have been off work due to ill health for more than 1 week but less than 6 months.

4. Please think about the **last 30 days** that you were in work. For each question, please put a cross in **one box on each line** that best describes **this job**.

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
a. My work is physically demanding.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. I can control the way I work.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. My work is repetitive.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. My work is boring and monotonous.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. My work colleagues are supportive.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. I can work at home sometimes.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. My hours of work are flexible.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. I can use aids and appliances to help me do my job.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. I can adapt my work area to help me do my job.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. There are opportunities to retrain and develop my skills.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Health care facilities are good in my place of work.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. In general, over the **last 30 days that you were in work**, how would you rate your average performance at work on a 0-10 scale, where 0 is 'worst performance' and 10 is 'best performance'?  
**(Please circle one number only)**

Worst performance										Best performance				
0	1	2	3	4	5	6	7	8	9	10				

Now we want you to think back to your employment status and follow these instructions.

If you are currently in **full-time or part-time paid work**  
**OR** have been off work due to ill health for more than 1 week but less than 6 months



please fill in questions 4 to 7, on pages 39 and 40

If you have been off work due to ill health for 6 months or more  
**OR** are unemployed



please fill in questions 8 and 9, on page 41

If you are **fully retired**



please fill in questions 10 to 15, on pages 42 to 44

If you are a **housewife**  
**OR** not working due to being on a formal training course  
**OR** because you look after/care for dependants



please go to Section D, on page 45

9. Do you choose to take part in paid or voluntary work?

Yes..... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in paid or voluntary work, as and when you have wanted?

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Do you choose to take part in education or training courses?

Yes..... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in education or training, as and when you have wanted?

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

11. Do you choose to take part in social activities?  
(Examples of social activities are community or religious activities, meeting up with friends, going to clubs)

Yes..... ☐ No.... ☐

If yes, during the past 4 weeks, have you taken part in social activities, as and when you have wanted?

All the time	Most of the time	Some of the time	A little of the time	None of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 3 - How do you do things

We would like to know about how you do things in your home and if you require the help of other people or special devices. Please answer each question.

1. The following questions are about activities you might do when looking after yourself during a typical day.  
Does your health now limit you in these activities? If so, how much?  
(Please put a cross in one box on each line)

	Yes, limited a lot	Yes, limited a little	No, not limited at all
a. Washing and bathing.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Looking after your skin, teeth, hair and nails.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Putting on and taking off clothes.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Using a toilet.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Eating and drinking.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Preparing food.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Washing and ironing clothes.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Doing housework.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Getting up from a chair.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. We would like to know about the job that you have done for **most of your working life**. Please tell us:
- a. The **title** of the job you have done for **most of your working life**  
(e.g. assembly worker, lawyer, office manager, shop assistant, van driver):  
.....
- b. The **type of work** of the job you have done for **most of your working life**  
(e.g. banking, building, catering, farming, office, pottery, retail):  
.....
- c. The **year** you **started** the job you have done for **most of your working life**:  
 (For example, 1958)
- d. Are you **still working** in the job you have done for **most of your working life**?  
(Please put a cross in one box only)  
Yes..... ☐ No.... ☐
- e. If **no**, the **year** you **finished** the job you have done for **most of your working life**?  
 (For example, 2002)
3. Thinking about the cost of living as it affects you, which of these descriptions best describes your situation?  
(Please put a cross in one box only)
- Find it a strain to get by from week to week..... ☐
- Have to be careful with money..... ☐
- Able to manage without much difficulty..... ☐
- Quite comfortably off..... ☐

### Part 3 - Work

We are interested in whether people's ability to work is affected by ill health or certain aspects of their job.

#### 1. What is your current employment status?

**(Please put a cross in one box only)**

In full-time paid work..... ☐

In part-time paid work but not retired..... ☐

In part-time paid work and partly retired..... ☐

Off work due to ill health for more than 1 week  
but less than 6 months..... ☐

Off work due to ill health for 6 months or more..... ☐

Fully retired due to reaching retirement age..... ☐

Fully retired due to taking early retirement..... ☐

Fully retired due to taking early retirement  
because of ill health..... ☐

Not working due to looking after/caring for  
dependants or family..... ☐

Not working due to attending a full-time formal  
education or training course..... ☐

Unemployed or seeking work..... ☐

Housewife or looking after the home..... ☐

#### 2. During the **past 4 weeks** have you had help from anyone else, such as family (including your spouse), friends, a health professional (e.g. nurse, occupational therapist, physiotherapist) or someone from social services (e.g. home help, care worker), to do the following activities?

**(Please put a cross in the box for any that apply to you)**

	No, and I didn't need help	No, <b>but</b> I have needed help	Yes, I've had help from family or friends	Yes, I've had help from a health professional or from social services
a. Washing and bathing.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Looking after your skin, teeth, hair, nails.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Putting on and taking off clothes.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Using a toilet.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Eating and drinking.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Preparing food.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Washing and ironing clothes.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Doing housework.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Getting up from a chair.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Moving around outside the home/going places.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Sorting out bills and paperwork.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

#### 3. During the past 4 weeks, on average, how many **hours per week** did you receive help **from friends and family** (including your spouse)?

hours per week

(For example, 10 hours per week)

4. During the past 4 weeks, on average, how many **hours per week** did you receive help **from a health professional** (e.g. nurse) **and/or from social services** (e.g. home-help, care worker)?

hours per week

(For example, 10 hours per week)

5. During the past 4 weeks have you used any of the following aids and appliances to help with your daily activities?

*(Please put a cross in one box on each line)*

	Yes	No
a) A walking stick or frame.....	<input type="checkbox"/>	<input type="checkbox"/>
b) A wheelchair or electric scooter.....	<input type="checkbox"/>	<input type="checkbox"/>
c) Stair lift or through-floor lift.....	<input type="checkbox"/>	<input type="checkbox"/>
d) Extra hand-rails to get into your home or upstairs.....	<input type="checkbox"/>	<input type="checkbox"/>
e) Aids to help you wash or dress.....	<input type="checkbox"/>	<input type="checkbox"/>
f) Aids to help you with cooking or eating.....	<input type="checkbox"/>	<input type="checkbox"/>
g) Raised chair or bed/special stool.....	<input type="checkbox"/>	<input type="checkbox"/>
h) Raised toilet seat.....	<input type="checkbox"/>	<input type="checkbox"/>
i) Hand-rails in your bathroom/bath seat/shower seat.....	<input type="checkbox"/>	<input type="checkbox"/>
j) Modified bath or shower.....	<input type="checkbox"/>	<input type="checkbox"/>
k) Adapted phone (e.g. big buttons).....	<input type="checkbox"/>	<input type="checkbox"/>

7. Is there any one special person you know that you feel very close to; someone you feel you can share confidences and feelings with?

Yes.... ☐

No.... ☐

If yes, how often do you see or talk with this person?

Daily	Weekly	Monthly	Several times a year	Once a year or less
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

8. How often do you go to religious meetings or services?

More than once a week	Once a week	1 to 3 times per month	Less than once per month	Never or almost never
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. How many hours **each week** do you participate in any groups such as social or work group, church-connected group, self-help group, charity, public service or community group?

None	1 to 2 hours	3 to 5 hours	6 to 10 hours	11 to 15 hours	16 or more hours
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 2 - Friends and family

We are interested in the contact you may have with your friends and family. Please answer each question and put a cross in one box on each line.

1. How many living children do you have?

None	1 to 2	3 to 5	6 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. How many of your children do you see at least once a month?

None	1 to 2	3 to 5	6 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. Apart from your children, how many relatives do you have with whom you feel close?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. How many close relatives do you see at least once a month?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. How many close friends do you have?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. How many of these friends do you see at least once a month?

None	1 to 2	3 to 5	6 to 9	10 or more
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 4 - Access and environment

We would like you to tell us how easy it is for you to get hold of, or gain access to, things when you need them in your daily life, and about your neighbourhood.

1. When you need them, do you have access to.....  
(Please put a cross in one box on each line)

	Yes	No
a) A car.....	<input type="checkbox"/>	<input type="checkbox"/>
b) Public transport.....	<input type="checkbox"/>	<input type="checkbox"/>
c) Your doctor (GP).....	<input type="checkbox"/>	<input type="checkbox"/>
d) A chemist.....	<input type="checkbox"/>	<input type="checkbox"/>
e) Internet and e-mail.....	<input type="checkbox"/>	<input type="checkbox"/>

2. Which of the following best describes your home?  
(Please put a cross in one box only)

Terraced house.....	<input type="checkbox"/>	Flat/apartment.....	<input type="checkbox"/>
Semi-detached house....	<input type="checkbox"/>	Bungalow.....	<input type="checkbox"/>
Detached house.....	<input type="checkbox"/>	Residential/care home....	<input type="checkbox"/>

3. Thinking about your home, please indicate how much you agree or disagree with each of the following statements.  
(Please put a cross in one box on each line)

	Strongly disagree	Disagree	Neither disagree nor agree	Agree	Strongly agree
a. It is easy to get to my toilet or bathroom.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. It is easy to move around my toilet or bathroom.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. It is easy to get in and out of my home.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



4. Thinking about your local neighbourhood, please indicate how much you agree or disagree with each of the following statements.

**(Please put a cross in one box on each line)**

	Strongly disagree	Disagree	Neither disagree nor agree	Agree	Strongly agree
a. Hills or steep slopes are common in my neighbourhood.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Hills and steep slopes make it difficult to move around outside my home.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Public buildings and shops are difficult to move around in.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Lack of good pavements stops me from going out for a walk.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Lack of access to public parks or sports facilities stops me from going out.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Litter or vandalism are a problem in my area.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Pollution is a problem in my area.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Heavy traffic or speeding cars stop me going out.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. The weather stops me from going out.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. The local council are helpful in facilitating my daily activities.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. My fear of crime stops me going out.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
l. The views of my family prevent me from doing my usual activities.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
m. The attitudes and views of people in my neighbourhood prevent me from doing my usual activities.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 1 - About you

1. What is your date of birth?

/   /

(For example - if you were born on the 5th of June 1936, this would be entered as 05/06/36)

2. Are you : Female..... ☐

Male..... ☐

3. What is your current marital status?

**(Please put a cross in one box only)**

Married..... ☐

Widowed..... ☐

Separated..... ☐

Cohabiting..... ☐

Divorced..... ☐

Single..... ☐

4. Do you live alone?

Yes..... ☐

No..... ☐

5. What is your weight?

stones   lbs or    kgs

6. What is your height?

feet   inches or    cms

7. What is your current smoking status?

**(Please put a cross in one box only)**

Never smoked

Previously smoked

Currently smoking

☐

☐

☐

8. On average, how often do you drink alcohol?

**(Please put a cross in one box only)**

Daily or most days

Once or twice a week

Once or twice a month

Once or twice a year

Never


☐

☐

☐

☐

☐



32140



SECTION C: ABOUT YOU

This section contains general questions about yourself and your circumstances.

Please follow the instructions and answer ALL of the questions relevant to you.

Part 5 - How you have felt in the past week

The next questions are about how you feel at the moment. Please read each item and put a cross next to the reply that comes closest to how you have been feeling **in the past week**. Don't take too long over your replies; your immediate reaction to each item will usually be more accurate than a long thought out response.

1. I feel tense or 'wound up':

Most of the time	A lot of the time	From time to time, occasionally	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. I still enjoy the things I used to enjoy:

Definitely as much	Not quite as much	Only a little	Hardly at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. I get a sort of frightened feeling as if something awful is about to happen:

Very definitely and quite badly	Yes, but not too badly	A little, but it doesn't worry me	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. I can laugh and see the funny side of things:

As much as I always could	Not quite so much now	Definitely not so much now	Not at all
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Worrying thoughts go through my mind:

A great deal of the time	A lot of the time	Not too often	Very little
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

6. I feel cheerful:

Never	Not often	Sometimes	Most of the time
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. I can sit at ease and feel relaxed:

Definitely  
☐

Usually  
☐

Not often  
☐

Not at all  
☐

8. I feel as if I am slowed down:

Nearly all the time  
☐

Very often  
☐

Sometimes  
☐

Not at all  
☐

9. I get a sort of frightened feeling like 'butterflies' in my stomach:

Not at all  
☐

Occasionally  
☐

Quite often  
☐

Very often  
☐

10. I have lost interest in my appearance:

Definitely  
☐

I don't take  
as much care  
as I should  
☐

I may not take  
quite as much  
care  
☐

I take just as  
much care  
as ever  
☐

11. I feel restless as if I have to be on the move:

Very much indeed  
☐

Quite a lot  
☐

Not very much  
☐

Not at all  
☐

12. I look forward with enjoyment to things:

As much as  
I ever did  
☐

Rather less  
than I used to  
☐

Definitely less  
than I used to  
☐

Hardly at all  
☐

13. I get sudden feelings of panic:

Very often indeed  
☐

Quite often  
☐

Not very often  
☐

Not at all  
☐

14. I can enjoy a good book or radio or television programme:

Often  
☐

Sometimes  
☐

Not often  
☐

Very seldom  
☐

**Thank you for completing Section B**

**Please continue with Section C on page 32**

**Part 6 - Treatments for pain**

Which of the following treatments have you used in the last **12 months** because of **pain**?

*(Please put a cross in one box on each line)*

	Yes	No
a. Acupuncture.....	<input type="checkbox"/>	<input type="checkbox"/>
b. Aromatherapy.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Bandages or other types of support.....	<input type="checkbox"/>	<input type="checkbox"/>
d. Cold therapy (e.g. ice pack).....	<input type="checkbox"/>	<input type="checkbox"/>
e. Copper bracelets.....	<input type="checkbox"/>	<input type="checkbox"/>
f. Electrotherapy (e.g.TENS machine).....	<input type="checkbox"/>	<input type="checkbox"/>
g. Exercises.....	<input type="checkbox"/>	<input type="checkbox"/>
h. Heat therapy (e.g. heat pack or lamp).....	<input type="checkbox"/>	<input type="checkbox"/>
i. Injection into a joint.....	<input type="checkbox"/>	<input type="checkbox"/>
j. Magnets.....	<input type="checkbox"/>	<input type="checkbox"/>
k. Massage therapy.....	<input type="checkbox"/>	<input type="checkbox"/>
l. A surgical operation.....	<input type="checkbox"/>	<input type="checkbox"/>
m. Osteopathic manipulation.....	<input type="checkbox"/>	<input type="checkbox"/>
n. Physiotherapy.....	<input type="checkbox"/>	<input type="checkbox"/>
o. Occupational therapy.....	<input type="checkbox"/>	<input type="checkbox"/>

**Part 6 - Specific health problems**

We would like to know about some specific health problems.

1. Have you **ever** fractured or broken any of the following?  
*(Please put a cross in the box for any that apply to you)*

- Hip..... ☐
- Wrist..... ☐
- Other bones..... ☐

2. Do you suffer from any of the following?  
*(Please put a cross in the box for any that apply to you)*

- a. Chest problems..... ☐
- b. Heart problems..... ☐
- c. Deafness..... ☐
- d. Problems with eyesight  
(excluding the need for glasses)..... ☐
- e. Raised blood pressure..... ☐
- f. Diabetes..... ☐
- g. Stroke..... ☐
- h. Cancer..... ☐
- i. Liver disease..... ☐
- j. Kidney disease..... ☐
- k. Circulation problems in the legs..... ☐

3. Thinking back over **the past 3 months**, have you suffered from any of the following?

**(Please put a cross in the box for any that apply to you)**

- a. A fall or falls..... ☐
- b. Difficulty remembering things..... ☐
- c. Cough with spit..... ☐
- d. Breathless when walking..... ☐
- e. Dizziness or unsteadiness..... ☐
- f. Weakness in an arm or leg..... ☐
- g. Shaking in your hands..... ☐
- h. Swelling in your ankles or feet..... ☐

4. Thinking back over the **past 4 weeks**, did you?

**(Please put a cross in one box on each line)**

- |  | Not<br>at all            | On some<br>nights        | On most<br>nights        |
|--|--------------------------|--------------------------|--------------------------|
| a. Have trouble falling asleep.....  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| b. Wake up several times per night.....  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| c. Have trouble staying asleep.....  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| d. Wake up after your usual amount of<br>sleep feeling tired and worn out..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

5. What would you consider to be your two most important health problems **at the moment?**

- a. ....
- b. ....

## Part 5 - Medicines for pain

Please tell us about any tablets, pills or creams you may have used to reduce **pain** in the past **4 weeks**. For each medicine you have used, please put a cross to show if your doctor prescribed it, or if you bought it without a prescription. If you have used medicines prescribed by your doctor **and** which you have also bought without prescription, please put a cross in **both** boxes.

### TABLETS, PILLS

	Medicine prescribed by your doctor	Medicine bought <b>without</b> a prescription
a. Paracetamol (e.g. Panadol).....	<input type="checkbox"/>	<input type="checkbox"/>
b. Aspirin.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Ibuprofen tablets (e.g. Nurofen, Brufen).....	<input type="checkbox"/>	<input type="checkbox"/>
d. Naproxen (e.g. Naprosyn, Synflex).....	<input type="checkbox"/>	<input type="checkbox"/>
e. Diclofenac (e.g. Voltarol).....	<input type="checkbox"/>	<input type="checkbox"/>
f. Celecoxib (Celebrex), Etoricoxib (Arcoxia), Meloxicam (Mobic) or Lumiracoxib (Prexige).....	<input type="checkbox"/>	<input type="checkbox"/>
g. Tramacet.....	<input type="checkbox"/>	<input type="checkbox"/>
h. Co-proxamol (e.g. Distalgesic).....	<input type="checkbox"/>	<input type="checkbox"/>
i. Co-dydramol.....	<input type="checkbox"/>	<input type="checkbox"/>
j. Co-codamol (e.g. Solpadol, Kapake).....	<input type="checkbox"/>	<input type="checkbox"/>
k. Tramadol (e.g. Zydol, Zamadol).....	<input type="checkbox"/>	<input type="checkbox"/>
l. Dihydrocodeine (e.g. DF 118 Forte).....	<input type="checkbox"/>	<input type="checkbox"/>
m. Glucosamine.....	<input type="checkbox"/>	<input type="checkbox"/>
n. Chondroitin sulphate.....	<input type="checkbox"/>	<input type="checkbox"/>
o. Fish oil (e.g. cod liver oil).....	<input type="checkbox"/>	<input type="checkbox"/>
p. Herbal remedies (e.g. evening primrose oil)....	<input type="checkbox"/>	<input type="checkbox"/>

### CREAMS, GELS, RUB-ONS, SPRAYS

E.g. Emulgel, Feldene, Ibuleve, Movelat, Powergel ☐ ☐

## Part 4 - Coping with pain

This question is about how you **cope** with **pain**.


Below is a list of things that people have reported doing when they feel pain. Please cross a box for each statement to indicate **how often** you use that way of **cop**ing **when you are in pain** (where 0 is you "never do that", and 6 is you "always do that" when you are experiencing pain).

*(Please put a cross in one box on each line)*


When I have pain.....	I never do that 0	1	2	3	4	5	I always do that 6
a. I think of things I enjoy.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. I just think of it as some other sensation, such as numbness...	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. It is terrible and I feel it is never going to get any better.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. I don't pay any attention to it.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. I pray for the pain to stop.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. I tell myself I can't let the pain stand in the way of what I have to do.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. I do something active, like household chores or projects....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Other (please state).....							

6. We are interested in whether your big toes are straight or angled sideways because this might be related to your ability to move around.

First, please look at your left big toe whilst standing without shoes and socks on. Ignore the positioning and the gaps between your other toes and try to focus **only** on your big toe. Select from the first set of pictures below labelled from A to E which one best shows the angle of your left big toe. Please **circle** the letter of that picture.



Now do the same for your right big toe joint using the set of pictures below labelled from F to J. Again please **circle** the letter of the picture that best shows the angle of your right big toe.



Please put a cross in the box to show whether you agree (yes box) or disagree (no box) with each of the following statements.

(Please put a cross in one box on each line)

	Yes	No
a. I am confused and start to do more than one thing at a time.....	<input type="checkbox"/>	<input type="checkbox"/>
b. I have more minor accidents than usual (e.g. I drop things, I trip and fall, or bump into things).....	<input type="checkbox"/>	<input type="checkbox"/>
c. I react slowly to things that are said or done.....	<input type="checkbox"/>	<input type="checkbox"/>
d. I do not finish things that I start.....	<input type="checkbox"/>	<input type="checkbox"/>
e. I have difficulty reasoning and solving problems (e.g. making plans, making decisions, or learning new things)..	<input type="checkbox"/>	<input type="checkbox"/>
f. I sometimes get confused (e.g. I do not know where I am, who is around, or what day it is).....	<input type="checkbox"/>	<input type="checkbox"/>
g. I forget a lot (e.g. things that happened recently, where I put things, or to keep appointments).....	<input type="checkbox"/>	<input type="checkbox"/>
h. I do not keep my attention on any activity for long.....	<input type="checkbox"/>	<input type="checkbox"/>
i. I make mistakes more than usual.....	<input type="checkbox"/>	<input type="checkbox"/>
j. I have difficulty doing things which involve thought and concentration.....	<input type="checkbox"/>	<input type="checkbox"/>

Thank you for completing Section A

Please continue with Section B


Part 3 - Body Chart

This question is about recent pain you may have had in any part of your body. By pain we also mean ache, discomfort or stiffness. Please do not include pain due to a feverish illness such as flu. If you are a woman please do not include pain related to your monthly period.

1. In the past 4 weeks, have you had pain that has lasted for one day or longer in any part of your body?

(Please put a cross in one box only)

Yes ....	<input type="checkbox"/>	→ Please shade in the diagram below any pain that has lasted for one day or longer in the past 4 weeks
No ....	<input type="checkbox"/>	→ Please continue with Part 4



On this page we want you to think about **HOW INTENSE (how severe)** the pain in different parts of your body has been in the **last 6 months**.

2. In the **past 6 months, on average, how intense were each of these pains** rated on a 0-10 scale where 0 is "no pain" and 10 is "pain as bad as could be"? (That is, your usual pain at times you were experiencing pain.)

*For each pain, please put a cross in one box. For pains that do not apply to you please put a cross in box 0 to indicate No Pain.*

	No Pain 0	1	2	3	4	5	6	7	8	9	Pain as bad as could be 10
a. Headache	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Neck pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Shoulder pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Elbow pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Hand pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Chest pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Abdominal pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Back pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Hip pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Knee pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Foot pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**SECTION B: PAINS AND JOINT PROBLEMS**

This section is about any pains you may have or any problems with your joints.

Please fill in all the parts even if you do not suffer from any pains or joint problems.

Please follow the instructions for each part.



Part 1 - Specific joint pains and problems in the last year

We are interested to know how many people suffer from specific joint pains and problems. Please answer each of the following questions about your hands, hips, knees, and feet, **even if you do not suffer from any problems**.

We would like you to tell us if you have had **any problems** with your **hands**. 'Hand problems' relate to **any hand symptoms** you may have experienced; for example, **pain or stiffness**, or any **knobbly swellings** on your hands, including your fingers and thumbs.

1. Have you had any problems with your **HANDS**, including your fingers and thumbs, over the **last year**?  
*(Please put a cross in one box only)*
- Yes.... ☐ No.... ☐
2. Have you had any pain in your **HANDS**, including in your fingers and thumbs, over the **last year**?  
*(Please put a cross in one box only)*
- Yes.... ☐ No.... ☐

We are also interested in any pain you may have had in your legs.

3. Have you had pain in the last year in and **around the HIP**?  
*(Please put a cross in one box only)*
- Yes.... ☐ No.... ☐
4. Have you had pain in the last year in and **around the KNEE**?  
*(Please put a cross in one box only)*
- Yes.... ☐ No.... ☐
5. Have you had pain in the last year in and **around the FOOT**?  
*(Please put a cross in one box only)*
- Yes.... ☐ No.... ☐

Part 2 - Pain in different parts of the body in the last 6 months

This is about pain in different parts of your body in the last 6 months. Please complete each of the following questions **even if you have not suffered pain in any of these areas** in the last 6 months.

The first question is about **HOW OFTEN** you have had pain in different parts of the body in the **last 6 months**.


1. On how many days in the **last 6 months** have you had.....?  
*(For each pain, please put a cross in one box)*

	No days	1 to 30 days	31 to 89 days	90 or more days
a. Headache	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Neck pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Shoulder pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Elbow pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Hand pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Chest pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Abdominal pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. Back pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. Hip pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Knee pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Foot pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



## **Appendix 4: Baseline Regional Pains Questionnaire**

# **Questionnaire about Hand, Hip Knee & Foot Problems**



**SECTION E - CONTINUING TO HELP WITH THIS STUDY**

Thank you very much for completing this second questionnaire.

Your continued involvement in this study is greatly appreciated.

We may contact you again about the possibility of taking a further part in this study. Would you be willing to be contacted by telephone? Giving us permission to contact you by telephone does not mean that you must take part.

Yes, I am happy to be contacted by telephone.....

☐

My telephone number is.....

No, I do not want to be contacted by telephone.....

☐

**Even if you do not want to be contacted again by telephone, the answers in your questionnaire will still be very useful to us.**

Please check that you have answered all of the questions that apply to you.

Please return this questionnaire in the envelope provided (no stamp needed).

If you have any questions, telephone Ross Wilkie on 01782 583904.

**Thank you for taking the time to fill in this questionnaire**

## **Questionnaire about hand, hip, knee and foot problems**

### **Instructions for this questionnaire**

The aim of this questionnaire is to find out about joint pain and problems of local people. The answers you give in the questionnaire will be treated in the strictest confidence.

**If you have had HAND PROBLEMS AT ANY TIME IN THE LAST YEAR please fill in SECTION A on PAGE 3. For this questionnaire "hand problems" relate to any hand symptoms you may experience; for example, pain or stiffness, or any knobbly swellings on your hands, including your fingers and thumbs.**

**If you have had HIP PAIN AT ANY TIME IN THE LAST YEAR, please fill in SECTION B, on PAGE 25.**

**If you have had KNEE PAIN AT ANY TIME IN THE LAST YEAR, please fill in SECTION C, on PAGE 33.**

**If you have had FOOT PAIN AT ANY TIME IN THE LAST YEAR, please fill in SECTION D, on PAGE 41.**

**If you have more than one joint problem, please complete ALL the sections which apply to you**

**When you have filled in the relevant sections, please turn to the last page of this questionnaire, SECTION E, on PAGE 49.**



Please answer all of the questions unless the instructions ask you to do something else.

Please write in BLOCK CAPITALS where appropriate.

Most of the questions can be answered by putting a cross in a box like this



When you have finished please check that you have answered all of the questions and then return the questionnaire in the envelope enclosed. You do not need a stamp. Please return the questionnaire in the next two weeks.

More details about this project are available in the information sheet enclosed. If you would like further information about this project please contact Ross Wilkie, survey co-ordinator, on 01782 583904.

**Thank you for your help with this important research project**

# Section E

## For everyone to complete

**Please complete this section first.**

1. Are you:

Female.....☐

Male.....☐

2. What is your date of birth?

Day	Month	Year
<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

(For example - if you were born on the 5th June 1936, this would be entered as 05/06/36)

3. Please enter today's date:

Day	Month	Year
<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Because of pain in my feet:

15. My feet are worse in the morning

None of the time

☐

On some days

☐

On most/every day(s)

☐

16. My feet are more painful in the evening

None of the time

☐

On some days

☐

On most/every day(s)

☐

17. I get shooting pains in my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐

18. I am unable to carry out my previous work

None of  
the time

☐

On some days

☐

On most/every day(s)

☐

Not  
applicable

☐

19. I no longer do all my previous activities (sport, dancing, hill-walking, etc.)

None of  
the time

☐

On some days

☐

On most/every day(s)

☐

Not  
applicable

☐

# Section A

# HAND

Please complete this section if you have had hand problems at any time in the last year

THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN COMPLETING  
THIS QUESTIONNAIRE

PLEASE CONTINUE AND FILL IN SECTION E ON PAGE 47

**Because of pain in my feet:**

7. I avoid standing for a long time

None of the time

☐

On some days

☐

On most/every day(s)

☐

8. I catch the bus or use the car more often

None of the time

☐

On some days

☐

On most/every day(s)

☐

9. I need help with housework/shopping

None of the time

☐

On some days

☐

On most/every day(s)

☐

10. I still do everything but with more pain or discomfort

None of the time

☐

On some days

☐

On most/every day(s)

☐

11. I get irritable when my feet hurt

None of the time

☐

On some days

☐

On most/every day(s)

☐

12. I feel self-conscious about my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐

13. I get self-conscious about the shoes I have to wear

None of the time

☐

On some days

☐

On most/every day(s)

☐

14. I have constant pain in my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐

Below are some statements about problems people have because of pain in their feet.

For each statement indicate if this has applied to you during **the past month**. If so, was this only on some days or on most or every day in the past month?

*Please put a cross in one box on each line.*

**Because of pain in my feet:**

1. I avoid walking outside at all			
None of the time	On some days	On most/every day(s)	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
2. I avoid walking long distances			
None of the time	On some days	On most/every day(s)	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
3. I don't walk in a normal way			
None of the time	On some days	On most/every day(s)	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
4. I walk slowly			
None of the time	On some days	On most/every day(s)	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
5. I have to stop and rest my feet			
None of the time	On some days	On most/every day(s)	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
6. I avoid hard or rough surfaces when possible			
None of the time	On some days	On most/every day(s)	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	

**SECTION A - HAND PROBLEMS**

For this questionnaire 'hand problems' relate to **any hand symptoms** you may experience; for example, **pain or stiffness**, or any **knobbly swellings** on your hands, including your fingers and thumbs.

**Part 1**

1. Have you had any problems with your **hands**, including your fingers and thumbs, over the **last year**? (*Please put a cross in one box only*)

Yes..... ☐

No..... ☐

2. Have you had **hand pain**, including your fingers and thumbs, over the **last year**?

Yes..... ☐ Please continue with **Question 3**, below.

No..... ☐ Please go to **Part 2**, on page 6.

3. Please think back over the **last 12 months**. Please put a cross in one box only to show if you have pain, or have had pain:

in both hands (including fingers and thumbs)	the right hand only (including fingers and thumbs)	the left hand only (including fingers and thumbs)
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Again, please think back over the **last 12 months**. On how many days have you had this **hand pain**? (*Please put a cross in one box only*)

Less than 7 days..... ☐

1 to 4 weeks..... ☐

More than 1 month but less than 3 months..... ☐

3 months or more..... ☐



**Part 2**

1. These questions are about the severity of any **hand problems** you may have had during the **past month**.

*(Please put a cross in the box of the most appropriate answer)*

During the <b>past month</b> ...	Severe	Moderate	Mild	Very mild	None
a. How would you describe the hand <b>pain</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. How would you describe the hand <b>stiffness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. How would you describe the hand <b>aching</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. How would you describe the hand <b>tenderness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. How would you describe the hand <b>weakness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. How would you describe the hand <b>clumsiness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. How would you describe the <b>burning</b> sensation in your hand you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. How would you describe the <b>tingling</b> sensation in your hand you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. How would you describe the <b>numbness</b> in your hand you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**SECTION D - FOOT PAIN**

These questions are about any **pain** you may have experienced in your **FEET**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the foot?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both feet... ☐ in the right foot only... ☐ in the left foot only... ☐

3. Again please think back over the **last 12 months**. On how many days have you had this pain?

*(Please put a cross in one box only)*

<input type="checkbox"/>	Less than 7 days
<input type="checkbox"/>	1 - 4 weeks
<input type="checkbox"/>	More than 1 month but less than 3 months
<input type="checkbox"/>	3 months or more

4. Have you ever injured your foot badly enough to see a doctor about it?

<input type="checkbox"/>	No
<input type="checkbox"/>	Right foot only
<input type="checkbox"/>	Left foot only
<input type="checkbox"/>	Both feet

2. These questions are about **how often** you may have had **hand problems** during the **past month**.

*(Please put a cross in the box of the most appropriate answer)*

During the <b>past month</b> ...	All days	Most days	Some days	Few days	No days
a. How often did you have <b>severe pain</b> in your hands?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. How often did you have <b>pain in two or more hand joints</b> at the same time?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. How often did your hand joints feel <b>hot or warm</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. How often did the <b>morning stiffness</b> in your hands last for more than one hour from the time you woke up?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. How often did hand problems cause you to <b>drop objects</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. How often did your hand pain make it difficult for you to <b>sleep</b> ?.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. How often did your hand problems make you feel <b>frustrated</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. How often have you had to take <b>medication</b> for your hand symptoms?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Please continue with Part 3, on page 8.**





**Part 3**

This question is about any **recent pain** you may have had in your **hands**.

In the **past month**, have you had any ache or pain that has lasted for **one day or longer** in your hands? Please **do not** include pain due to feverish illness such as flu.

Yes..... ☐ Please shade in the diagrams below.

No..... ☐ Please go to **Part 4**, on page 9.

**Backs of Hands****Left****Right****Left****Palms of Hands****Right**

# Section D

# FOOT

**Please complete this section if you have had foot pain at any time in the last year**

**Part 4**


We are interested to know whether you have any finger nodes. A finger node is a firm, knobby swelling on the back of a finger joint.

**For example:**

A finger without nodes:



A finger with nodes:



Please look at your hands and then answer the following questions:

1. Do you think you have any finger **nodes/swellings**?


Yes..... ☐

Please continue with **question 2**, on page 10.

No..... ☐

Please go to **Part 5**, on page 10.

2. On the diagrams below please circle any finger **joints** on which you have nodes/swellings (you may have several nodes/swellings).



### Part 5

1. The following questions are about the **appearance** of your hands during the **past month**.

*(Please put a cross in the box that best describes how you have felt)*

	Strongly agree	Agree	Neither agree nor disagree	Disagree	Strongly disagree
a. I was <b>satisfied</b> with the appearance of my hands.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. The appearance of my hands sometimes made me <b>uncomfortable</b> in public.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. The appearance of my hands made me <b>depressed</b> ..	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. The appearance of my hands <b>interfered</b> with my usual social activities.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

19. Lying in bed

None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
----------------------------------	----------------------------------	--------------------------------------	------------------------------------	-------------------------------------

20. Getting in/out of bath

None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
----------------------------------	----------------------------------	--------------------------------------	------------------------------------	-------------------------------------

21. Sitting

None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
----------------------------------	----------------------------------	--------------------------------------	------------------------------------	-------------------------------------

22. Getting on/off toilet

None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
----------------------------------	----------------------------------	--------------------------------------	------------------------------------	-------------------------------------

23. Heavy domestic duties

None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
----------------------------------	----------------------------------	--------------------------------------	------------------------------------	-------------------------------------

24. Light domestic duties

None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
----------------------------------	----------------------------------	--------------------------------------	------------------------------------	-------------------------------------

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN COMPLETING  
THIS QUESTIONNAIRE**

**PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU (INCLUDING SECTION E ON PAGE 47)**

11. Standing

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. Bending to the floor

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13. Walking on flat

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. Getting in/out of car

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

15. Going shopping

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

16. Putting on socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

17. Rising from bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

18. Taking off socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 6

1. These questions refer to **HAND AND FINGER FUNCTION**.

*(Please put a cross in the box under the most appropriate answer for each question)*

During the <b>past month...</b>	All days	Most days	Some days	Few days	No days
a. Could you <b>easily</b> write with a pen or pencil?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Could you <b>easily</b> button a shirt or blouse?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Could you <b>easily</b> turn a key in a lock?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Could you <b>easily</b> tie a knot or a bow?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Could you <b>easily</b> open a new jar of food?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. These questions refer to **ARM FUNCTION**.

*(Please put a cross in the box of the most appropriate answer for each question)*

During the <b>past month...</b>	All days	Most days	Some days	Few days	No days
a. Could you <b>easily</b> wipe your mouth with a napkin?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Could you <b>easily</b> put on a pullover sweater?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Could you <b>easily</b> comb or brush your hair?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Could you <b>easily</b> scratch your low back with your hand?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Could you <b>easily</b> reach shelves that were above your head?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 7**

1. We are interested in finding out how your **hand problems** have affected your ability to carry out your **usual activities** over the **past week**.

*(Please put a cross in the box which best describes your usual abilities)*

During the <b>past week</b> ...	No difficulty	Mild difficulty	Moderate difficulty	Severe difficulty	Could not do at all due to problems
a. Could you <b>pick up</b> coins from a table top without sliding them?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Could you <b>hold</b> a book while reading?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Could you <b>clench</b> your fist?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Could you <b>self-care</b> after going to the toilet (e.g. pull-up your underclothes, wipe yourself)?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Could you <b>open</b> packaging (e.g. packets of food, parcels)?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 2**

The following questions concern the amount of **joint stiffness** (not pain) you have experienced in the **last 48 hours** in your knee. Stiffness is a sensation of restriction or slowness in the ease with which you move your joints.

*(Please put a cross in one box on each line)*

6. How severe is your stiffness after waking in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. How severe is your stiffness after sitting, lying or resting in the day?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 3**

The following questions concern your **physical function**. By this we mean your ability to move around and look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the **last 48 hours** due to your knee.

*(Please put a cross in one box on each line)*

**Question: What degree of difficulty do you have?**

8. Descending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. Ascending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Rising from sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

For all of the following questions please think back over the last 48 hours.

Part 1

The following questions concern the amount of **pain** you have experienced in your knee. For each question please enter the amount of pain experienced in the last 48 hours.

(Please put a cross in one box on each line)

Question: How much pain do you have?

1. Walking on a flat surface

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Going up or down stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. At night while in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Sitting or lying

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Standing upright

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 8

1. The following questions concern the amount of pain you have experienced in your hands. For each situation please enter the amount of pain experienced in the last week.

(Please put a cross in the box of the most appropriate answer)

QUESTION: How much pain do you have in your hands?

a. At rest (i.e. when not using your hands).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

b. When gripping objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

c. When lifting objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

d. When turning objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

e. When squeezing objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



2. The following question concerns the amount of joint stiffness (not pain) you have experienced in the last week in your hands. Stiffness is a sensation of restriction or slowness in the ease with which you move your hands.

*(Please put a cross in the box of the most appropriate answer)*

- a. How **severe** is stiffness in your hands **after first wakening** in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. The following questions concern your physical function. By this we mean your ability to move around and to look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the last week due to your hand problem.

*(Please put a cross in the box of the most appropriate answer)*

**QUESTION: How much difficulty do you have with the following?**

- a. Turning taps on.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- b. Turning a round door-knob or handle.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- c. Doing up buttons.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

### SECTION C - KNEE PAIN

These questions are about any **pain** you may have experienced in your **KNEES**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the knee?

Yes.....☐

No.....☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both knees..☐ in the right knee only..☐ in the left knee only..☐

3. Again please think back over the **last 12 months**. On how many days have you had this pain?

*(Please put a cross in one box only)*

<input type="checkbox"/>	Less than 7 days
<input type="checkbox"/>	1 - 4 weeks
<input type="checkbox"/>	More than 1 month but less than 3 months
<input type="checkbox"/>	3 months or more

4. Have you ever injured your knee badly enough to see a doctor about it?

<input type="checkbox"/>	No
<input type="checkbox"/>	Right knee only
<input type="checkbox"/>	Left knee only
<input type="checkbox"/>	Both knees

QUESTION: How much difficulty do you have with the following?

d. Fastening jewellery (e.g. watches, earrings, cufflinks, necklaces, brooches and bracelets).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

e. Opening a new jar.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

f. Carrying a full pot with one hand (e.g. carrying any reasonably heavy object such as a saucepan).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

g. Peeling vegetables/fruits.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

h. Picking up large heavy objects.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

i. Wringing out washcloths (e.g. squeezing a wet sponge or flannel).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 9**

1. Listed below are a number of **symptoms** that you may or may not have experienced since your hand problem.

The **first column** is about whether you have experienced any of these symptoms since you have had your hand problem.

The **second column** is about whether you believe that these symptoms are related to your hand problem.

**Please complete both columns by putting a cross in one box for each column, to indicate "Yes" or "No".**

	<u>First column</u>		<u>Second column</u>
	I have experienced this symptom since my hand problem		This symptom is related to my hand problem
a. Pain/ache.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
b. Sore Throat.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
c. Nausea.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
d. Breathlessness.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
e. Weight Loss.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
f. Fatigue.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
g. Stiff Joints.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
h. Sore Eyes.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
i. Wheeziness.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
j. Headaches.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
k. Upset Stomach.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
l. Sleep Difficulties.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
m. Dizziness.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	
n. Loss of Strength.....	Yes <input type="checkbox"/> No <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/>	

# Section C

## KNEE

**Please complete this section if you have had knee pain at any time in the last year**

2. We are interested in your own personal views of how you now see your current hand problem.

Please indicate how much you agree or disagree with the following statements about your hand problem by putting a cross in one box on each line.

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
a. My hand problem will last a short time.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. My hand problem is likely to be permanent rather than temporary...	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. My hand problem will last for a long time.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. This hand problem will pass quickly.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. I expect to have this hand problem for the rest of my life.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. My hand problem is a serious condition.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. My hand problem has major consequences on my life.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. My hand problem does not have much effect on my life.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. My hand problem strongly affects the way others see me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. My hand problem has serious financial consequences.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. My hand problem causes difficulties for those who are close to me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
l. There is a lot which I can do to control my hand symptoms.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
m. What I do can determine whether my hand problem gets better or worse.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
n. The course of my hand problem depends on me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
o. Nothing I do will affect my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
p. I have the power to influence my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
q. My actions will have no effect on the outcome of my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
r. My hand problem will improve in time.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
s. There is very little that can be done to improve my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
t. My treatment will be effective in curing my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
u. The negative effects of my hand problem can be prevented (avoided) by my treatment.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
v. My treatment can control my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
w. There is nothing which can help my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
x. The symptoms of my hand problem are puzzling to me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
y. My hand problem is a mystery to me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
z. I don't understand my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

19. Lying in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

20. Getting in/out of bath

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

21. Sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

22. Getting on/off toilet

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

23. Heavy domestic duties

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

24. Light domestic duties

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN COMPLETING  
THIS QUESTIONNAIRE**

**PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU (INCLUDING SECTION E ON PAGE 47)**

11. Standing

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. Bending to the floor

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13. Walking on flat

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. Getting in/out of car

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

15. Going shopping

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

16. Putting on socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

17. Rising from bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

18. Taking off socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
aa. My hand problem doesn't make any sense to me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
bb. I have a clear picture or understanding of my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
cc. The symptoms of my hand problem change a great deal from day to day.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
dd. My symptoms come and go in cycles.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
ee. My hand problem is very unpredictable.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
ff. I go through cycles in which my hand problem gets better and worse.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
gg. I get depressed when I think about my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
hh. When I think about my hand problem I get upset.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
ii. My hand problem makes me feel angry.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
jj. My hand problem does not worry me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
kk. Having this hand problem makes me feel anxious.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
ll. My hand problem makes me feel afraid.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. We are interested in what you consider may have been the cause of your hand problem. As people are very different, there is no correct answer for this question. We are most interested in your own views about the factors that caused your hand problem rather than what others including doctors or family may have suggested to you. Below is a list of possible causes for your hand problem. Please indicate how much you agree or disagree that they were causes for your hand problem by putting a cross in one box for each line.

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
<b>Possible causes...</b>					
a. Stress or worry.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Hereditary - it runs in my family.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. A Germ or virus.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Diet or eating habits.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Chance or bad luck.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Poor medical care in my past.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. Pollution in the environment.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. My own behaviour.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. My mental attitude e.g. thinking about life negatively.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
j. Family problems or worries caused my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
k. Overwork.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 2

The following questions concern the amount of **joint stiffness** (not pain) you have experienced in the **last 48 hours** in your hip. Stiffness is a sensation of restriction or slowness in the ease with which you move your joints.

*(Please put a cross in one box on each line)*

6. How severe is your stiffness after waking in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. How severe is your stiffness after sitting, lying or resting in the day?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 3

The following questions concern your **physical function**. By this we mean your ability to move around and look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the **last 48 hours** due to your hip.

*(Please put a cross in one box on each line)*

**Question: What degree of difficulty do you have?**

8. Descending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. Ascending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Rising from sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

For all of the following questions please think back over the last 48 hours.

Part 1

The following questions concern the amount of **pain** you have experienced in your hip. For each question please enter the amount of pain experienced in the **last 48 hours**.

(Please put a cross in one box on each line)

Question: How much pain do you have?

1. Walking on a flat surface

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Going up or down stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. At night while in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Sitting or lying

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Standing upright

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
l. My emotional state e.g. feeling down, lonely, anxious, empty.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
m. Ageing.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
n. Alcohol.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
o. Smoking.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
p. Accident or injury.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
q. My personality.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
r. Altered immunity (change in the body's ability to fight illness).	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Please list the three most important factors that you now believe caused YOUR hand problem, starting with the most important cause first and so on. You may use any of the items from the list above, or you may have additional ideas of your own.

The most important causes for me :-

1.	.....
2.	.....
3.	.....



**Part 10**

1. These are some general questions about your hands.

*(Please put a cross in one box for each question)*

a. Do you consider yourself to be...

Right-handed.. ☐

Left-handed.. ☐

Both.. ☐

b. Which hand gives you the most problem?

Right hand... ☐

Left hand ... ☐

Both... ☐

c. In the past, have you **ever** injured your hands badly enough to see a doctor?

No.. ☐

Right hand only.. ☐

Left hand only.. ☐

Both hands... ☐

d. In the past, have you **ever** had any hand operations?

No.. ☐

Right hand only.. ☐

Left hand only.. ☐

Both hands... ☐

e. In your opinion, has your past or present **job** involved excessive use of your hands?

Yes..... ☐

No..... ☐

f. In your opinion, have your past or present **hobbies or pastimes** involved excessive use of your hands?

Yes..... ☐

No..... ☐

**SECTION B - HIP PAIN**

These questions are about any **pain** you may have experienced in your **HIPS**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the hip?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both hips.. ☐

in the right hip only.. ☐

in the left hip only.. ☐

3. Again please think back over the **last 12 months**. On how many days have you had this pain?

*(Please put a cross in one box only)*

☐

Less than 7 days

☐

1 - 4 weeks

☐

More than 1 month but less than 3 months

☐

3 months or more

4. Have you ever injured your hip badly enough to see a doctor about it?

☐

No

☐

Right hip only

☐

Left hip only

☐

Both hips

g. How long have you had your hand problem?

(Please let us know the **number of years**, or if less than one year the **number of months**, you have had your hand problem)

Years....

Months (if less than 1 year).....

h. We are interested in finding out what you believe is the **main** cause of your hand problem.

(Please put a cross in **ONE** box only)

Osteoarthritis.....

Rheumatoid arthritis.....

Carpal tunnel syndrome.....

Raynaud's disease.....

Injury (e.g. broken bones, burns, sprains).....

Other.....

Don't know.....

i. This question refers to the overall impact of your hand problem.

Considering all the ways that your hand problem affects you, how well are you doing compared to other people your age?

Very wellWellFairPoorVery Poorly

- j. Have you consulted your G.P. (family doctor) in the **past 12 months** because of your hand problem?

*(Please put a cross in one box only)*

Yes..... ☐

No..... ☐

- k. Which of the following services or treatments have you used in the **past 12 months** because of your hand problem? For **each service** you have used please put a cross to show whether the NHS provided this, or if you had private treatment. If you have used NHS **and** private services, please cross both boxes. For any service that you have not used, please leave blank.

	Yes (NHS)	Yes (Private)
a. Physiotherapy.....	<input type="checkbox"/>	<input type="checkbox"/>
b. Occupational therapy.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Hospital specialist.....	<input type="checkbox"/>	<input type="checkbox"/>
d. Acupuncture.....	<input type="checkbox"/>	<input type="checkbox"/>
e. Osteopath/Chiropractor....	<input type="checkbox"/>	<input type="checkbox"/>
f. Drugs on prescription.....	<input type="checkbox"/>	<input type="checkbox"/>
g. Hand operation.....	<input type="checkbox"/>	<input type="checkbox"/>
h. Hand injection.....	<input type="checkbox"/>	<input type="checkbox"/>

i. Other (please state) .....

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN COMPLETING  
THIS QUESTIONNAIRE**

**PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU (INCLUDING SECTION E ON PAGE 47)**

# Section B


## HIP

**Please complete this section if you have had  
hip pain at any time in the last year**

## **Appendix 5: Regional Pains Questionnaire 3 years Follow-up**

# **Questionnaire about Hand, Hip Knee & Foot Problems**

**3 Year Follow-up**



LREC Number: 05/Q2604/20

Version 2, Date: 07/02/2005

## **Questionnaire about hand, hip, knee and foot problems**

### **Instructions for this questionnaire**

The aim of this questionnaire is to find out about joint pain and problems of local people. The answers you give in the questionnaire will be treated in the strictest confidence.

**If you have had HAND PROBLEMS AT ANY TIME IN THE LAST YEAR please fill in SECTION A on PAGE 4. For this questionnaire "hand problems" relate to any hand symptoms you may experience; for example, pain or stiffness, or any knobbly swellings on your hands, including your fingers and thumbs.**

**If you have had HIP PAIN AT ANY TIME IN THE LAST YEAR, please fill in SECTION B, on PAGE 20.**

**If you have had KNEE PAIN AT ANY TIME IN THE LAST YEAR, please fill in SECTION C, on PAGE 28.**

**If you have had FOOT PAIN AT ANY TIME IN THE LAST YEAR, please fill in SECTION D, on PAGE 36.**

**If you have more than one joint problem, please complete EACH section which applies to you**

**When you have filled in the sections relevant to you, please turn to the last page of this questionnaire, on PAGE 42.**

Thank you very much for completing this second questionnaire.

Your continued involvement in this study is greatly appreciated.

**Please check that you have answered all the questions that apply to you.**

Please return the questionnaire in the envelope provided (no stamp needed).

If you have any questions, please telephone Rosie Lacey on 01782 583921.

**Thank you again for taking the time to fill in this questionnaire.**

**Because of pain in my feet:**

15. My feet are worse in the morning

None of the time

☐

On some days

☐

On most/every day(s)

☐

16. My feet are more painful in the evening

None of the time

☐

On some days

☐

On most/every day(s)

☐

17. I get shooting pains in my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐

18. I am unable to carry out my previous work

None of  
the time

☐

On some days

☐

On most/every day(s)

☐

Not  
applicable

☐

19. I no longer do all my previous activities (sport, dancing, hill-walking, etc.)

None of  
the time

☐

On some days

☐

On most/every day(s)

☐

Not  
applicable

☐

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS QUESTIONNAIRE**

Please answer all of the questions unless the instructions ask you to do something else.

Please write in **BLOCK CAPITALS** where appropriate.

Most of the questions can be answered by putting a cross in a box like this:



When you have finished please check that you have answered all of the questions and then return the questionnaire in the envelope enclosed. You do not need a stamp. Please return the questionnaire in the next two weeks.

More details about this project are available in the information sheet enclosed. If you would like further information about this project please contact Rosie Lacey, survey co-ordinator, on 01782 583921.

**Thank you for your help with this research project**

**Please complete this section first.**

1. Are you:

Female.....

☐

Male.....

☐

2. What is your date of birth?

Day	Month	Year
<input type="text"/>	<input type="text"/>	<input type="text"/>

(For example - if you were born on the 5th June 1936, this would be entered as 05/06/36)

3. Please enter today's date:

Day	Month	Year
<input type="text"/>	<input type="text"/>	<input type="text"/>

**Because of pain in my feet:**

7. I avoid standing for a long time

None of the time

☐

On some days

☐

On most/every day(s)

☐

8. I catch the bus or use the car more often

None of the time

☐

On some days

☐

On most/every day(s)

☐

9. I need help with housework/shopping

None of the time

☐

On some days

☐

On most/every day(s)

☐

10. I still do everything but with more pain or discomfort

None of the time

☐

On some days

☐

On most/every day(s)

☐

11. I get irritable when my feet hurt

None of the time

☐

On some days

☐

On most/every day(s)

☐

12. I feel self-conscious about my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐

13. I get self-conscious about the shoes I have to wear

None of the time

☐

On some days

☐

On most/every day(s)

☐

14. I have constant pain in my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐



Below are some statements about problems people have because of pain in their feet.

For each statement indicate if this has applied to you during **the past month**. If so, was this only on some days or on most or every day in the past month?

**Please put a cross in one box on each line.**

**Because of pain in my feet:**

1. I avoid walking outside at all

None of the time

☐

On some days

☐

On most/every day(s)

☐

2. I avoid walking long distances

None of the time

☐

On some days

☐

On most/every day(s)

☐

3. I don't walk in a normal way

None of the time

☐

On some days

☐

On most/every day(s)

☐

4. I walk slowly

None of the time

☐

On some days

☐

On most/every day(s)

☐

5. I have to stop and rest my feet

None of the time

☐

On some days

☐

On most/every day(s)

☐

6. I avoid hard or rough surfaces when possible

None of the time

☐

On some days

☐

On most/every day(s)

☐

# Section A

## HAND

**Please complete this section if you have had hand problems at any time in the last year**



27188

## **SECTION D - FOOT PAIN**

These questions are about any **pain** you may have experienced in your **FEET**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the foot?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both feet... ☐ in the right foot only... ☐ in the left foot only... ☐

3. Again please think back over the **last 12 months**. On how many days have you had this pain?

***(Please put a cross in one box only)***

- ☐ Less than 7 days
- ☐ 1 - 4 weeks
- ☐ More than 1 month but less than 3 months
- ☐ 3 months or more

4. Have you ever injured your foot badly enough to see a doctor about it?

- ☐ No
- ☐ Right foot only
- ☐ Left foot only
- ☐ Both feet

## SECTION A - HAND PROBLEMS

For this questionnaire 'hand problems' relate to **any hand symptoms** you may experience; for example, **pain or stiffness**, or any **knobbly swellings** on your hands, including your fingers and thumbs.

### Part 1

1. Have you had any **problems** with your **hands**, including your fingers and thumbs, over the **last year**? (*Please put a cross in one box only*)

Yes..... ☐

No..... ☐

2. Have you had **hand pain**, including your fingers and thumbs, over the **last year**?

Yes..... ☐ Please continue with **Question 3**, below.

No..... ☐ Please go to **Part 2**, on page 7.

3. Please think back over the **last 12 months**. Please put a cross in one box only to show if you have pain, or have had pain:

in both hands  
(including fingers  
and thumbs)

☐

the right hand only  
(including fingers  
and thumbs)

☐

the left hand only  
(including fingers  
and thumbs)

☐

4. Again, please think back over the **last 12 months**. On how many days have you had this **hand pain**? (*Please put a cross in one box only*)

Less than 7 days..... ☐

1 to 4 weeks..... ☐

More than 1 month but less than 3 months..... ☐

3 months or more..... ☐

## Part 2

1. These questions are about the **severity** of any **hand problems** you may have had during the **past month**.

*(Please put a cross in the box of the most appropriate answer)*

During the <b>past month</b> ...	Severe	Moderate	Mild	Very mild	None
a. How would you describe the hand <b>pain</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. How would you describe the hand <b>stiffness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. How would you describe the hand <b>aching</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. How would you describe the hand <b>tenderness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. How would you describe the hand <b>weakness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. How would you describe the hand <b>clumsiness</b> you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. How would you describe the <b>burning</b> sensation in your hand you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. How would you describe the <b>tingling</b> sensation in your hand you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i. How would you describe the <b>numbness</b> in your hand you usually had?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

# Section D FOOT

**Please complete this section if you have had foot pain at any time in the last year**

2. These questions are about **how often** you may have had **hand problems** during the **past month**.

*(Please put a cross in the box of the most appropriate answer)*

During the <b>past month</b> ...	All days	Most days	Some days	Few days	No days
a. How often did you have <b>severe pain</b> in your hands?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. How often did you have <b>pain in two or more hand joints</b> at the same time?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. How often did your hand joints feel <b>hot or warm</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. How often did the <b>morning stiffness</b> in your hands last for more than one hour from the time you woke up?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. How often did hand problems cause you to <b>drop objects</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. How often did your hand pain make it difficult for you to <b>sleep</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
g. How often did your hand problems make you feel <b>frustrated</b> ?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
h. How often have you had to take <b>medication</b> for your hand symptoms?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Please continue with Part 3, on page 9



Part 3


This question is about any **recent pain** you may have had in your **hands**.

In the **past month**, have you had any ache or pain that has lasted for **one day or longer** in your hands? Please **do not** include pain due to feverish illness such as flu. **(Please put a cross in one box only)**


Yes..... ☐ Please shade in the diagrams below.

No..... ☐ Please go to **Part 4**, on page 10.

Backs of Hands



Palms of Hands



Question: What degree of difficulty do you have?

19. Lying in bed	None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
20. Getting in/out of bath	None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
21. Sitting	None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
22. Getting on/off toilet	None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
23. Heavy domestic duties	None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>
24. Light domestic duties	None <input type="checkbox"/>	Mild <input type="checkbox"/>	Moderate <input type="checkbox"/>	Severe <input type="checkbox"/>	Extreme <input type="checkbox"/>

© Dr Nicholas Bellamy. All rights reserved 1996

THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION

PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU



**Question: What degree of difficulty do you have?**

11. Standing

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. Bending to the floor

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13. Walking on flat

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. Getting in/out of car

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

15. Going shopping

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

16. Putting on socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

17. Rising from bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

18. Taking off socks/stockings


None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 4**

We are interested to know whether you have any finger nodes. A finger node is a firm, knobby swelling on the back of a finger joint.

**For example:**

A finger without nodes:



A finger with nodes:



Please look at your hands and then answer the following questions:


1. Do you think you have any finger **nodes/swellings**?

Yes..... ☐

Please continue with **question 2**, on page 11.


No..... ☐

Please go to **Part 5**, on page 11.



27188

2. On the diagrams below please circle any finger **joints** on which you have nodes/swellings (you may have several nodes/swellings).



**Part 5**

1. These questions refer to **HAND AND FINGER FUNCTION**.  
*(Please put a cross in the box under the most appropriate answer for each question)*

During the <b>past month...</b>	All days	Most days	Some days	Few days	No days
a. Could you <b>easily</b> write with a pen or pencil?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Could you <b>easily</b> button a shirt or blouse?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Could you <b>easily</b> turn a key in a lock?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Could you <b>easily</b> tie a knot or a bow?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Could you <b>easily</b> open a new jar of food?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 2**

The following questions concern the amount of **joint stiffness** (not pain) you have experienced in the **last 48 hours** in your knee. Stiffness is a sensation of restriction or slowness in the ease with which you move your joints.  
*(Please put a cross in one box on each line)*

6. How <b>severe</b> is your stiffness after <b>first waking</b> in the morning?	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. How <b>severe</b> is your stiffness after sitting, lying or resting <b>later in the day</b> ?	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 3**

The following questions concern your **physical function**. By this we mean your ability to move around and look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the **last 48 hours** due to your knee.  
*(Please put a cross in one box on each line)*

**Question: What degree of difficulty do you have?**

8. Descending stairs	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Ascending stairs	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. Rising from sitting	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



For all of the following questions please think back over the last 48 hours.

### Part 1

The following questions concern the amount of **pain** you have experienced in your knee. For each question please enter the amount of pain experienced in the **last 48 hours**.

*(Please put a cross in one box on each line)*

**Question: How much pain do you have?**

1. Walking on a flat surface

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Going up or down stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. At night while in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Sitting or lying

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Standing upright

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

### Part 6

1. The following questions concern the amount of pain you have experienced in your hands. For each situation please enter the amount of **pain** experienced in the **last week**.

*(Please put a cross in the box of the most appropriate answer)*

**QUESTION: How much pain do you have in your hands?**

a. At rest (i.e. when not using your hands).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

b. When gripping objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

c. When lifting objects with your hands.


None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

d. When turning objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

e. When squeezing objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



27188

2. The following question concerns the amount of **joint stiffness** (not pain) you have experienced **in the last week** in your **hands**.  
Stiffness is a sensation of restriction or slowness in the ease with which you move your hands.

*(Please put a cross in the box of the most appropriate answer)*

- a. How **severe** is stiffness in your hands **after first wakening** in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. The following questions concern your **physical function**. By this we mean your ability to move around and to look after yourself.  
For each of the following activities, please indicate the **degree of difficulty** you have experienced **in the last week** due to your **hand problem**.

*(Please put a cross in the box of the most appropriate answer)*

**QUESTION: How much difficulty do you have with the following?**

- a. Turning taps on.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- b. Turning a round door-knob or handle.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- c. Doing up buttons.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## **SECTION C - KNEE PAIN**

These questions are about any **pain** you may have experienced in your **KNEES**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the knee?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both knees.. ☐ in the right knee only.. ☐ in the left knee only.. ☐

3. Again please think back over the **last 12 months**. On how many days have you had this **knee pain**?

*(Please put a cross in one box only)*

<input type="checkbox"/>	Less than 7 days
<input type="checkbox"/>	1 - 4 weeks
<input type="checkbox"/>	More than 1 month but less than 3 months
<input type="checkbox"/>	3 months or more

4. Have you ever injured your knee badly enough to see a doctor about it?

<input type="checkbox"/>	No
<input type="checkbox"/>	Right knee only
<input type="checkbox"/>	Left knee only
<input type="checkbox"/>	Both knees



# Section C

## KNEE

Please complete this section if you have had knee pain at any time in the last year.



**QUESTION: How much difficulty do you have with the following?**

d. Fastening jewellery (e.g. watches, earrings, cufflinks, necklaces, brooches and bracelets).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

e. Opening a new jar.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

f. Carrying a full pot with one hand (e.g. carrying any reasonably heavy object such as a saucepan).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

g. Peeling vegetables/fruits.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

h. Picking up large heavy objects.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

i. Wringing out washcloths (e.g. squeezing a wet sponge or flannel).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 7

We are interested in your own personal views of how you now see your current **hand problem**.

Please indicate how much you agree or disagree with the following statements about your hand problem by **putting a cross in one box on each line**.

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
1. My symptoms come and go in cycles.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. I don't understand my hand problem.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. My hand problem will last a long time.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. What I do can determine whether my hand problem gets better or worse.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. My hand problem is a serious condition.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. My hand problem has major consequences on my life.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. My hand problem does not have much effect on my life.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. My hand problem strongly affects the way others see me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

# Section C

## KNEE

**Please complete this section if you have had knee pain at any time in the last year**

	Strongly disagree	Disagree	Neither agree nor disagree	Agree	Strongly agree
9. My hand problem has serious financial consequences.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. My hand problem causes difficulties for those who are close to me.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11. There is a lot which I can do to control my hand symptoms.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

### Part 8

These are some general questions about your hands.

*(Please put a cross in one box for each question)*

1. Do you consider yourself to be...

Right-handed...	<input type="checkbox"/>	Left-handed...	<input type="checkbox"/>	Both...	<input type="checkbox"/>
-----------------	--------------------------	----------------	--------------------------	---------	--------------------------

2. Which hand gives you the most problem?

Right hand...	<input type="checkbox"/>	Left hand ...	<input type="checkbox"/>	Both...	<input type="checkbox"/>
---------------	--------------------------	---------------	--------------------------	---------	--------------------------

3. Over the past **three years**, have you injured your hands badly enough to see a doctor?

No..	<input type="checkbox"/>	Right hand only..	<input type="checkbox"/>	Left hand only..	<input type="checkbox"/>	Both hands..	<input type="checkbox"/>
------	--------------------------	-------------------	--------------------------	------------------	--------------------------	--------------	--------------------------

4. Over the past **three years**, have you had any hand operations?

No..	<input type="checkbox"/>	Right hand only..	<input type="checkbox"/>	Left hand only..	<input type="checkbox"/>	Both hands..	<input type="checkbox"/>
------	--------------------------	-------------------	--------------------------	------------------	--------------------------	--------------	--------------------------

5. How long have you had your **hand problem**?

(Please let us know the **number of years**, or if less than one year, the **number of months**, you have had your hand problem)

Years....

Months (if less than 1 year).....

6. Thinking overall about your hand problem, during the **past month** how would you describe your **hand problem(s)** in general?

None          Very mild          Mild          Moderate          Severe

☐          ☐          ☐          ☐          ☐

7. In the last **two weeks**, how bothersome has your hand problem been?

Not at all          Slightly          Moderately          Very much          Extremely

☐          ☐          ☐          ☐          ☐

8. If you had to spend the rest of your life with the hand problems you have right now, how would you feel about it?

Very dissatisfied          Somewhat dissatisfied          Neither satisfied nor dissatisfied          Somewhat satisfied          Very satisfied

☐          ☐          ☐          ☐          ☐

9. This question refers to the overall impact of your hand problem.

Considering all the ways that your hand problem affects you, how well are you doing compared to other people your age?

Very well          Well          Fair          Poor          Very poorly

☐          ☐          ☐          ☐          ☐

**Question: What degree of difficulty do you have?**

19. Lying in bed

None          Mild          Moderate          Severe          Extreme  
☐          ☐          ☐          ☐          ☐

20. Getting in/out of bath

None          Mild          Moderate          Severe          Extreme  
☐          ☐          ☐          ☐          ☐

21. Sitting

None          Mild          Moderate          Severe          Extreme  
☐          ☐          ☐          ☐          ☐

22. Getting on/off toilet

None          Mild          Moderate          Severe          Extreme  
☐          ☐          ☐          ☐          ☐

23. Heavy domestic duties

None          Mild          Moderate          Severe          Extreme  
☐          ☐          ☐          ☐          ☐

24. Light domestic duties

None          Mild          Moderate          Severe          Extreme  
☐          ☐          ☐          ☐          ☐

© Dr Nicholas Bellamy. All rights reserved 1996

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION**

**PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU**

Question: What degree of difficulty do you have?

11. Standing

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. Bending to the floor

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13. Walking on flat

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. Getting in/out of car

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

15. Going shopping

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

16. Putting on socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

17. Rising from bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

18. Taking off socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Have you consulted your G.P. (family doctor) in the **past 12 months** because of your hand problem?

(Please put a cross in one box only)

Yes..... ☐

No..... ☐

11. Which of the following services or treatments have you used in the **past 12 months** because of your hand problem? For **each service** you have used please put a cross to show whether the NHS provided this, or if you had private treatment. If you have used NHS **and** private services, please cross both boxes. For any service that you have not used, please leave blank.

	Yes (NHS)	Yes (Private)
a. Physiotherapy.....	<input type="checkbox"/>	<input type="checkbox"/>
b. Occupational therapy.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Hospital specialist.....	<input type="checkbox"/>	<input type="checkbox"/>
d. Acupuncture.....	<input type="checkbox"/>	<input type="checkbox"/>
e. Osteopath/Chiropractor....	<input type="checkbox"/>	<input type="checkbox"/>
f. Drugs on prescription.....	<input type="checkbox"/>	<input type="checkbox"/>
g. Hand operation.....	<input type="checkbox"/>	<input type="checkbox"/>
h. Hand injection.....	<input type="checkbox"/>	<input type="checkbox"/>
i. Other (please state) .....		

THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION

PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU

## Part 2

The following questions concern the amount of **joint stiffness** (not pain) you have experienced in the **last 48 hours** in your hip. Stiffness is a sensation of restriction or slowness in the ease with which you move your joints.

*(Please put a cross in one box on each line)*

6. How **severe** is your stiffness after **first waking** in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. How **severe** is your stiffness after sitting, lying or resting **later in the day**?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## Part 3

The following questions concern your **physical function**. By this we mean your ability to move around and look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the **last 48 hours** due to your hip.

*(Please put a cross in one box on each line)*

**Question: What degree of difficulty do you have?**

8. Descending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. Ascending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Rising from sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



For all of the following questions please think back over the last 48 hours.

### Part 1

The following questions concern the amount of **pain** you have experienced in your hip. For each question please enter the amount of pain experienced in the **last 48 hours**.

*(Please put a cross in one box on each line)*

**Question: How much pain do you have?**

1. Walking on a flat surface

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Going up or down stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. At night while in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Sitting or lying

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Standing upright

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

# Section B

## HIP

**Please complete this section if you have had hip pain at any time in the last year**



27188

## **SECTION B - HIP PAIN**

These questions are about any **pain** you may have experienced in your **HIPS**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the hip?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both hips.. ☐ in the right hip only.. ☐ in the left hip only.. ☐

3. Again please think back over the **last 12 months**. On how many days have you had this **hip pain**?

***(Please put a cross in one box only)***

- ☐ Less than 7 days
- ☐ 1 - 4 weeks
- ☐ More than 1 month but less than 3 months
- ☐ 3 months or more

4. Have you ever injured your hip badly enough to see a doctor about it?

- ☐ No
- ☐ Right hip only
- ☐ Left hip only
- ☐ Both hips


## **Appendix 6: Regional Pains Questionnaire 6 years Follow-up**

# Questionnaire about Joint Pain

6-Year Follow-up

REC Number: 06/Q2604/20

Version 3, Date: 04/02/2008



## Questionnaire about Joint Pain

### Instructions for this questionnaire

The aim of this questionnaire is to find out more about joint pain and the problems people have with their joints.

**PLEASE READ ALL THE INSTRUCTIONS BELOW BEFORE YOU START TO COMPLETE THE QUESTIONNAIRE**

- If you have had **HAND PROBLEMS** (pain or stiffness, or knobbly swellings on your hands, including fingers and thumbs) **AT ANY TIME IN THE LAST YEAR** → please fill in **SECTION A** on **PAGE 5**
- If you have had **HIP PAIN AT ANY TIME IN THE LAST YEAR** → please fill in **SECTION B** on **page 15**
- If you have had **KNEE PAIN AT ANY TIME IN THE LAST YEAR** → please fill in **SECTION C** on **PAGE 23**
- If you have had **FOOT PAIN AT ANY TIME IN THE LAST YEAR** → please fill in **SECTION D** on **PAGE 31**
- Whatever your joint pain, if you are in full-time or part-time paid work **OR** you have been off work due to ill health for more than 1 week but less than six months → please fill in **SECTION E** on **page 37**

**If you have more than one joint problem, please complete EACH section which applies to you**

When you have filled in the sections relevant to you, please complete **SECTION F ON PAGE 43**, and then turn to the last page of this questionnaire, **PAGE 47**.

Thank you very much for completing this second questionnaire.

Your continued involvement in this study is greatly appreciated.

**Please check that you have answered all the questions that apply to you.**

Please return the questionnaire in the envelope provided (no stamp needed).

If you have any questions, please telephone Rosie Lacey on 01782 584721.

**Thank you again for taking the time to fill in this questionnaire.**

Please answer all of the questions unless the instructions ask you to do something else.

Please write in **BLOCK CAPITALS** where appropriate.

Most of the questions can be answered by putting a cross in a box like this: ☐

When you have finished please check that you have answered all of the questions that are relevant to you, and then return the questionnaire in the envelope enclosed. You do not need a stamp. Please return the questionnaire in the next two weeks.

More details about this project are available in the information sheet enclosed. If you would like further information about this project please contact Rosie Lacey, survey co-ordinator, on 01782 584721.

**Thank you for your help with this research project**

**Please complete this section first.**

1. Are you:

Female..... ☐

Male..... ☐

Day      Month      Year

2. What is your date of birth?

/  /

(For example - if you were born on the 5th June 1936, this would be entered as 05/06/36)

Day      Month      Year

3. Please enter today's date:

/  /

SECTION F - YOUR VIEWS ABOUT WHAT SHOULD BE RESEARCHED

We are interested in finding out your views about what **type of research** you think we should be doing on **joint pain in older adults**. You might find it helpful to think about what services are missing, what could be improved, or what is important to you.

1. Which topics should be made priority for joint pain research?

- Please look at the list of topics below and select **THREE** which are the most important to you.
- Then please put a cross against the topic you think is the most important in the box for "1st most important".
- For the next most important of your three choices, please cross "2nd most important".
- And finally put a cross against your third choice under "3rd most important".

Topic	1st most important	2nd most important	3rd most important
Injections in the joint.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Reduced swelling.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Tablets.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Diet / weight loss.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Complementary therapy.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Keeping active.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Aids and adaptations.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Joint replacement.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Counselling.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Removal of fluid / debris.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Improved mobility.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Education and advice.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Physical therapy.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

(Please check that you have only crossed 3 boxes)

2. Other (Please write below topics not listed above that you think should be a priority for joint pain research)

.....

.....

.....

# Section A

## HAND

**Please complete this section if you have  
had hand problems at any time in the last year**





# Section F

## **YOUR VIEWS ABOUT WHAT SHOULD BE RESEARCHED**

This section is for everyone to complete

Section A  
HAND

Please complete this section if you have  
had hand problems at any time in the last year



29727

9. Do you think joint pain will limit your ability to work before you reach 68 years old??

*(Please put a cross in one box only)*

I am 68 or older and I am still working..... ☐

It won't limit me, and I will work until I am 68..... ☐

Joint pain will increasingly limit my ability to work, but I will be able to work until I am 68..... ☐

Joint pain will stop me from working before I am 68..... ☐

Don't know..... ☐

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION**

**PLEASE CONTINUE AND FILL IN SECTION F**

## **SECTION A - HAND PROBLEMS**

For this questionnaire 'hand problems' relate to **any hand symptoms** you may experience; for example, **pain or stiffness**, or any **knobbly swellings** on your hands, including your fingers and thumbs.

### **Part 1**

1. Have you had any **problems** with your **hands**, including your fingers and thumbs, over the **last year**? *(Please put a cross in one box only)*

Yes..... ☐

No..... ☐

2. Have you had **hand pain**, including your fingers and thumbs, over the **last year**? *(Please put a cross in one box only)*

Yes..... ☐ Please continue with **Question 3**, below.

No..... ☐ Please go to **Part 2**, on page 8.

3. Please think back over the **last 12 months**. Please put a cross in **one** box only to show if you have pain, or have had pain:

in both hands  
(including fingers  
and thumbs)

☐

the right hand only  
(including fingers  
and thumbs)

☐

the left hand only  
(including fingers  
and thumbs)

☐

4. Again, please think back over the **last 12 months**. On how many days have you had this **hand pain**? *(Please put a cross in one box only)*

Less than 7 days..... ☐

1 to 4 weeks..... ☐

More than 1 month but less than 3 months..... ☐

3 months or more..... ☐

## Part 2

We are interested to know whether you have any finger nodes. A finger node is a firm, knobby swelling on the back of a finger joint.

### For example:

A finger without nodes:




A finger with nodes:



Please look at your hands and on the diagrams below circle any finger **joints** on which you have nodes or swellings (you may have several nodes/swellings)

Left

Right



6. To what extent can you **change or adapt** your normal work situation, if your joint pain is really bad (e.g. take a break or do lighter duties)?  
(Please put a cross in one box only)

A great deal	To some extent	A little	Very little
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. Have you taken time off work during the **last 6 months** for any health problems **other** than **joint pain**?  
(Please put a cross in one box only)

Yes..... ☐ (Please answer the question below to give us more details)

No..... ☐ (Please go to Question 8)

If **yes**, please write in the **number** of months, weeks or days you were off work due to any health problems **other** than **joint pain** in the last 6 months.

<input type="text"/> <input type="text"/> months	<input type="text"/> <input type="text"/> weeks	<input type="text"/> <input type="text"/> days
--	---	--

8. Do you think your joint pain will limit your ability to work before you reach retirement age (65 years for men, 60 years for women)?  
(Please put a cross in one box only)

I am older than retirement age and I am still working..... ☐

No, it won't limit me and will not cause me to retire early..... ☐

Joint pain will limit me, but will not cause me to retire early..... ☐

Joint pain will limit me and cause me to retire early..... ☐

4. We would like to know how joint pain affects your work. Please **put a cross in only ONE box** by the **statement** that best describes your **joint pain**.

I can do my job without getting more joint pain..... ☐

I can do my job but it gives me more joint pain..... ☐

For more than a quarter of my usual working day,  
joint pain prevents me from doing my job..... ☐

For more than half of my usual working day, joint pain  
prevents me from doing my job..... ☐

Joint pain prevents me from working at all..... ☐

5. Please indicate how much you agree or disagree with each statement.  
*(Please put a cross in one box on each line)*

	Strongly disagree	Disagree	Uncertain	Agree	Strongly agree
a. Because of my joint pain, the stresses of my job are much harder to handle.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Despite having joint pain, I am able to finish hard tasks in my work.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. My joint pain distracts me from taking pleasure in my work.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
d. Due to my joint pain I feel hopeless about finishing certain work tasks.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
e. Despite my joint pain, I am able to concentrate on achieving my goals at work.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
f. Despite my joint pain, I feel energetic enough to complete all my work?.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 3**

1. The following questions concern the amount of pain you have experienced in your hands. For each situation please enter the amount of **pain** experienced **in the last week**.

*(Please put a cross in one box on each line)*

**QUESTION: How much pain do you have in your hands?**

a. At rest (i.e. when not using your hands).

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

b. When gripping objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

c. When lifting objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

d. When turning objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

e. When squeezing objects with your hands.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. The following question concerns the amount of **joint stiffness** (not pain) you have experienced **in the last week** in your **hands**.  
Stiffness is a sensation of restriction or slowness in the ease with which you move your hands.

*(Please put a cross in the box of the most appropriate answer)*

How **severe** is stiffness in your hands **after first wakening** in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. The following questions concern your **physical function**. By this we mean your ability to move around and to look after yourself.  
For each of the following activities, please indicate the **degree of difficulty** you have experienced **in the last week** due to your **hand problem**.

*(Please put a cross in one box on each line)*

**QUESTION: How much difficulty do you have with the following?**

- a. Turning taps on.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- b. Turning a round door-knob or handle.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

- c. Doing up buttons.

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

## SECTION E - WORK

1. We would like to know if your **current** job has been affected by joint pain.  
*(Please put a cross in one box on each line)*

	Yes	No
a. I have joint pain and the way I do my job and the amount of work I do has <b>not</b> been affected.....	<input type="checkbox"/>	<input type="checkbox"/>
b. I have <b>changed</b> the way I do my job due to joint pain.....	<input type="checkbox"/>	<input type="checkbox"/>
c. I have <b>reduced</b> the number of hours I work due to joint pain...	<input type="checkbox"/>	<input type="checkbox"/>
d. I am currently <b>off work</b> due to <b>joint pain</b> .....	<input type="checkbox"/>	<input type="checkbox"/>
e. I am currently <b>off work</b> due to a <b>health problem other than joint pain</b> .....	<input type="checkbox"/>	<input type="checkbox"/>

2. In the last **6 months**, how much has your joint pain **changed** your ability to work, where 0 is 'no change' and 10 is 'complete change'?  
*(Please circle one number only)*

No change											Complete change
0	1	2	3	4	5	6	7	8	9	10	

3. Have you taken time off work during the **last 6 months** because of your **joint pain**?

*(Please put a cross in one box only)*

Yes..... ☐ *(Please answer the question below to give us more details)*

No..... ☐ *(Please go to Question 4)*

If **yes**, please write in the **number** of months, weeks or days you were off work due to your **joint pain** in the last 6 months.

<input type="text"/> <input type="text"/>	months	<input type="text"/> <input type="text"/>	weeks	<input type="text"/> <input type="text"/>	days
---	--------	---	-------	---	------

**QUESTION: How much difficulty do you have with the following?**

***(Please put a cross in one box on each line)***

d. Fastening jewellery (e.g. watches, earrings, cufflinks, necklaces, brooches and bracelets).

None

☐

Mild

☐

Moderate

☐

Severe

☐

Extreme

☐

e. Opening a new jar.

None

☐

Mild

☐

Moderate

☐

Severe

☐

Extreme

☐

f. Carrying a full pot with one hand (e.g. carrying any reasonably heavy object such as a saucepan).

None

☐

Mild

☐

Moderate

☐

Severe

☐

Extreme

☐

g. Peeling vegetables/fruits.

None

☐

Mild

☐

Moderate

☐

Severe

☐

Extreme

☐

h. Picking up large heavy objects.

None

☐

Mild

☐

Moderate

☐

Severe

☐

Extreme

☐

i. Wringing out washcloths (e.g. squeezing a wet sponge or flannel).

None

☐

Mild

☐

Moderate

☐

Severe

☐

Extreme

☐

**Part 4**

These are some general questions about your hands.

*(Please put a cross in one box on each line)*

1. Do you consider yourself to be...

Right-handed.... ☐

Left-handed.... ☐

Both.... ☐

2. Considering all the ways that your hand problem affects you, how well are you doing compared to other people your age?

Very well

Well

Fair

Poor

Very poorly

☐☐☐☐☐

3. Have you consulted your G.P. (family doctor) in the **past 12 months** because of your hand problem?

*(Please put a cross in one box only)*

Yes..... ☐

No..... ☐

# Section E

## WORK

Please complete this section **ONLY** if you are **currently** working in full-time or part-time paid work, or you have been off work due to ill health for more than 1 week but **less** than 6 months.



**Because of pain in my feet:**

15. My feet are worse in the morning

None of the time                      On some days                      On most / every day(s)

☐    ☐    ☐

16. My feet are more painful in the evening

None of the time                      On some days                      On most / every day(s)

☐    ☐    ☐

17. I get shooting pains in my feet

None of the time                      On some days                      On most / every day(s)

☐    ☐    ☐

18. I am unable to carry out my previous work

None of the time                      On some days                      On most / every day(s)                      Not applicable

☐    ☐    ☐    ☐

19. I no longer do all my previous activities (sport, dancing, hill-walking, etc.)

None of the time                      On some days                      On most / every day(s)                      Not applicable

☐    ☐    ☐    ☐

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION**

**PLEASE CONTINUE AND FILL IN SECTION E IF YOU ARE IN  
FULL-TIME OR PART-TIME PAID WORK OR YOU HAVE BEEN  
OFF WORK DUE TO ILL HEALTH FOR MORE THAN 1 WEEK BUT  
LESS THAN 6 MONTHS**

4. Which of the following services or treatments have you used in the **past 12 months** because of your hand problem? For **each service** you have used please put a cross to show whether the NHS provided this, or if you had private treatment. If you have used NHS **and** private services, please cross both boxes. For any service that you have not used, please leave blank.

	Yes (NHS)	Yes (Private)
a. Physiotherapy.....	<input type="checkbox"/>	<input type="checkbox"/>
b. Occupational therapy.....	<input type="checkbox"/>	<input type="checkbox"/>
c. Hospital specialist.....	<input type="checkbox"/>	<input type="checkbox"/>
d. Acupuncture.....	<input type="checkbox"/>	<input type="checkbox"/>
e. Osteopath/Chiropractor....	<input type="checkbox"/>	<input type="checkbox"/>
f. Drugs on prescription.....	<input type="checkbox"/>	<input type="checkbox"/>
g. Hand operation.....	<input type="checkbox"/>	<input type="checkbox"/>
h. Hand injection.....	<input type="checkbox"/>	<input type="checkbox"/>
i. Other (please state) .....		

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION**

**PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU (INCLUDING SECTION F ON PAGE 43)**



Because of pain in my feet:

7. I avoid standing for a long time	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. I catch the bus or use the car more often	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. I need help with housework / shopping	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. I still do everything but with more pain or discomfort	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11. I get irritable when my feet hurt	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12. I feel self-conscious about my feet	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
13. I get self-conscious about the shoes I have to wear	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
14. I have constant pain in my feet	None of the time	On some days	On most / every day(s)
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Below are some statements about problems people have because of **pain in their feet**.

For each statement indicate if this has applied to you during **the past month**. If so, was this only on some days or on most or every day in the past month?

*(Please put a cross in one box on each line)*

**Because of pain in my feet:**

1. I avoid walking outside at all

None of the time

☐

On some days

☐

On most / every day(s)

☐

2. I avoid walking long distances

None of the time

☐

On some days

☐

On most / every day(s)

☐

3. I don't walk in a normal way

None of the time

☐

On some days

☐

On most / every day(s)

☐

4. I walk slowly

None of the time

☐

On some days

☐

On most / every day(s)

☐

5. I have to stop and rest my feet

None of the time

☐

On some days

☐

On most / every day(s)

☐

6. I avoid hard or rough surfaces when possible

None of the time

☐

On some days

☐

On most / every day(s)

☐

# Section B

## HIP

**Please complete this section if you have had hip pain at any time in the last year**

## SECTION D - FOOT PAIN

These questions are about any **pain** you may have experienced in your **FEET**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the foot?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both feet... ☐ in the right foot only... ☐ in the left foot only... ☐

3. Again please think back over the **last 12 months**. On how many days have you had this **foot pain**?  
(Please put a cross in one box only)

Less than 7 days..... ☐

1 to 4 weeks..... ☐

More than 1 month but less than 3 months..... ☐

3 months or more..... ☐

4. Have you consulted your GP (family doctor) in the **last 12 months** because of your foot pain?  
(Please put a cross in one box only)

Yes..... ☐

No..... ☐

## **SECTION B - HIP PAIN**

These questions are about any **pain** you may have experienced in your **HIPS**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the hip?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both hips.. ☐ in the right hip only.. ☐ in the left hip only.. ☐

3. Again please think back over the **last 12 months**. On how many days have you had this **hip pain**?

**(Please put a cross in one box only)**

Less than 7 days..... ☐

1 to 4 weeks..... ☐

More than 1 month but less than 3 months..... ☐

3 months or more..... ☐

4. Have you consulted your GP (family doctor) in the **last 12 months** because of your hip pain?

**(Please put a cross in one box only)**

Yes..... ☐

No..... ☐

For all of the following questions please think back over the last 48 hours.

### Part 1

The following questions concern the amount of **pain** you have experienced in your hip. For each question please enter the amount of pain experienced in the **last 48 hours**.

*(Please put a cross in one box on each line)*

**Question: How much pain do you have?**

1. Walking on a flat surface

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Going up or down stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. At night while in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Sitting or lying

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

5. Standing upright

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

# Section D

# FOOT

**Please complete this section if you have had foot pain at any time in the last year**

**Part 2**

The following questions concern the amount of **joint stiffness** (not pain) you have experienced in the **last 48 hours** in your hip. Stiffness is a sensation of restriction or slowness in the ease with which you move your joints.  
*(Please put a cross in one box on each line)*

6. How **severe** is your stiffness after **first waking** in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. How **severe** is your stiffness after sitting, lying or resting **later in the day**?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Part 3**

The following questions concern your **physical function**. By this we mean your ability to move around and look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the **last 48 hours** due to your **hip**.  
*(Please put a cross in one box on each line)*

**Question: What degree of difficulty do you have?**

8. Descending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. Ascending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Rising from sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>



Question: What degree of difficulty do you have?

11. Standing

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

12. Bending to the floor

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

13. Walking on flat

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

14. Getting in/out of car

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

15. Going shopping

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

16. Putting on socks/stockings

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

17. Rising from bed

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

18. Taking off socks/stockings

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

Question: What degree of difficulty do you have?

19. Lying in bed

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

20. Getting in/out of bath

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

21. Sitting

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

22. Getting on/off toilet

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

23. Heavy domestic duties

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

24. Light domestic duties

None Mild Moderate Severe Extreme  
☐ ☐ ☐ ☐ ☐

© Dr Nicholas Bellamy. All rights reserved 1996

THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION

PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU (INCLUDING SECTION F ON PAGE 43)

© Dr Nicholas Bellamy. All rights reserved 1996



**Question: What degree of difficulty do you have?**

11. Standing

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

12. Bending to the floor

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13. Walking on flat

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

14. Getting in/out of car

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

15. Going shopping

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

16. Putting on socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

17. Rising from bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

18. Taking off socks/stockings

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Question: What degree of difficulty do you have?**

19. Lying in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

20. Getting in/out of bath

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

21. Sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

22. Getting on/off toilet

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

23. Heavy domestic duties

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

24. Light domestic duties

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

© Dr Nicholas Bellamy. All rights reserved 1996

**THANK YOU VERY MUCH FOR YOUR TIME AND HELP IN  
COMPLETING THIS SECTION**

**PLEASE CONTINUE AND FILL IN THE OTHER SECTIONS  
RELEVANT TO YOU (INCLUDING SECTION F ON PAGE 43)**



29727

© Dr Nicholas Bellamy. All rights reserved 1996



Part 2

The following questions concern the amount of **joint stiffness** (not pain) you have experienced in the **last 48 hours** in your knee. Stiffness is a sensation of restriction or slowness in the ease with which you move your joints.  
*(Please put a cross in one box on each line)*

6. How **severe** is your stiffness after **first wakening** in the morning?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

7. How **severe** is your stiffness after sitting, lying or resting **later in the day**?

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Part 3

The following questions concern your **physical function**. By this we mean your ability to move around and look after yourself. For each of the following activities, please indicate the degree of difficulty you have experienced in the **last 48 hours** due to your knee.  
*(Please put a cross in one box on each line)*

Question: What degree of difficulty do you have?

8. Descending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

9. Ascending stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

10. Rising from sitting

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

For all of the following questions please think back over the last 48 hours.

### Part 1

The following questions concern the amount of **pain** you have experienced in your knee. For each question please enter the amount of pain experienced in the **last 48 hours**.

*(Please put a cross in one box on each line)*

**Question: How much pain do you have?**

1. Walking on a flat surface

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

2. Going up or down stairs

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. At night while in bed

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

4. Sitting or lying

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>


5. Standing upright

None	Mild	Moderate	Severe	Extreme
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

# Section C

## KNEE

**Please complete this section if you have had knee pain at any time in the last year**



29727

### **SECTION C - KNEE PAIN**

These questions are about any **pain** you may have experienced in your **KNEES**.

Please answer all the questions in this section.

1. Have you had pain in the **last year** in or around the knee?

Yes..... ☐

No..... ☐

2. Please think back over the **last 12 months**. Please put a cross in one box to show if you have pain, or have had pain:

in both knees.. ☐ in the right knee only.. ☐ in the left knee only.. ☐

3. Again please think back over the **last 12 months**. On how many days have you had this **knee pain**?

*(Please put a cross in one box only)*

Less than 7 days..... ☐

1 to 4 weeks..... ☐

More than 1 month but less than 3 months..... ☐

3 months or more..... ☐

4. Have you consulted your GP (family doctor) in the **last 12 months** because of your knee pain?

*(Please put a cross in one box only)*

Yes..... ☐

No..... ☐

## **Appendix 7: QUIPS-LBP Tool**

## QUality In Prognosis Studies (QUIPS) – Low Back Pain: Quality Appraisal Exercise

**Primary Goal:** Generate a list of specific issues related to potential biases in LBP prognosis studies

**Steps (for each assigned study):**

- Review the methods of the assigned prognosis study
- Decide if the study methods satisfy each general statement below for the 6 biases
- Explain **WHY** you chose that rating under ‘*Comments*’ (good or bad methods achieved)
- Provide explanation (important points or exceptions) relevant to assess LBP studies
- Citations to methods publications are encouraged and will be extremely helpful

Study (First author, year): \_\_\_\_\_

<b>Bias related to Study Participation</b>	
<b>1.</b> The source population or population of interest is adequately described for key characteristics.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i> For example, what are the 'key characteristics' that need to be described for the source population?
<b>2.</b> The sampling frame and recruitment are adequately described, possibly including: <ul style="list-style-type: none"> <li>- Methods to identify the sample (number and type used, e.g. referral patterns in healthcare settings)</li> <li>- Time period of recruitment</li> <li>- Place of recruitment (setting and geographic location)</li> </ul> <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>3.</b> Inclusion and exclusion criteria are adequately described, including explicit diagnostic criteria or 'zero time' description.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>4.</b> There is adequate participation in the study by eligible individuals.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>5.</b> The baseline study sample (i.e. individuals entering into the study) is adequately described for key characteristics.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>6. Study Participation Summary:</b> Is the following statement satisfied based on responses to the above questions, "The study sample represents the population of interest on key characteristics, sufficient to limit potential bias to the results"?  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear	<i>Comments:</i> Which of the above questions did you consider to make this judgment?

<b>Bias related to Study Attrition</b>	
<p><b>7.</b> Response rate (i.e. proportion of study sample completing the study and providing outcome data) is adequate.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>8.</b> Attempts to collect information on drop-outs are described.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>9.</b> Reasons for 'loss to follow-up' are provided.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>10.</b> Subjects lost to follow-up are adequately described for key characteristics.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>11.</b> There are no important differences between completers and non-completers on key characteristics and outcomes.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>12. Study Attrition Summary:</b>  The following statement is satisfied based on responses to the above questions, "Loss to follow-up (from sample to study population) is not associated with key characteristics (i.e. the study data adequately represents the sample), sufficient to limit potential bias"?</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear</p>	<i>Comments:</i>

<b>Bias related to Prognostic Factor Measurement</b>	
<p><b>13.</b> A clear definition or description of the prognostic factor measured is provided (e.g. including dose, level, duration of exposure, and clear specification of the method of measurement).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>14.</b> Continuous variables are reported or appropriate (i.e. not data dependent) cut-points are used.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>15.</b> The prognostic factor measure and method used is adequately valid and reliable to limit misclassification bias (e.g. may include relevant outside sources of information on measurement properties, also characteristics such as blind measurement, limited reliance on recall).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>16.</b> Adequate proportion of sample has complete data for all relevant outcome periods.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>17.</b> The method and setting of measurement is the same for all study participants.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>18.</b> Appropriate methods are used if imputation is used for missing prognostic factor data.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>19. Prognostic Factor Measurement Summary:</b>  The following statement is satisfied based on responses to the above questions, "The prognostic factor of interest is adequately measured in study participants to sufficiently limit potential bias"?</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear</p>	<i>Comments:</i>

<b>Bias related to Outcome Measurement</b>	
<p><b>20.</b> A clear definition of the outcome of interest is provided including duration of follow-up, the level and extent of the outcome construct.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<p><i>Comments:</i></p>
<p><b>21.</b> The outcome measure and method used is adequately valid and reliable to limit misclassification bias (e.g. may include relevant outside sources of information on measurement properties, also characteristics such as blind measurement, confirmation of outcome with valid and reliable test).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<p><i>Comments:</i></p>
<p><b>22.</b> The method and setting of measurement is the same for all study participants.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<p><i>Comments:</i></p>
<p><b>23. Outcome Measurement Summary:</b> The following statement is satisfied based on responses to the above questions, "The outcome of interest is adequately measured in study participants to sufficiently limit potential bias"?</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear</p>	<p><i>Comments:</i></p>



<b>Bias related to Confounding Measurement and Account</b>	
<p><b>24.</b> All important confounders, including treatments (key variables in conceptual model) are measured.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>25.</b> Clear definitions of the important confounders measured are provided (e.g. including dose, level, and duration of exposures).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>26.</b> Measurement of all important confounders is adequately valid and reliable (e.g. may include relevant outside sources of information on measurement properties, also characteristics such as blind measurement, limited reliance on recall).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>27.</b> The method and setting of confounding measurement is the same for all study participants.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>28.</b> Appropriate methods are used if imputation is used for missing confounder data.</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>29.</b> Important potential confounders are accounted for in the study design (i.e. matching for key variables, stratification, or initial assembly of comparable groups).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>30.</b> Important potential confounders are accounted for in the analysis (i.e. appropriate adjustment).</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear   <input type="checkbox"/> Not relevant</p>	<i>Comments:</i>
<p><b>31. Confounding Measurement and Account Summary:</b>  The following statement is satisfied based on responses to the above questions, "Important potential confounders are appropriately accounted for, limiting potential bias with respect to the prognostic factor of interest"?</p> <p><input type="checkbox"/> Yes   <input type="checkbox"/> Partly   <input type="checkbox"/> No   <input type="checkbox"/> Unclear</p>	<i>Comments:</i>

<b>Bias related to Analysis</b>	
<b>32.</b> There is sufficient presentation of data to assess the adequacy of the analysis.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>33.</b> The strategy for model building (i.e. inclusion of variables) is appropriate and based on a conceptual framework or model.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>34.</b> The selected model is adequate for the design of the study.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>35.</b> There is no selective reporting of results.  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/> Not relevant	<i>Comments:</i>
<b>36. Analysis Summary:</b> The following statement is satisfied based on the responses to the above questions, "The statistical analysis is appropriate for the design of the study, limiting potential for presentation of invalid results"?  <input type="checkbox"/> Yes <input type="checkbox"/> Partly <input type="checkbox"/> No <input type="checkbox"/> Unclear	<i>Comments:</i>

Overall comments on methods in LBP prognosis studies: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Comments on this project: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**THANK YOU!**

## **Appendix 8: Published conference abstracts**

# Medical or Research Professionals / Clinicians - Abstract Submission

*Epidemiology, health services and outcome research*

EULAR11-5561

## SELF-CARE NEEDS IN COMMUNITY-DWELLING OLDER ADULTS WITH JOINT PAIN: A DESCRIPTIVE CROSS-SECTIONAL STUDY

Y. Greenhill<sup>1,\*</sup>, R. Wilkie<sup>1</sup>, R. Beech<sup>1</sup>, G. Peat<sup>1</sup>

<sup>1</sup>Arthritis Research UK Primary Care Centre, KEELE UNIVERSITY, Staffordshire, United Kingdom

**I confirm that my abstract complies with the rules and contains original data:** Yes

**My abstract has been or will be presented at a scientific meeting during a 12 months period prior to EULAR 2011:**

No

**Are you applying for a Travel Bursary?:** Yes

**Background:** Self-care restriction is a frequent form of disability in older adults and is associated with joint pain. The need for personal help and assistance and its effects on self-care are under-researched.

**Objectives:** To describe the extent of the need for and receipt of help and assistance in older adults with joint pain and explore the relationship with meeting self-care needs “as and when I want”.

**Methods:** Cross-sectional analysis of data from a population-based cohort study of joint pain in adults aged 50 years and over: the North Staffordshire Osteoarthritis Project. Eligible participants responded to a postal survey, reported hand, hip, knee or foot pain in the previous 12 months and difficulties with washing and bathing. Personal help and assistance for self-care was measured by a single item with responses classed as (i) no expressed need for help/ assistance, (ii) expressed need but non-receipt of help and (iii) expressed need and receipt of help. Using an item from the Keele Assessment of Participation (KAP), all participants were asked how often their self-care needs were met “as and when they wanted”. Frequency of expressed need for and receipt of, personal help/assistance were described and related to the KAP responses using binary logistic regression. Results are expressed as crude odds ratios (OR) with 95% confidence intervals and after adjustment for severity of difficulties with washing and bathing limitation and covariates independently associated with self-care restriction in the general population.

**Results:** Of 588 older adults who were included in the analysis (mean age was 73 (SD=9.4); 61% were female). 279 (47%) expressed a need for help/assistance, of whom 175 received help from family or friends (n=157) or health professionals (n=18). Those expressing a need for help but not receiving any were most likely to feel their self-care needs were seldom met “as and when they wanted” (crude OR 6.3; 95% CI: 3.7, 10.7) (Table1). Of 175 receiving help, 68 (39%) still perceived their self-care needs to be not fully met. This pattern of associations persisted after adjusting for the severity of difficulties and for selected covariates.

Table1. Associations between perceived self-care needs not being met “as and when wanted” and expressed need for and receipt of, personal help or assistance for self-care in 588 community-dwelling older adults with joint pain and difficulties in self-care.

	KAP item: Self-care needs seldom met “as and when wanted”		
	n (%)	OR (95%CI)	aOR (95%CI)*
Expressed need and receipt of help/assistance			
No expressed need (n=309)	35 (12)	1	1
Expressed need-non-receipt of help (n=104)	45 (50)	6.3 (3.7, 10.7)	8.8 (4.5, 17.3)
Expressed need-receipt of help (n=175)	68 (39)	4.9 (3.7, 10.7)	5.4 (2.9, 10.2)

**Image/Graph:**

**Conclusions:** Among older adults with joint pain and difficulties in self-care tasks, there is a high level of expressed need for personal help/assistance which, when received, is provided largely by informal networks of friends and family rather than formal health or social care. The receipt of personal help/assistance contributes to having their self-care needs met “as and when they want” but remains insufficient in many cases. Additional provision of personal assistance may reduce self-care restriction, however further investigation into the reasons for self-care restriction is required to explain why it occurs when the need for personal help is met.

**Disclosure of Interest:** None Declared

**SAT0575**

**POPULATION BASED PROSPECTIVE STUDY OF SELF-CARE RESTRICTION IN OLDER ADULTS WITH JOINT PAIN**

Y. Prior<sup>1,2,\*</sup>, R. Wilkie<sup>1</sup>, R. Beech<sup>1</sup>, G. Peat<sup>1</sup>

<sup>1</sup>Arthritis Research UK Primary Care Sciences, Keele University, Newcastle-under-lyme, <sup>2</sup>Centre for Health Sciences Research, University of Salford, Manchester, United Kingdom

**Background:** Self-care restriction is a frequent form of disability in older adults and is associated with joint pain. The natural history of self-care restriction in older adults with joint pain is unknown.

**Objectives:** To (i) describe the frequency of onset and persistence of self-care restriction (ii) establish whether joint-specific characteristics are associated with onset and persistence (iii) identify potential predictors of onset and persistence of self-care restriction in community-dwelling older adults with joint pain.

**Methods:** Data was collected in a population-based prospective cohort study (baseline and 3-year follow-up) of joint pain in adults aged 50 years and over in the North Staffordshire Osteoarthritis Project (n=3147). Perceived self-care participation restriction was measured using one item from the Keele Assessment of Participation ("During the past 4 weeks my self-care needs (examples are washing, bathing, toileting, dressing, feeding and maintaining health) have been met, as and when I have wanted") at baseline and three year follow-up. A single item at baseline was used to classify chronicity. Severity of hand, hip, knee, and foot pain were assessed at baseline by single items and subscale scores from the AUSCAN (hand), WOMAC (hip/knee) and MFPDI (foot). Binary logistic regression was used to estimate the association between joint specific characteristics and onset and persistence of self-care restriction at 3 years, before and after adjusting for covariates previously shown to be associated with restricted self-care in the general population (age, activity limitation (SF-36 PF-10), depression (HAD), cognitive complaint (SIP), perceived adequacy of income and educational attainment). Results were summarised as odds ratios (OR) with 95% confidence intervals (CI).

**Results:** The frequencies of onset and persistence of self-care restriction at 3 years were 7.5% (95%CI: 6.6, 8.6) and 39.7% (34.2, 45.4) respectively. Chronic joint pain, severe pain and stiffness were associated with onset and persistence of self-care restriction at 3 years (i.e. for onset, severe vs. non-severe pain: crude OR 3.0 (2.1, 4.0), chronic vs. non-chronic pain (<3 months): 1.9 (1.3, 2.8), severe vs. non-severe stiffness: 2.7 (1.9, 3.7); for persistence, 2.0 (1.0, 3.8); 1.4 (1.0, 2.8); 1.7 (1.0, 2.8) respectively). For all joint pain factors, adjustment for the factors associated with self-care restriction in the general population attenuated all associations. Being aged 80 years and over, lower educational attainment, higher levels of activity limitation, depression and cognitive impairment were independently associated with onset. Only depression was associated with persistence of self-care restriction (adj. OR 2.4 (1.1, 5.4)) after adjustment for all other covariates.

**Conclusions:** Perceived self-care restriction, as assessed by the KAP, changes over time. Older adults with chronic and severe peripheral joint pain have an increased risk of developing self-care restriction and of this persisting. In a multivariable model these joint-specific characteristics were found not to be independent predictors, suggesting that, in the absence of significant bias or confounding, the effect of joint pain on self-care restriction may be through its effects on activity limitation, depression, and cognitive complaint.

**Disclosure of Interest:** None Declared